

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

A pilot evaluation of the Sleep Ninja – a smartphone-application for adolescent insomnia symptoms

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-026502
Article Type:	Research
Date Submitted by the Author:	10-Sep-2018
Complete List of Authors:	Werner-Seidler, Aliza ; University of New South Wales, Black Dog Institute Wong, Quincy; University of New South Wales, Black Dog Institute; Western Sydney University School of Science and Health, School of Social Sciences and Psychology Johnston, Lara; University of New South Wales, Black Dog Institute O'Dea, B; University of New South Wales, Black Dog Institute Torok, Michelle; University of New South Wales, Black Dog Institute Christensen, Helen; University of New South Wales - Randwick Campus, Black Dog Institute
Keywords:	insomnia, adolescent mental health, cognitive-behaviour therapy for insomnia, ehealth

SCHOLARONE™
Manuscripts

1
2
3
4
5 **A pilot evaluation of the Sleep Ninja – a smartphone-application for adolescent**
6
7 **insomnia symptoms**
8
9
10
11
12
13
14
15

16 Aliza Werner-Seidler¹; a.werner-seidler@blackdog.org.au

17 Quincy Wong^{1,2}; quincy.wong@blackdog.org.au

18 Lara Johnston¹; l.johnston@blackdog.org.au

19 Bridianne O’Dea¹; b.odea@blackdog.org.au

20 Michelle Torok¹; m.torok@blackdog.org.au

21 Helen Christensen¹; h.christensen@blackdog.org.au
22
23
24
25
26
27
28
29
30
31
32

33 Affiliations:

34 ¹Black Dog Institute, University of New South Wales, Sydney, Australia

35 ²School of Social Sciences and Psychology, Western Sydney University, Sydney, NSW,
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Corresponding author: Dr Aliza Werner-Seidler; Black Dog Institute, Hospital Road,
Randwick, New South Wales, Australia; a.werner-seidler@blackdog.org.au; 02 9382 380.

Word Count: 5102

Keywords: Insomnia, adolescent mental health, cognitive-behaviour therapy for insomnia,
eHealth

Abstract

Objectives: The aim of this study was to test the feasibility, acceptability and preliminary effects of a recently developed smartphone application, Sleep Ninja, for adolescent sleep difficulties.

Setting: The study was conducted online with Australian individuals recruited through the community.

Participants: Participants were 50 young people aged 12-16 years with at least mild symptoms of insomnia.

Design: A single-arm pre-post design was used to evaluate feasibility, acceptability and sleep and mental health variables at baseline and post-intervention.

Intervention: Cognitive Behaviour Therapy for Insomnia (CBT-I) informed the development of the Sleep Ninja. The core strategies covered by the app are psychoeducation, stimulus-control, sleep hygiene, cognitive therapy and relaxation techniques. It includes six training sessions (lessons), a sleep tracking function, recommended bedtimes based on sleep guidelines, reminders to start a wind-down routine each night, a series of sleep tips, and general information about sleep. Users progress through each training session and conclude the six-week program with a black belt in sleep.

Outcome measures: Feasibility was evaluated based on consent rates, adherence and attrition, acceptability was assessed using questionnaires and a post-study interview, and sleep, depression and anxiety variables were assessed at baseline and post-intervention.

Results: Data indicated that the Sleep Ninja is a feasible intervention and is acceptable to young people. Findings showed there were significant improvements on sleep variables including insomnia (within-group effect size $d=-0.90$), sleep quality ($d=-0.46$), depression ($d=-0.36$) and anxiety ($d=-0.41$).

1
2
3 **Conclusions:** The Sleep Ninja is a promising intervention for the treatment of sleep
4 difficulties among adolescents. A follow-up randomised controlled trial is now warranted.
5

6
7 **Trial registration:** Australian New Zealand Clinical Trials Registry

8
9 (#ACTRN12617000141347).
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Summary

Strengths and limitations of this study

- This is the first study to evaluate app-delivered Cognitive Behavioural Therapy for Insomnia in adolescents with sleep difficulties.
- The intervention being tested, Sleep Ninja, was developed with input from young people, is fully automated and does not require internet coverage to function.
- The evaluation included measures of feasibility and acceptability as well as detailed semi-structured interviews about participants' experience with the app.
- As a preliminary study, this study did not include a control group.

1
2
3 Clinical insomnia is a sleep disturbance characterised by difficulty falling asleep,
4 staying asleep or waking up too early, with associated daytime impairment (American
5 Psychiatric Association, 2013). It effects approximately 4% of adolescents (Ohayon, Roberts,
6 Zulley, Smirne, & Priest, 2000), however sub-threshold symptoms are very common, with
7 approximately 25% of young people reporting some degree of sleep disturbance (Carskadon,
8 1990; Ohayon et al., 2000).

9
10
11
12
13
14
15
16 Depression and insomnia are closely linked, with comorbidity levels as high as 73%
17 in young people (Liu et al., 2007). Insomnia is not only a symptom of depression, but is a
18 common precursor, with high quality longitudinal data having established insomnia is an
19 independent risk factor for depression onset (Baglioni et al., 2011; Franzen & Buysse, 2008;
20 Riemann & Voderholzer, 2003). For example, a recent meta-analysis found that insomnia
21 was associated with a greater than two-fold increase in depression risk (Baglioni et al., 2011).
22 Depression is now the leading cause of disability in young people worldwide (Gore et al.,
23 2011), and approximately 1 in 5 young people will experience an episode of depression by
24 the time they are 18 years old. The demand for mental health services dramatically outweighs
25 the availability of these services, and current treatments alleviate only 13% of the disease
26 burden (Andrews, Issakidis, Sanderson, Corry, & Lapsley, 2004). Innovative and effective
27 ways to address the depression crisis in young people are needed, and one way to do this may
28 be by targeting sleep disturbance.

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44 In line with this suggestion, there is an emerging literature showing that targeting
45 insomnia in individuals with concurrent insomnia and depression improves both sleep and
46 depression outcomes (Ashworth et al., 2015; Manber et al., 2008; Wagley, Rybarczyk, Nay,
47 Danish, & Lund, 2013). This suggests there may be value in targeting sleep to improve
48 insomnia symptoms, with potential downstream effects on depression. To our knowledge,
49 there have been only two studies, one in adults and one in adolescents, testing the hypothesis
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that treating insomnia can prevent depressive symptoms. In the adult study, insomnia
4
5 treatment led to a significant reduction in depression following the intervention and at 6-
6
7 month and 18-month follow-up, relative to an active control group (Batterham et al., 2017;
8
9 Christensen et al., 2016). In the youth study, a face-to-face insomnia intervention was
10
11 delivered to secondary school students who had insomnia and anxiety. Results showed
12
13 improvements on sleep and anxiety outcomes, but not symptoms of depression (Blake et al.,
14
15 2016). Data from the two year follow-up from this study has not yet been published
16
17 (Waloszek et al., 2015).
18
19

20 The gold standard treatment for insomnia is cognitive behaviour therapy for insomnia
21
22 (CBT-I), recommended by major medical authorities (Australasian Sleep Association,
23
24 American College of Physicians), and typically involves a combination of strategies that
25
26 include psychoeducation, stimulus control, sleep restriction, cognitive therapy, and sleep
27
28 hygiene. The effectiveness of CBT-I is established among adolescents and adults, and there is
29
30 evidence from the adult literature to support the use of digitally delivered CBT-I to treat
31
32 insomnia symptoms (Okajima, Komada, & Inoue, 2011; Zachariae, Lyby, Ritterband, &
33
34 O'Toole, 2016). Although in its infancy, there are now two studies (one pilot and one
35
36 randomised controlled trial [RCT]) that have examined the use of internet-delivered CBT-I
37
38 for adolescents with subthreshold insomnia symptoms and insomnia, respectively, both with
39
40 positive results (de Bruin, Bogels, Oort, & Meijer, 2015; de Bruin, Oort, Bogels, & Meijer,
41
42 2014). However, there are no digitally delivered CBT-I programs that are commercially
43
44 available for youth at present (Werner-Seidler, Johnston, & Christensen, 2018).
45
46
47

48 Delivering sleep interventions via digital formats may be particularly well-suited to
49
50 young people. Today's adolescents have grown up in the digital age and are more
51
52 comfortable with technology than any previous generation. The move from face-to-face
53
54 therapies to web-based platforms has been positive — young people overwhelmingly (97%)
55
56
57
58
59
60

1
2
3 show a preference for digital delivery when given the choice between face-to-face and digital
4 CBT-I (de Bruin et al., 2014). This preference may in part be explained by the fact that young
5 people are notoriously reluctant to seek help for psychological issues, for reasons that include
6 stigma and a preference to manage the problem themselves (Gulliver, Griffiths, &
7 Christensen, 2010). Given that fewer than 40% of youth with a mental health problem access
8 professional help (Rickwood, Deane, & Wilson, 2007), focusing on factors such as sleep
9 disturbance as a risk factor for other more stigmatised disorders like depression, offers a
10 potentially more acceptable approach specifically for adolescents.

11
12 To meet the need for a widely available adolescent CBT-I program, we used a
13 participatory design process to develop an automated smartphone application (app; Sleep
14 Ninja) to target symptoms of adolescent sleep problems (Werner-Seidler et al., 2017). Using
15 smartphones to deliver CBT-I offers unprecedented convenience, removing barriers such as
16 cost and accessibility. Smartphone ownership levels are now at an all-time high (estimated to
17 be >80%), with approximately 95% of teenagers in English-speaking, developed countries
18 (Australia, Britain, US) having access to a smartphone (Ofcom, 2017; Pew Research Center,
19 2018; Triton Digital, 2017).

20
21 The goal of this pilot study was to examine the acceptability, feasibility and
22 preliminary effects of this intervention delivered to adolescents. In line with the guidelines on
23 the development of behavioural interventions (Craig et al., 2008; Gitlin, 2013), the primary
24 purpose of this study was to investigate recruitment rates, uptake, intervention completion,
25 reasons for non-adherence, and participant retention. The secondary aim was to use both
26 quantitative and qualitative methods to determine the acceptability of the app among young
27 people with sleep problems and allow for the refinement of the intervention prior to a formal
28 randomised evaluation. A final aim was to examine the impact of the Sleep Ninja app on
29 sleep outcomes and mental health symptoms. We used a single-arm, pre-post design to
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 address these objectives. It was hypothesised that the app would be a feasible modality in
4
5 which to deliver the automated sleep intervention, as measured by uptake, completion and
6
7 retention rates, that the app would be acceptable to young people, and that its use would be
8
9 associated with improvement in sleep and mental health symptoms.
10

11 12 13 **Method**

14
15 This trial was prospectively registered on the Australian New Zealand Clinical Trials
16
17 Registry (#ACTRN12617000141347).
18

19 20 **Participants**

21
22 Fifty participants were recruited via media and social media channels, including the
23
24 Black Dog Institute's website and paid Facebook advertisements that targeted potential
25
26 participants and their parents. Inclusion criteria were: aged 12-16 years, presence of at least
27
28 mild insomnia, operationalised by endorsement of at least one of the following symptoms
29
30 over the preceding two-week period: difficulty falling asleep, difficulty staying asleep or
31
32 problems waking up too early. These items are the first three questions on the Insomnia
33
34 Severity Index (Bastien, Vallieres, & Morin, 2001), and were chosen to include a participant
35
36 group with at least mild levels of insomnia. For study inclusion, participants also needed to
37
38 own a smartphone running iOS or Android, have a valid email address, access to the internet,
39
40 and be able to provide personal and parental consent.
41
42
43

44 45 **Measures**

46
47 **Insomnia Severity Index (ISI).** The ISI is a psychometrically sound, seven-item self-
48
49 report measure of insomnia symptoms over the previous two weeks (Bastien et al., 2001;
50
51 Chung, Kan, & Yeung, 2011). Responses are reported on a Likert scale from 0 to 4,
52
53 producing total scores of 0 to 28 (Bastien et al., 2001; Chung et al., 2011). Cut-off scores are
54
55 as follows: 0-7 reflects no clinically significant insomnia, 8-14 indicates subthreshold
56
57
58
59
60

1
2
3 insomnia, 15-21 suggests moderate severity insomnia, and 22-28 indicates severe insomnia
4
5 (Chahoud, Chahine, Salameh, & Sauleau, 2017).
6

7 **Pittsburgh Sleep Quality Index (PSQI).** The PSQI is a widely used self-report 19-
8
9 item scale that assesses usual sleep habits and experiences over the preceding month and has
10
11 been validated in adolescent samples (de la Vega et al., 2015). There are seven sub-scales
12
13 which are sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep
14
15 disturbances, use of sleeping medications, daytime dysfunction (Cole et al., 2006). Each
16
17 component is scored from 0 (no difficulty) to 3 (severe difficulty), which are summed to
18
19 obtain a Global PSQI score ranging from 0 to 21 (Guo, Sun, Liu, & Wu, 2016).
20
21

22 **Patient Health Questionnaire (PHQ-8).** The PHQ-8 assesses depressive symptoms
23
24 in the preceding two weeks (Kroenke et al., 2009). The questions are identical to those asked
25
26 in the PHQ-9 with the exclusion of the last item which asks about suicide. The PHQ-8 is
27
28 comparable to the PHQ-9 in terms of diagnosing depressive disorders (Kroenke, Spitzer, &
29
30 Williams, 2001). Each item is scored on a 4-point scale and summed together to form a total
31
32 depression score ranging from 0 to 24. Scores correspond to the following cut-offs: 0-9
33
34 indicates minimal symptoms, 10-14 indicates mild symptoms, 15-19 reflects moderate
35
36 symptoms, and 20-24 is indicative of severe depression. The PHQ-8 has demonstrated good
37
38 sensitivity (70%) and high specificity (98%) for major depression for scores ≥ 10 (Kroenke et
39
40 al., 2009).
41
42

43 **Generalised Anxiety Disorder 7-item (GAD-7).** The GAD-7 evaluates symptoms of
44
45 generalised anxiety disorder (Spitzer, Kroenke, Williams, & Lowe, 2006). All items are
46
47 scored on a scale from 0 (not at all) to 3 (nearly every day). The scores on each item are
48
49 summed together to derive a total score, ranging from 0 to 21 of which 0-4 indicates minimal
50
51 anxiety, 5-9 mild anxiety, 10-14 moderate anxiety, and 15-21 severe anxiety.
52
53
54
55
56
57
58
59
60

1
2
3 **Expectations of Success.** A four-item scale was developed for this study to assess
4 participants' motivation and expectations for improving their sleep with an app (e.g., *I am*
5 *confident that people could learn skills for improving sleep from an app*). The Expectation of
6 Success measure was scored on a five-point scale and total scores were computed by
7 summing each item, ranging from 0 to 16. Higher scores on this scale indicate greater
8 confidence and readiness to target sleep using a smartphone app.
9

10
11
12 **Acceptability of the Intervention.** The Acceptability of the Intervention scale is a
13 seven-item measure that was developed by the research team to assess participants' attitudes
14 and behaviours associated with using the app (e.g., *How much did you learn from the app and*
15 *Would you recommend this program to others?*). This measure was informed by similar
16 acceptability measures commonly used in the field (Thorndike et al., 2008). Each item was
17 designed to assess a different domain (such as app usefulness, ease of use, comprehension),
18 and therefore item scores were considered separately.
19

20
21
22 **Reasons for Non-Adherence.** The Reasons for Non-Adherence measure is a 23-item
23 scale that was developed by the researchers to assess the degree to which different reasons
24 impacted on participants' use of the app. Examples of domains assessed included technical
25 problems with the app and difficulty with the material. The scores on each item were
26 considered separately.
27

28
29
30 **Sleep Diary.** The ten-item Sleep Diary was developed by the research team
31 incorporating the questions from the Consensus Sleep Diary (Carney et al., 2012), with the
32 addition of two questions regarding daytime naps and use of sleep medication. Participants
33 answered 10 questions which included bedtime, time taken to fall asleep (sleep onset latency;
34 SOL), number and duration of night-time awakenings (number of awakenings; NWAK,
35 duration of wakefulness after sleep onset; WASO), time of final awakening, time participants
36 got out of bed for the day, subjective sleep quality, how refreshed participants felt upon
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 awakening, duration of any daytime naps and use of sleep medication. From the sleep diary
4 we calculated time in bed (TIB), total sleep time (TST; calculated by subtracting SOL,
5 WASO and time between waking and getting up in the morning, from TIB) and sleep
6 efficiency (SE; calculated by taking the percentage of TST/TIB).
7
8
9

10
11 **Post-Study Interview.** After study completion, participants were invited to attend a
12 face-to-face or telephone interview to provide feedback on their experience. Interviews were
13 semi-structured and explored participants' opinions about the study in general, and
14 specifically in relation to the intervention. Questions were open-ended, and flexible enough to
15 explore ideas that were raised during each interview. Interviews were audio recorded and
16 then transcribed verbatim by the interviewer. The interview content was pragmatically coded
17 into relevant themes by the same researcher, with oversight and guidance provided by the
18 research team.
19
20
21
22
23
24
25
26
27

28 **Intervention – ‘Sleep Ninja’**

29
30
31 The Sleep Ninja app was derived from CBT-I and developed by our team, as a fully-
32 automated smartphone app. Young people contributed to the content, functionality and
33 accessibility/user experience of the app through a series of focus groups (Werner-Seidler et
34 al., 2017). The core strategies included in the app were: psychoeducation, stimulus-control,
35 sleep hygiene, cognitive therapy and relaxation techniques. Sleep restriction was deliberately
36 omitted because some support (parental and/or professional) is likely to be required to
37 successfully implement sleep restriction (particularly in young people).
38
39
40
41
42
43
44
45

46 The structure of the Sleep Ninja app includes six training sessions (lessons), a sleep
47 tracking function, recommended bedtimes based on sleep guidelines, reminders to start a
48 wind-down routine each night, a series of sleep tips, and general information about sleep. The
49 home screen has 3 options: Train, Track, and More (see Figure 1). Users complete training
50 sessions which are delivered through a chat-bot format where the sleep ninja essentially acts
51
52
53
54
55
56
57
58
59
60

1
2
3 as a sleep coach. Training sessions take approximately 5-10 minutes to complete, and cover:
4
5 psychoeducation (training 1), stimulus control (training 2), sleep hygiene (training 3),
6
7 identifying and planning for high-risk situations (training 4), cognitive therapy (training 5),
8
9 and a final review session (training 6). The user interacts with the app through a forced
10
11 choice chat-bot format which is responsive to the input of the user, meaning it personalises
12
13 information and recommendations based on the selections and sleep profile recorded by the
14
15 participant. Users level up and reach their next “belt” by completing one training session and
16
17 tracking their sleep for 3 nights (out of a 7-night period). As there were six training sessions
18
19 to complete, the app was made available for six weeks (42 days) before it locked. Users finish
20
21 the program with a black belt in sleep.
22
23

24 **Procedure**

25
26 All procedures were approved by the University of New South Wales Human
27
28 Research Ethics Committee (HC#16702). Participants were encouraged to download consent
29
30 forms if they met the eligibility criteria listed on the study website and submit this directly to
31
32 the research team, once completed. Those who provided written informed consent and that of
33
34 a parent or guardian were then enrolled in the trial and invited to complete the screening
35
36 questions to verify study eligibility, before completing baseline questionnaires which
37
38 included: demographics, ISI, PSQI, PHQ-8, GAD-7 and Expectation of Success. Participants
39
40 could then access the first day of the online seven-day sleep diary. Another diary entry
41
42 became available each day for the following six days and participants were reminded to
43
44 complete entries via text-message. At the completion of seven entries in the diary,
45
46 participants were given access to the Sleep Ninja app on their personal smartphone devices.
47
48 Participants could use the app for six weeks before the post study questionnaire was made
49
50 available, which included the same battery as baseline with the omission of the Expectations
51
52 of Success questionnaire and with the addition of the Acceptability and Reasons for Non-
53
54
55
56
57
58
59

1
2
3 Adherence questionnaires. Participants then completed another seven-day sleep diary, which
4 was delivered in the same format and schedule as baseline. After the study had finished,
5 participants were invited to participate in a face-to-face or telephone interview to provide
6 feedback on their experience of participating in the study. Participants were reimbursed for
7 their time with giftcards to the value of \$10 each for completing baseline and post-study
8 assessment schedules; \$20 for a telephone interview and \$30 for a face-to-face interview.
9
10
11
12
13
14
15

16 **Patient and Public Involvement**

17
18 Prior to this study, a separate group of young people were consulted in a series of
19 focus groups to inform the design, features and structure of the app (for more information
20 please see Werner-Seidler et al., 2017). As a feasibility and acceptability study, participants
21 were asked to report on their experiences with respect to both the app and the study
22 procedures, via questionnaires and an in-depth semi-structured interview. Given that a key
23 objective of this study was to assess the acceptability of the Sleep Ninja app, participants'
24 perspectives were of critical importance. A one-page lay summary of the study results has
25 been sent to all participants.
26
27
28
29
30
31
32
33
34

35 **Statistical Analyses**

36
37 Statistical significance was set at $\alpha=.05$. Sleep diary variables were averaged across
38 each time point (baseline, post-study) to create summary scores. All questionnaire and sleep
39 diary variables were initially screened for excessive skew (>3) or kurtosis (>8 ; Kline, 2011).
40 Four sleep diary entries did not pass screening and were further scrutinised (baseline: WASO
41 entry; post-study: SOL, TST, SE entries). Examination of these four entries revealed each had
42 an extreme value (z -scores ranged from $|4.11|$ to $|6.05|$) and a decision was made to remove
43 these four values. Subsequently all variables had satisfactory skew and kurtosis.
44
45
46
47
48
49
50
51

52 Questionnaire and sleep diary variables were examined using multilevel modelling.
53
54 This modelling approach handles missing data by incorporating all available data from each
55
56
57
58
59
60

subject into the analysis. Given the aims of our study, our interest centred on the main effect of time (i.e., change from baseline- to post). Random effects were modelled for intercept and time. Models were respecified with a random effect for intercept only in cases where there was no variation in individual baseline- to post-study changes. Within-group effect sizes were computed as the modelled mean difference between baseline and post-study divided by the sample standard deviation at baseline.

Results

Baseline Characteristics

See Table 1 for characteristics of the study sample. Participants had a mean age of 13.71 ($SD=1.35$), were spread across school grade, and nearly all were born in Australia and living in the city. Most participants reported difficulty falling asleep, with about half also reporting problems staying asleep or waking up too early, and about a quarter of the sample were receiving treatment for sleep or a mental health problem.

Table 1. Demographic Variables

Characteristics	Sample ($N = 50$)
Age in years, mean (SD , range)	13.71 (1.35, 12-16)
Age in years, n (%)	
12	10 (20.4%)
13	15 (30.6%)
14	7 (14.3%)
15	12 (24.5%)
16	5 (10.2%)
Female, n (%)	33 (66%)
Born in Australia, n (%)	47 (94%)
Live in the city, n (%)	44 (88%)
Sleep problems, n (%)	
Difficulty falling asleep	47 (94%)
Difficulty staying asleep	28 (56%)
Problems waking up too early	28 (56%)
Receiving treatment for sleep or mental health problem, n (%)	13 (26%)

Note. One participant did not indicate their age, so $n = 49$ for age.

Recruitment Rate

There were more than 300 enquires made to the research team about participation in this trial. Of these, 60 individuals indicated eligibility and returned consent forms. Ten of these participants were not enrolled in the trial; four did not meet inclusion criteria and six withdrew prior to the trial. Reasons for withdrawing were: a change of mind ($n=1$), a lack of time ($n=2$), considering participation a chore ($n=1$), confidentiality concerns ($n=1$). One participant did not provide a reason. Therefore, 89% ($n=50$) of the 56 young people who provided consent and met screening criteria continued to trial. See Figure 2 CONSORT diagram for details.

Expectation of Success

Overall, participants were optimistic about using the app, with a mean score of 12.90 ($SD=2.09$) out of a possible 16 points. Every single participant agreed that in principle, people could learn skills for improving sleep from an app, and indicated that they felt that study participation was important. All participants reported that improving their sleep habits were important, with 49% indicating it was 'very important'. Finally, the sample demonstrated their readiness for change by indicating that they were at least moderately ready to improve their sleep patterns using an app.

Retention

Of the 50 participants in the study sample, 47 (94%) completed the baseline questionnaire and sleep diaries and were invited to download the Sleep Ninja. At post-study, 34 participants completed the post-study battery (72% retention). Participants who had available data at both time-points did not differ significantly from those who only had baseline data on any of the questionnaires or sleep diary measures (all F s < 2.58 , $p > .115$).

Uptake and Adherence

Forty-five participants (96%) who completed the baseline assessment downloaded the Sleep Ninja. Program usage data indicated that of these, 82% completed the first lesson, 51% completed four of the six lessons, and 33% completed all six. Participants were accurate in their reporting of app use, with approximately 80% of participants indicating that they completed ‘most’ or ‘almost all’ of the app.

Acceptability

Survey responses on the Acceptability of the Intervention questionnaire indicated that young people overwhelmingly reported that the app was ‘easy’ or ‘very easy’ to use (97%). The majority of participants (59%) indicated that they learnt ‘a fair bit’ from the app, and 28% reported that they learnt ‘a great deal’, while 12% did not learn very much or almost nothing. Participants found the app to be either ‘useful’ or ‘very useful’ (78%), while the remainder (22%) did not find it useful. Most respondents (72%) reported changing their behaviour after using the app, and examples of behaviour change included changes to their pre-bedtime routine (22%), keeping more consistent sleep-wake cycles (65%), getting up earlier in the morning (22%), and restricting the use of their bed for sleep (30%). More than half of the participants reported that they would use this kind of app in the future (56%), and encouragingly, 91% would recommend the Sleep Ninja app to a friend.

The interview mirrored the findings of the questionnaire in terms of acceptability and usefulness. However, there were some aspects of the app that users felt could be improved. Specifically, interviewees expressed a desire for improved explanation of the different app sections and what they needed to do each time they opened the app. Participants commonly reported wanting to be able to personalise their user experience more, including skipping information they knew, seeking more information around difficult or unfamiliar topics, accessing information in different formats (e.g. video/audio), being able to speed up or slow

down the Sleep Ninja's speech, and being able to update their wind-down activity choices and the time the reminder appeared.

Reasons for Non-Adherence

Results from this questionnaire indicated that young people were overwhelmingly happy to use an app to receive help for their sleep issues (84%), did not have technical issues with its use (75%), and felt they had the technical skills to use the app (90%). Participants all reported that the material was relevant and conceptually easy to understand. The main reasons participants reported not using the app was that they felt it took too long to work through (53%), there was too much text to read (47%), and that it was too repetitive (59%).

Preliminary Effects

Sleep and Mental Health Questionnaire Outcomes. Table 2 shows the results for the questionnaire measures. As predicted, from baseline- to post-study, there was a significant decrease in insomnia severity measured by the ISI, $\beta=-4.29$, $p<.001$, $d=-0.90$, and sleep quality on the PSQI, $\beta=-1.88$, $p<.001$; $d=-0.46$. Similarly, mental health measures showed a decrease in both depression on the PHQ-A, $\beta=-2.60$, $p<.001$, $d=-0.36$ and anxiety on the GAD-7, $\beta=-2.56$, $p<.001$, $d=-0.41$.

Table 2. Questionnaire Measures

Outcome	Pre-intervention		Post-intervention		Modelled change from pre- to post-intervention	<i>d</i>
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	β [95% CI]	
ISI	50	14.12 (4.75)	34	9.62 (5.23)	-4.29 [-5.63, -2.95]***	-0.90
PSQI	50	10.43 (4.12)	33	8.03 (4.08)	-1.88 [-2.85, -0.90]***	-0.46
PHQ-A	49	13.04 (7.24)	32	9.88 (7.53)	-2.60 [-3.99, -1.22]***	-0.36
GAD-7	49	9.92 (6.19)	32	7.09 (6.13)	-2.56 [-3.59, -1.52]***	-0.41

Note. Raw means (SDs) are presented. Cohen's *d* values are time effects for pre-intervention to post-intervention using the modelled mean difference divided by the sample pre-intervention SD. ISI = Insomnia Severity Index; PSQI = Pittsburgh Sleep Quality Index; PHQ-A = Patient Health Questionnaire modified for Adolescents; GAD-7 = Generalised Anxiety Disorder 7-item.

*** $p < .001$.

1
2
3 *Sleep Diary Outcomes.* Results for the sleep diary entries are shown in Table 3. At
4 baseline, participants reported taking an average of one hour and 12 minutes to fall asleep,
5 spent an average of 9 hours and 39 minutes in bed, woke up an average of 1.47 times, and
6 slept for a total of seven hours and 40 minutes. Overall sleep efficiency was just above 80%.
7 Results from the analysis indicated that as predicted, there was a significant decrease of 21
8 minutes in how long participants took to fall asleep (SOL; $\beta=-0.37$, $p=.032$) and how
9 frequently they woke during the night reducing to an average of 0.87 times (NWAK; $\beta=-0.46$,
10 $p=.011$). There were also improvements in total sleep time of 33 minutes (TST; $\beta=0.53$,
11 $p=.005$), SE ($\beta=5.25$, $p=.016$), how refreshing sleep was ($\beta=0.43$, $p=.001$) and sleep quality
12 ($\beta=0.31$, $p=.018$). There were no significant differences in TIB, WASO or medication use (all
13 $ps>.05$). Within-group Cohen's d effect sizes ranged from small to medium (0.31-0.68).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Sleep Diary Measures

Outcome	Pre-intervention		Post-intervention		Modelled change from pre- to post-intervention	
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	β [95% CI]	<i>d</i>
Sleep Diary						
TIB (h:min)	48	9:39 (1:11)	29	9:38 (0:55)	-0.01 [-0.42, 0.41]	-0.01
SOL (h:min)	48	1:12 (1:01)	28	0:51 (0:47)	-0.37 [-0.70, -0.03]*	-0.36
WASO (h:min)	47	0:14 (0:14)	29	0:12 (0:14)	-0.04 [-0.12, 0.05]	-0.17
TST (h:min)	48	7:40 (1:09)	28	8:13 (1:06)	0.53 [0.17, 0.90]**	0.46
NWAK	48	1.47 (1.50)	29	0.87 (1.35)	-0.46 [-0.81, -0.11]*	-0.31
SE (%)	48	80.12 (12.01)	28	85.64 (11.30)	5.25 [1.03, 9.47]*	0.44
Sleep refreshingness	48	2.37 (0.63)	29	2.78 (0.78)	0.43 [0.19, 0.68]***	0.68
Sleep quality	48	2.84 (0.74)	29	3.10 (0.82)	0.31 [0.06, 0.56]*	0.42
Use of medication	48	0.12 (0.30)	29	0.11 (0.30)	-0.01 [-0.02, 0.01]	-0.03

Note. Raw means (SDs) are presented. Cohen's *d* values are time effects for pre-intervention to post-intervention using the modelled mean difference divided by the sample pre-intervention SD. Time variables (TIB, SOL, WASO, TST) are expressed in hours:minutes. SE is expressed as a percentage. Refreshingness of sleep is rated on a Likert scale from 1 = *Exhausted* to 5 = *Very refreshed*. Quality of sleep is rated on a Likert scale from 1 = *Very Poor* to 5 = *Very Good*. Use of medication is expressed as a proportion of days medication was used to help with sleep. TIB = Time in bed; SOL = Sleep onset latency; NWAK = Number of awakenings; WASO = Wake after sleep onset; TST = Total sleep time; SE = Sleep efficiency.

* $p < .05$, ** $p < .01$, *** $p < .001$.

Discussion

The purpose of this pilot study was to evaluate the feasibility, acceptability and preliminary effects of the Sleep Ninja app on sleep and mental health symptoms, for use among adolescents with insomnia symptoms. A secondary objective of the study was to gather information in order to refine aspects of the app before evaluating it in a larger trial. Our findings confirmed that young people with sleep disturbance were optimistic about using the app and could complete baseline questionnaires and sleep diaries using an automated digital format without assistance. Feasibility was confirmed based on uptake, completion and retention rates with young people volunteering for the study, downloading the app and completing most of the lessons. This provides evidence that the Sleep Ninja app is a feasible intervention to deliver to young people experiencing sleep difficulties.

Our results also indicate that the app was acceptable and engaging as demonstrated through the acceptability questionnaire and post-study interviews. Overall, adolescents reported that the app was easy to use and contained material relevant to their sleep issues, which resulted in sleep-related behaviour change by the conclusion of the study. Findings identified several areas for improvement and refinement of the app. Specifically, the main reason participants stopped using the app was due to the amount of text presented in the app and the repetitive nature of the material. Participants also requested an improved onboarding process to explain the app components, as well as a more tailored experience. Therefore, refining the app taking these points into account is likely to increase adherence and engagement with the content.

Efficacy outcomes showed that insomnia symptoms improved significantly from baseline to post-study, effectively moving participants from the lower cut off for clinical insomnia, firmly into the sub-threshold symptom level. There was an improvement in self-reported sleep quality, with a medium effect size ($d=-0.46$) suggesting app use improves

1
2
3 quality of sleep. This suggests that the program has benefits beyond just insomnia, which is
4 consistent with research indicating that insomnia and other sleep disturbances (e.g. delayed
5 sleep phase disorder, irregular sleep wake presentations) share common processes and
6 symptoms (Harvey, 2009). The improvements detected on the two sleep questionnaire
7 measures (ISI, PQSI) were corroborated by those found in the sleep diaries, with participants
8 falling asleep more quickly at night, waking less frequency, sleeping for longer and reporting
9 improved sleep efficiency and quality. Findings on these sleep outcomes are consistent with
10 the two studies which have evaluated web-delivered CBT-I for adolescents with insomnia (de
11 Bruin et al., 2015; de Bruin et al., 2014), and show for the first time that CBT-I, when
12 delivered to adolescents by smartphone-app, confer benefits for sleep disturbance. It is
13 encouraging that the within-group effect size obtained in the current study for insomnia ($d=-$
14 0.90) is comparable to within-group effect size found for digitally-delivered CBT-I in a
15 randomised trial ($d=-0.92$) (de Bruin et al., 2015). In this randomised study, the intervention
16 group was found to be superior to the waitlist control, suggesting that a within-group effect
17 size of this magnitude is likely to reflect improvement over and above what would be
18 expected based on a standard placebo effect (de Bruin et al., 2015). Beyond the sleep
19 outcomes, we also found that there were decreases in depression and anxiety symptoms
20 following the completion of the intervention suggesting that there may be value in using this
21 app to address mental illness. The magnitude of the decrease in depression scores is notable,
22 with individuals moving from the moderate range into the mild range, and a within-group
23 effect size that is comparable to other adolescent depression prevention trials (Horowitz,
24 Garber, Ciesla, Young, & Mufson, 2007). These findings provide proof-of-principle evidence
25 that the Sleep Ninja app may be useful in addressing depression. As a pilot study, we did not
26 test the specific hypothesis that targeting insomnia will decrease depression risk, but given
27 our encouraging findings, a follow-up randomised controlled trial which follows participants
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 over time and assesses depression risk is now warranted. This study makes a unique
4 contribution to the literature by showing that smartphone delivery of CBT-I is a promising
5 format in which to deliver this gold-standard intervention. This is the first study that we are
6 aware of that has evaluated app-delivered CBT-I in young people (Werner-Seidler et al.,
7 2018), and only the second study that has tested mobile phone delivery of CBT-I, the first
8 study being conducted in adults, with positive effects on sleep outcomes (Horsch et al.,
9 2017). Using smartphones to deliver interventions such as this offer a myriad of advantages,
10 including immediate connectivity to automated interactive applications that can be accessed
11 anytime, anywhere. Sleep Ninja has been developed so that it does not rely on internet access
12 for use, which is likely to be important to young people who may have limited data plans, and
13 individuals who do not have optimal internet coverage. Not requiring internet coverage to
14 access digital programs represents a new wave of flexibility in the delivery of health
15 interventions and the automated nature of the intervention means it can be delivered without
16 professional support. It is notable that we had a 72% retention rate, which is at the upper level
17 of that detected for digital interventions which has been shown in a meta-analysis to range
18 between 43-99% (Helen Christensen, Griffiths, & Farrer, 2009), with retention rates typically
19 lower in non-supported interventions (Andersson & Cuijpers, 2009).

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
There are several study limitations that need to be considered. First, participants in
this trial were required to have relatively minor levels of insomnia symptoms for study entry.
This decision was made by balancing inclusiveness against the requirement of sleep
disturbance to ensure participants were motivated to use the app. Moreover, given the study
focus was on feasibility and acceptability, we felt it would be prudent to establish these
factors before targeting a more severe participant group. That said, while we set the threshold
for entry relatively low, both the mean and the median converged on an ISI score of just
above 14, indicating that participants were at the junction between having subthreshold and

1
2
3 clinical levels of insomnia symptoms (cut-off score is 15). Therefore, it is relatively unlikely
4 that there would have been a floor effect as the data showed there was sufficient room to
5 detect symptom-improvement. The high mean symptom level also suggests that the results
6 may generalise to a group with clinical levels of insomnia. Second, we did not include a
7 control group. Again, as the study goal was to establish feasibility and acceptability, it was
8 not necessary to include a control group for this purpose. However, this design is not able to
9 attribute causality to the intervention and a controlled study is now needed. Finally, this
10 study relied exclusively on subjectively reported sleep outcomes. Although objective
11 measures such as polysomnography provide the most accurate way to assess sleep, there is
12 evidence that subjectively measured sleep variables could be more closely associated with
13 functional outcomes (Wilson, Fung, Walker, & Barnes, 2013). Moreover, subjectively
14 experienced sleep quality and parameters have consistently shown to be strongly associated
15 with psychological wellbeing (Fuligni & Hardway, 2006; Lund, Reider, Whiting, & Prichard,
16 2010), suggesting that perception of sleep is as important, if not more so, than objective
17 measures. We are currently investigating how inbuilt smartphone sensors such as the
18 accelerometer might be used to provide a more objective estimate of sleep.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
This study provides preliminary evidence supporting the feasibility, acceptability and
effects of a fully-automated app that targets adolescent sleep disturbance. The Sleep Ninja
intervention shows promise both as a sleep-focused intervention, but also potentially to
reduce risk for depression onset.

Figure Legends

Figure 1. Example screens from the Sleep Ninja app. From left: Homescreen, Training Session Access and Progress Record, Tracking and Bedtime Setting, More Information

Figure 2. Participant Flow

For peer review only

Acknowledgements

Sincere thanks to Tobias Reid, Jacinto Santamaria and Cesar Anonuevo for their tireless IT support on this project. Thanks also to Dean Winder for assistance in recruitment. The authors would also like to thank the participants and their families for their enthusiasm in this project. AW-S is supported by a NSW Health Early-to-Mid Career Fellowship, MT is supported by an early-career NHMRC Fellowship #1138710 and HC is supported by NHMRC Fellowship #1056964.

Author Statement

AW-S, BO, MT and HC conceived of the study and the trial design. AW-S designed the study with input from all authors, and oversaw the management of the trial. LJ led trial recruitment, managed the day-to-day running of the trial and conducted the participant interviews. QW conducted the analyses with assistance from AW-S and LJ. All authors contributed to the preparation of the manuscript.

Competing Interests Statement

No competing interests to declare.

Funding Statement

This work was supported by a grant awarded to the Black Dog Institute by the Corella Foundation.

Data Sharing Statement: No additional unpublished data from this study is publicly available.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed. ed.). Washington DC: Author.
- Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: A meta-analysis. *Cognitive Behaviour Therapy*, 38(4), 196-205. doi:10.1080/16506070903318960
- Andrews, G., Issakidis, C., Sanderson, K., Corry, J., & Lapsley, H. (2004). Utilising survey data to inform public policy: Comparison of the cost-effectiveness of treatment of ten mental disorders. *The British Journal of Psychiatry*, 184(6), 526-533.
- Ashworth, D. K., Sletten, T. L., Junge, M., Simpson, K., Clarke, D., Cunnington, D., & Rajaratnam, S. M. (2015). A randomized controlled trial of cognitive behavioral therapy for insomnia: An effective treatment for comorbid insomnia and depression. *Journal of Counselling Psychology*, 62(2), 115-123. doi:10.1037/cou0000059
- Baglioni, C., Battagliese, G., Feige, B., Spiegelhalder, K., Nissen, C., Voderholzer, U., ... Riemann, D. (2011). Insomnia as a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *Journal of Affective Disorders*, 135(1-3), 10-19. doi:10.1016/j.jad.2011.01.011
- Bastien, C. H., Vallieres, A., & Morin, C. M. (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine*, 2(4), 297-307.
- Batterham, P. J., Christensen, H., Mackinnon, A. J., Gosling, J. A., Thorndike, F. P., Ritterband, L. M., ... Griffiths, K. M. (2017). Trajectories of change and long-term outcomes in a randomised controlled trial of internet-based insomnia treatment to prevent depression. *BJPsych Open*, 3(5), 228-235. doi:10.1192/bjpo.bp.117.005231
- Blake, M., Waloszek, J. M., Schwartz, O., Raniti, M., Simmons, J. G., Blake, L., ... Allen, N. B. (2016). The SENSE study: Post intervention effects of a randomized controlled

- 1
2
3 trial of a cognitive-behavioral and mindfulness-based group sleep improvement
4 intervention among at-risk adolescents. *Journal of Consulting and Clinical*
5 *Psychology*, 84(12), 1039-1051. doi:10.1037/ccp0000142
6
7
8
9 Carney, C. E., Buysse, D. J., Ancoli-Israel, S., Edinger, J. D., Krystal, A. D., Lichstein, K. L.,
10 & Morin, C. M. (2012). The Consensus Sleep Diary: Standardizing prospective sleep
11 self-monitoring. *Sleep*, 35(2), 287-302. doi:10.5665/sleep.1642
12
13
14
15 Carskadon, M. A. (1990). Patterns of sleep and sleepiness in adolescents. *Pediatrician*, 17(1),
16 5-12.
17
18
19
20 Chahoud, M., Chahine, R., Salameh, P., & Sauleau, E. A. (2017). Reliability, factor analysis
21 and internal consistency calculation of the Insomnia Severity Index (ISI) in French
22 and in English among Lebanese adolescents. *eNeurologicalSci*, 7, 9-14.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Christensen, H., Batterham, P. J., Gosling, J. A., Ritterband, L. M., Griffiths, K. M.,
Thorndike, F. P., ... Mackinnon, A. J. (2016). Effectiveness of an online insomnia
program (SHUTi) for prevention of depressive episodes (the GoodNight Study): A
randomised controlled trial. *Lancet Psychiatry*, 3(4), 333-341. doi:10.1016/s2215-
0366(15)00536-2
- Christensen, H., Griffiths, K. M., & Farrer, L. (2009). Adherence in internet interventions for
anxiety and depression: Systematic review. *Journal of Medical Internet Research*,
11(2), e13.
- Chung, K. F., Kan, K. K.-K., & Yeung, W.-F. (2011). Assessing insomnia in adolescents:
Comparison of Insomnia Severity Index, Athens Insomnia Scale and Sleep Quality
Index. *Sleep Medicine*, 12(5), 463-470.

- 1
2
3 Cole, J. C., Motivala, S. J., Buysse, D. J., Oxman, M. N., Levin, M. J., & Irwin, M. R. (2006).
4
5 Validation of a 3-factor scoring model for the Pittsburgh Sleep Quality Index in older
6
7 adults. *Sleep*, 29(1), 112.
8
- 9 Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2008).
10
11 Developing and evaluating complex interventions: The new Medical Research
12
13 Council guidance. *BMJ*, 337, a1655. doi:10.1136/bmj.a1655
14
- 15 de Bruin, E. J., Bogels, S. M., Oort, F. J., & Meijer, A. M. (2015). Efficacy of cognitive
16
17 behavioral therapy for insomnia in adolescents: A randomized controlled trial with
18
19 internet therapy, group therapy and a waiting list condition. *Sleep*, 38(12), 1913-1926.
20
21 doi:10.5665/sleep.5240
22
23
- 24 de Bruin, E. J., Oort, F. J., Bogels, S. M., & Meijer, A. M. (2014). Efficacy of internet and
25
26 group-administered cognitive behavioral therapy for insomnia in adolescents: A pilot
27
28 study. *Behavioral Sleep Medicine*, 12(3), 235-254.
29
30 doi:10.1080/15402002.2013.784703
31
32
- 33 de la Vega, R., Tome-Pires, C., Sole, E., Racine, M., Castarlenas, E., Jensen, M. P., & Miro,
34
35 J. (2015). The Pittsburgh Sleep Quality Index: Validity and factor structure in young
36
37 people. *Psychological Assessment*, 27(4), e22-27. doi:10.1037/pas0000128
38
- 39 Franzen, P. L., & Buysse, D. J. (2008). Sleep disturbances and depression: Risk relationships
40
41 for subsequent depression and therapeutic implications. *Dialogues in Clinical*
42
43 *Neuroscience*, 10(4), 473-481.
44
45
- 46 Fuligni, A. J., & Hardway, C. (2006). Daily variation in adolescents' sleep, activities, and
47
48 psychological well-being. *Journal of Research on Adolescence*, 16(3), 353-378.
49
50 doi:10.1111/j.1532-7795.2006.00498.x
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Gitlin, L. N. (2013). Introducing a new intervention: An overview of research phases and
4 common challenges. *American Journal of Occupational Therapy*, 67(2), 177-184.
5
6 doi:10.5014/ajot.2013.006742
7
8
9 Gore, F. M., Bloem, P. J. N., Patton, G. C., Ferguson, J., Joseph, V., Coffey, C., ... Mathers,
10 C. D. (2011). Global burden of disease in young people aged 10-24 years: A
11 systematic analysis. *The Lancet*, 377(9783), 2093-2102. doi:10.1016/S0140-
12 6736(11)60512-6
13
14
15
16
17 Gulliver, A., Griffiths, K. M., & Christensen, H. (2010). Perceived barriers and facilitators to
18 mental health help-seeking in young people: A systematic review. *BMC Psychiatry*,
19 10, 113-113. doi:10.1186/1471-244X-10-113
20
21
22
23
24 Guo, S., Sun, W., Liu, C., & Wu, S. (2016). Structural validity of the Pittsburgh Sleep
25 Quality Index in Chinese undergraduate students. *Frontiers in Psychology*, 7, 1126.
26
27 doi:10.3389/fpsyg.2016.01126
28
29
30
31 Harvey, A. G. (2009). A transdiagnostic approach to treating sleep disturbance in psychiatric
32 disorders. *Cognitive Behaviour Therapy*, 38 Suppl 1, 35-42.
33
34 doi:10.1080/16506070903033825
35
36
37
38 Horowitz, J. L., Garber, J., Ciesla, J. A., Young, J. F., & Mufson, L. (2007). Prevention of
39 depressive symptoms in adolescents: A randomized trial of cognitive-behavioral and
40 interpersonal prevention programs. *Journal of Consulting and Clinical Psychology*,
41 75(5), 693-706. doi:10.1037/0022-006x.75.5.693
42
43
44
45
46 Horsch, C. H., Lancee, J., Griffioen-Both, F., Spruit, S., Fitrianie, S., Neerincx, M. A., ...
47 Brinkman, W. P. (2017). Mobile phone-delivered cognitive behavioral therapy for
48 insomnia: A randomized waitlist controlled trial. *Journal of Medical Internet*
49 *Research*, 19(4), e70. doi:10.2196/jmir.6524
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: Validity of a brief
4 depression severity measure. *Journal of General Internal Medicine*, *16*(9), 606-613.
5
6 doi:10.1046/j.1525-1497.2001.016009606.x
7
8
9 Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B. W., Berry, J. T., & Mokdad, A. H.
10 (2009). The PHQ-8 as a measure of current depression in the general population.
11
12 *Journal of Affective Disorders*, *114*(1), 163-173. doi:10.1016/j.jad.2008.06.026
13
14
15 Liu, X., Buysse, D. J., Gentzler, A. L., Kiss, E., Mayer, L., Kapornai, K., ... Kovacs, M.
16 (2007). Insomnia and hypersomnia associated with depressive phenomenology and
17
18 comorbidity in childhood depression. *Sleep*, *30*(1), 83-90.
19
20
21
22 Lund, H. G., Reider, B. D., Whiting, A. B., & Prichard, J. R. (2010). Sleep patterns and
23
24 predictors of disturbed sleep in a large population of college students. *Journal of*
25
26 *Adolescent Health*, *46*(2), 124-132. doi:10.1016/j.jadohealth.2009.06.016
27
28
29 Manber, R., Edinger, J. D., Gress, J. L., San Pedro-Salcedo, M. G., Kuo, T. F., & Kalista, T.
30 (2008). Cognitive behavioral therapy for insomnia enhances depression outcome in
31
32 patients with comorbid major depressive disorder and insomnia. *Sleep*, *31*(4), 489-
33
34 495.
35
36
37 Ofcom. (2017). *Communications Market Report 2017*. Retrieved from
38
39 https://www.ofcom.org.uk/__data/assets/pdf_file/0017/105074/cmr-2017-uk.pdf
40
41
42 Ohayon, M. M., Roberts, R. E., Zulley, J., Smirne, S., & Priest, R. G. (2000). Prevalence and
43
44 patterns of problematic sleep among older adolescents. *Journal of the American*
45
46 *Academy of Child & Adolescent Psychiatry*, *39*(12), 1549-1556.
47
48 doi:10.1097/00004583-200012000-00019
49
50
51 Okajima, I., Komada, Y., & Inoue, Y. (2011). A meta-analysis on the treatment effectiveness
52
53 of cognitive behavioral therapy for primary insomnia. *Sleep and Biological Rhythms*,
54
55 *9*(1), 24-34. doi:10.1111/j.1479-8425.2010.00481.x
56
57
58
59
60

- 1
2
3 Pew Research Center. (2018). *Teens, Social Media & Technology*. Retrieved from
4 [http://assets.pewresearch.org/wp-](http://assets.pewresearch.org/wp-content/uploads/sites/14/2018/05/31102617/PI_2018.05.31_TeensTech_FINAL.pdf)
5 [content/uploads/sites/14/2018/05/31102617/PI_2018.05.31_TeensTech_FINAL.pdf](http://assets.pewresearch.org/wp-content/uploads/sites/14/2018/05/31102617/PI_2018.05.31_TeensTech_FINAL.pdf)
6
7
8
9 Rickwood, D. J., Deane, F. P., & Wilson, C. J. (2007). When and how do young people seek
10 professional help for mental health problems. *The Medical Journal of Australia*,
11 *187*(7 Suppl), S35-S39.
12
13
14
15 Riemann, D., & Voderholzer, U. (2003). Primary insomnia: A risk factor to develop
16 depression? *Journal of Affective Disorders*, *76*(1-3), 255-259.
17
18
19
20 Spitzer, R. L., Kroenke, K., Williams, J. B., & Lowe, B. (2006). A brief measure for
21 assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*,
22 *166*(10), 1092-1097. doi:10.1001/archinte.166.10.1092
23
24
25
26 Thorndike, F. P., Saylor, D. K., Bailey, E. T., Gonder-Frederick, L., Morin, C. M., &
27 Ritterband, L. M. (2008). Development and perceived utility and impact of an internet
28 intervention for insomnia. *E-journal of Applied Psychology : Clinical and Social*
29 *Issues*, *4*(2), 32-42.
30
31
32
33
34
35 Triton Digital. (2017). *The Infinite Dial Australia*. Retrieved from
36 http://info.tritondigital.com/hubfs/Infinite_Dial_Australia_2017.pdf?t=149990026677
37
38
39
40
41
42 Wagley, J. N., Rybarczyk, B., Nay, W. T., Danish, S., & Lund, H. G. (2013). Effectiveness of
43 abbreviated CBT for insomnia in psychiatric outpatients: Sleep and depression
44 outcomes. *Journal of Clinical Psychology*, *69*(10), 1043-1055.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 to prevent depression and improve cardiac health in adolescence. *BMC Psychology*, 3,
4
5 39. doi:10.1186/s40359-015-0096-x

6
7 Werner-Seidler, A., Johnston, L., & Christensen, H. (2018). Digitally-delivered cognitive-
8
9 behavioural therapy for youth insomnia: A systematic review. *Internet Interventions*,
10
11 11, 71-78. doi:10.1016/j.invent.2018.01.007

12
13 Werner-Seidler, A., O'Dea, B., Shand, F., Johnston, L., Frayne, A., Fogarty, A. S., &
14
15 Christensen, H. (2017). A smartphone app for adolescents with sleep disturbance:
16
17 Development of the Sleep Ninja. *JMIR Mental Health*, 4(3), e28.
18
19 doi:10.2196/mental.7614

20
21
22 Wilson, D. L., Fung, A., Walker, S. P., & Barnes, M. (2013). Subjective reports versus
23
24 objective measurement of sleep latency and sleep duration in pregnancy. *Behavioral*
25
26 *Sleep Medicine*, 11(3), 207-221. doi:10.1080/15402002.2012.670674

27
28
29 Zachariae, R., Lyby, M. S., Ritterband, L. M., & O'Toole, M. S. (2016). Efficacy of internet-
30
31 delivered cognitive-behavioral therapy for insomnia - A systematic review and meta-
32
33 analysis of randomized controlled trials. *Sleep Medicine Reviews*, 30, 1-10.
34
35 doi:10.1016/j.smr.2015.10.004




Figure 1. Example screens from the Sleep Ninja app. From left: Homescreen, Training Session Access and Progress Record, Tracking and Bedtime Setting, More Information

139x238mm (300 x 300 DPI)




Figure 2. Participant Flow

187x238mm (300 x 300 DPI)

BMJ Open

A pilot evaluation of the Sleep Ninja – a smartphone-application for adolescent insomnia symptoms

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-026502.R1
Article Type:	Research
Date Submitted by the Author:	24-Jan-2019
Complete List of Authors:	Werner-Seidler, Aliza ; University of New South Wales, Black Dog Institute Wong, Quincy; University of New South Wales, Black Dog Institute; Western Sydney University School of Science and Health, School of Social Sciences and Psychology Johnston, Lara; University of New South Wales, Black Dog Institute O'Dea, B; University of New South Wales, Black Dog Institute Torok, Michelle; University of New South Wales, Black Dog Institute Christensen, Helen; University of New South Wales - Randwick Campus, Black Dog Institute
Primary Subject Heading:	Mental health
Secondary Subject Heading:	Mental health
Keywords:	insomnia, adolescent mental health, cognitive-behaviour therapy for insomnia, ehealth

SCHOLARONE™
Manuscripts

1
2
3
4
5 **A pilot evaluation of the Sleep Ninja – a smartphone-application for adolescent**
6 **insomnia symptoms**
7
8
9

10
11
12
13
14
15
16
17 Aliza Werner-Seidler¹; a.werner-seidler@blackdog.org.au

18
19 Quincy Wong^{1, 2}; quincy.wong@blackdog.org.au

20
21 Lara Johnston¹; l.johnston@blackdog.org.au

22
23 Bridianne O’Dea¹; b.odea@blackdog.org.au

24
25 Michelle Torok¹; m.torok@blackdog.org.au

26
27
28 Helen Christensen¹; h.christensen@blackdog.org.au
29
30
31
32
33
34

35 **Affiliations:**

36
37 ¹Black Dog Institute, University of New South Wales, Sydney, Australia

38
39 ²School of Social Sciences and Psychology, Western Sydney University, Sydney, NSW,
40
41 Australia.
42
43
44
45

46
47 Corresponding author: Dr Aliza Werner-Seidler; Black Dog Institute, Hospital Road,
48
49 Randwick, New South Wales, Australia; a.werner-seidler@blackdog.org.au; 02 9382 380.
50

51
52 Word Count: 5102
53

54
55 **Keywords:** Insomnia, adolescent mental health, cognitive-behaviour therapy for insomnia,
56
57 eHealth
58
59
60

Abstract

Objectives: The aim of this study was to test the feasibility, acceptability and preliminary effects of a recently developed smartphone application, Sleep Ninja, for adolescent sleep difficulties.

Setting: The study was conducted online with Australian individuals recruited through the community.

Participants: Participants were 50 young people aged 12-16 years with sleep difficulties.

Design: A single-arm pre-post design was used to evaluate feasibility, acceptability and sleep and mental health variables at baseline and post-intervention.

Intervention: Cognitive Behaviour Therapy for Insomnia (CBT-I) informed the development of the Sleep Ninja. The core strategies covered by the app are psychoeducation, stimulus-control, sleep hygiene, and sleep-related cognitive therapy. It includes six training sessions (lessons), a sleep tracking function, recommended bedtimes based on sleep guidelines, reminders to start a wind-down routine each night, a series of sleep tips, and general information about sleep. Users progress through each training session and conclude the six-week program with a black belt in sleep.

Outcome measures: Feasibility was evaluated based on consent rates, adherence and attrition, acceptability was assessed using questionnaires and a post-study interview, and sleep, depression and anxiety variables were assessed at baseline and post-intervention.

Results: Data indicated that the Sleep Ninja is a feasible intervention and is acceptable to young people. Findings showed there were significant improvements on sleep variables including insomnia (within-group effect size $d=-0.90$), sleep quality ($d=-0.46$), depression ($d=-0.36$) and anxiety ($d=-0.41$).

Conclusions: The Sleep Ninja is a promising intervention that could assist adolescents who experience sleep difficulties. A follow-up randomised controlled trial is now warranted.

Trial registration: Australian New Zealand Clinical Trials Registry

(#ACTRN12617000141347).

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Summary

Strengths and limitations of this study

- This is the first study to evaluate app-delivered Cognitive Behavioural Therapy for Insomnia in adolescents with sleep difficulties.
- The intervention being tested, Sleep Ninja, was developed with input from young people, is fully automated and does not require internet coverage to function.
- The evaluation included measures of feasibility and acceptability as well as detailed semi-structured interviews about participants' experience with the app.
- As a preliminary study, this study did not include a control group.

1
2
3 Clinical insomnia is a sleep disturbance characterised by difficulty falling asleep,
4 staying asleep or waking up too early, with associated daytime impairment [1]. It effects
5 approximately 4% of adolescents [2], however sub-threshold symptoms are common, with
6 approximately 25% of young people reporting some degree of sleep disturbance [2, 3].
7
8
9

10
11
12 Depression and insomnia are closely linked, with comorbidity levels as high as 73%
13 in young people [4]. Insomnia is not only a symptom of depression, but is a common
14 precursor, with high quality longitudinal data having established insomnia as an independent
15 risk factor for depression onset [5, 6, 7]. For example, a recent meta-analysis found that
16 insomnia was associated with a greater than two-fold increase in depression risk [5].
17
18 Although depression has multiple causes and maintaining factors that go beyond the presence
19 of sleep problems, the literature suggests that sleep plays an important role, and targeting
20 sleep in the context of depression may have wide-reaching benefits.
21
22
23
24
25
26
27
28
29

30
31 There is emerging evidence that addressing insomnia in individuals with concurrent
32 insomnia and depression improves both sleep and depression outcomes [8, 9, 10]. This
33 suggests there may be value in targeting sleep to improve insomnia symptoms, with potential
34 downstream effects on depression. To our knowledge, there have been three studies testing
35 the hypothesis that targeting insomnia can prevent depressive symptoms. In an adult study,
36 insomnia treatment led to a reduction in depression following the intervention and at 6 and
37 18-month follow-up, relative to an active control group [11, 12]. In a youth study, a face-to-
38 face insomnia intervention was delivered to secondary school students and results showed
39 improvements on sleep and anxiety outcomes, but not symptoms of depression [13]. Data
40 from the two-year follow-up from this study has not yet been published [14]. In a second
41 youth study, a sleep intervention delivered either in group format or digitally led to decreased
42 depressive symptoms at both 2 and 12-month follow-up, an effect that was mediated by
43 improvements in insomnia [15].
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The gold standard treatment for insomnia is cognitive behaviour therapy for insomnia
4 (CBT-I; Australasian Sleep Association, American College of Physicians), and there is
5
6 accumulating evidence to support the use of digitally delivered CBT-I in both adults and
7
8 young people [16, 17, 18, 19]. Delivering sleep interventions via digital formats may be
9
10 particularly well-suited to young people, with adolescents showing a strong preference (97%)
11
12 for digital delivery when given the choice between face-to-face and digital CBT-I [17]. This
13
14 preference may in part be explained by the fact that young people are reluctant to seek help
15
16 for psychological issues, for reasons that include stigma and a preference to manage the
17
18 problem themselves [20]. Sleep is typically less stigmatised than disorders like depression,
19
20 suggesting that it may be more appealing to adolescents. Currently, there are no digital CBT-I
21
22 programs that are commercially available for youth [21]. To overall objective of this study
23
24 was to evaluate a newly-developed digital CBT-I program for adolescents with sleep
25
26 difficulties.
27
28
29
30
31
32

33 The aim of this pilot study was to examine the acceptability, feasibility and
34
35 preliminary effects of an intervention (Sleep Ninja) delivered to adolescents via smartphones.
36
37 In line with the guidelines on the development of behavioural interventions [22, 23], the
38
39 primary purpose of this study was to investigate recruitment rates, uptake, intervention
40
41 completion, reasons for non-adherence, and participant retention. The secondary aim was to
42
43 use both quantitative and qualitative methods to determine the acceptability of the app among
44
45 young people with sleep difficulties and allow for the refinement of the intervention prior to a
46
47 formal randomised evaluation. A final aim was to examine the impact of the Sleep Ninja app
48
49 on sleep outcomes and mental health symptoms. We used a single-arm, pre-post design to
50
51 address these aims. It was hypothesised that the app would be a feasible modality in which to
52
53 deliver the automated sleep intervention, as measured by uptake, completion and retention
54
55
56
57
58
59
60

1
2
3 rates, that the app would be acceptable to young people, and that its use would be associated
4
5 with improvement in sleep and mental health symptoms.
6
7
8
9

10 **Method**

11
12 This trial was prospectively registered on the Australian New Zealand Clinical Trials
13
14 Registry (#ACTRN12617000141347).
15
16

17 **Participants**

18
19 Fifty participants were recruited via media and social media channels, including the
20
21 Black Dog Institute's website and paid Facebook advertisements that targeted potential
22
23 participants and their parents. Inclusion criteria were: aged 12-16 years, presence of at least
24
25 mild insomnia, operationalised by endorsement of at least one of the following symptoms
26
27 over the preceding two-week period: difficulty falling asleep, difficulty staying asleep or
28
29 waking up too early. These items are the first three questions on the Insomnia Severity Index
30
31 [24], and were chosen to include a participant group with at least mild levels of insomnia. For
32
33 study inclusion, participants also needed to own a smartphone running iOS or Android, have
34
35 a valid email address, access to the internet, and be able to provide personal and parental
36
37 consent.
38
39
40
41
42
43
44

45 **Measures**

46
47 **Insomnia Severity Index (ISI).** The ISI is a psychometrically sound, seven-item self-
48
49 report measure of insomnia symptoms over the previous two weeks [24, 25]. Responses are
50
51 reported on a Likert scale from 0 to 4, producing total scores of 0 to 28 [24, 25]. Cut-off
52
53 scores are as follows: 0-7 reflects no clinically significant insomnia, 8-14 indicates
54
55 subthreshold insomnia, 15-21 suggests moderate severity insomnia, and 22-28 indicates
56
57 severe insomnia [24]. The ISI was designed for use in adults but has been widely
58
59
60

1
2
3 administered to, and validated in, adolescent samples [25, 26, 27]. In one adolescent
4 validation study, reliability was strong (Cronbach's $\alpha=0.83$), and test-retest reliability was
5 acceptable, $r=.79$ [25].
6
7

8
9
10 **Pittsburgh Sleep Quality Index (PSQI).** The PSQI is a widely used self-report 19-
11 item scale that assesses usual sleep habits and experiences over the preceding month and has
12 been validated in adolescent samples, with strong internal consistency ($\alpha=.72$) and test-retest
13 reliability over a 6-week period ($r=.81$) [28]. There are seven sub-scales which are sleep
14 quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of
15 sleeping medications, daytime dysfunction [29]. Each component is scored from 0 (no
16 difficulty) to 3 (severe difficulty), which are summed to obtain a Global PSQI score ranging
17 from 0 to 21 [30].
18
19

20
21
22 **Patient Health Questionnaire – Adolescent Version (PHQ-A).** The PHQ-A
23 assesses depressive symptoms in the preceding two weeks in adolescents, and has been
24 adapted from the widely used PHQ-9 designed for adults [31]. This measure has excellent
25 psychometric properties, including $\alpha= 0.89$, and test-retest reliability of $r=0.84$ [32]. In this
26 study, we used the 8-item version in which the questions are identical to those asked in the
27 PHQ-9 with the exclusion of the last item which asks about suicide and is comparable to the
28 PHQ-9 in terms of diagnosing depressive disorders [32, 33]. Each item is scored on a 4-point
29 scale and summed together to form a total depression score ranging from 0 to 24. Scores
30 correspond to the following cut-offs: 0-9 indicates minimal symptoms, 10-14 indicates mild
31 symptoms, 15-19 reflects moderate symptoms, and 20-24 is indicative of severe depression
32 [31]. The PHQ-A has demonstrated good sensitivity (73%) and high specificity (94%) for
33 major depressive disorder [31].
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55
56 **Generalised Anxiety Disorder 7-item (GAD-7).** The GAD-7 evaluates symptoms of
57 generalised anxiety disorder [34]. All items are scored on a scale from 0 (not at all) to 3
58
59
60

1
2
3 (nearly every day). The scores on each item are summed together to derive a total score,
4 ranging from 0 to 21 of which 0-4 indicates minimal anxiety, 5-9 mild anxiety, 10-14
5 moderate anxiety, and 15-21 severe anxiety [34]. The GAD-7 has good sensitivity (89%) and
6 specificity (82%) for GAD scores >10 [34]. The measure has also been validated in
7 adolescent populations with Cronbach's $\alpha=0.90$ and high convergent and discriminant
8 validity [35].
9
10
11
12
13
14
15
16

17 **Expectations of Success.** A four-item scale was developed for this study to assess
18 participants' motivation and expectations for improving their sleep with an app (e.g., *I am*
19 *confident that people could learn skills for improving sleep from an app*). The four items
20 assessed perceived confidence, importance, usefulness and readiness to change. The
21 Expectation of Success measure was scored on a five-point scale and total scores were
22 computed by summing each item, ranging from 0 to 16. Higher scores on this scale indicate
23 greater confidence and readiness to target sleep using a smartphone app.
24
25
26
27
28
29
30
31
32

33 **Acceptability of the Intervention.** The Acceptability of the Intervention scale is a
34 seven-item measure that was developed by the research team to assess participants' attitudes
35 and behaviours associated with using the app (e.g., *How much did you learn from the app* and
36 *Would you recommend this program to others?*). This measure was informed by similar
37 acceptability measures commonly used in the field [36]. Each item was designed to assess a
38 different domain. The first four domains to be assessed were app completion, ease of use,
39 amount learnt and usefulness, with each being scored on an ordinal scale from 0-3. The final
40 three items assessed behaviour change, whether the participant would use an app like this in
41 the future, and whether they would recommend it to a friend, and were scored dichotomously
42 as either yes or no, with an option for participants to describe the nature of their behaviour
43 change if yes. As each question assessed a different domain, item scores were considered
44 separately.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Reasons for Non-Adherence.** The Reasons for Non-Adherence measure is a 23-item
4 scale that was adapted from a previous measure [37] to assess the degree to which different
5 reasons impacted on participants' use of the app. There are four domains assessed:
6 phone/internet/technical issues (*e.g., My phone wasn't working or was having problems*);
7 personal issues (*e.g., I didn't think I deserved help*); intervention-general issues (*e.g., I wasn't*
8 *convinced the app would be helpful*); and intervention-specific issues (*e.g., There was too*
9 *much text to read*). Participants responded on an ordinal scale indicating whether each item
10 played no, a little or major part in why they stopped or had difficulty using the app as
11 intended. The scores on each item were considered separately.
12
13
14
15
16
17
18
19
20
21
22
23

24 **Sleep Diary.** The ten-item Sleep Diary was developed by the research team
25 incorporating the questions from the Consensus Sleep Diary [38], with the addition of two
26 questions regarding daytime naps and use of sleep medication. Participants answered 10
27 questions which included bedtime, time taken to fall asleep (sleep onset latency; SOL),
28 number and duration of night-time awakenings (number of awakenings; NWAK, duration of
29 wakefulness after sleep onset; WASO), time of final awakening, time participants got out of
30 bed for the day, subjective sleep quality, how refreshed participants felt upon awakening,
31 duration of any daytime naps and use of sleep medication. Sleep diaries were completed
32 electronically with pre-set categories from which users selected responses from a drop-down
33 menu. A clock scroller was used to enter the time and/or duration of all sleep-related
34 activities and all times were entered in 12-hour format to minimise errors associated with 24-
35 hour time. Restrictions were set to ensure participants could not enter a wake time earlier than
36 bedtime and visa versa. All questions required answers for the sleep diary to be submitted.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 time between waking and getting up in the morning, from TIB) and sleep efficiency (SE;
4
5 calculated by taking the percentage of TST/TIB).
6
7

8 **Post-Study Interview.** After study completion, participants were invited to attend a
9
10 face-to-face or telephone interview to provide feedback on their experience. Interviews were
11
12 semi-structured and explored participants' opinions about the study in general, and
13
14 specifically in relation to the intervention. Questions were open-ended, and flexible enough to
15
16 explore ideas that were raised during each interview. Interviews were audio recorded and
17
18 then transcribed verbatim by the interviewer. The interview content was pragmatically coded
19
20 into relevant themes by the same researcher, with oversight and guidance provided by the
21
22 research team.
23
24
25

26 **Intervention – ‘Sleep Ninja’**

27
28 The Sleep Ninja app was derived from CBT-I and developed by our team, as a fully-
29
30 automated smartphone app. A participatory design process was used whereby young people
31
32 contributed to the content, functionality and accessibility/user experience of the app through a
33
34 series of focus groups [39]. The core strategies included in the app were: psychoeducation,
35
36 stimulus-control, sleep hygiene and sleep-focused cognitive therapy. Sleep restriction, which
37
38 aims to increase sleep efficiency by reducing the amount of time spent in bed, was
39
40 deliberately omitted because some support (parental and/or professional) is likely to be
41
42 required to successfully implement sleep restriction, particularly in young people. Although
43
44 sleep restriction did not comprise part of the app, there was instead a focus on the importance
45
46 or regular sleep-wake cycles. The app teaches users about the importance of consistent sleep
47
48 and wake times, and recommended bedtimes are calculated based on the time they need to
49
50 wake up (according to sleep guidelines). This strategy draws from transdiagnostic approaches
51
52 to target sleep difficulties that go beyond insomnia (e.g., delayed sleep phase and irregular
53
54
55
56
57
58
59
60

1
2
3 sleep presentations; [40]) and may therefore be useful to adolescents experiencing a broad
4
5 range of sleep difficulties.
6

7
8 The structure of the Sleep Ninja app includes six training sessions (lessons), a sleep
9
10 tracking function, recommended bedtimes based on sleep guidelines, reminders to start a
11
12 wind-down routine each night, a series of sleep tips, and general information about sleep. The
13
14 home screen has 3 options: Train, Track, and More (see Figure 1). Users complete training
15
16 sessions which are delivered through a chat-bot format where the sleep ninja essentially acts
17
18 as a sleep coach. Training sessions take approximately 5-10 minutes to complete, and cover:
19
20 (i) psychoeducation, information about circadian rhythms and the importance of keeping
21
22 regular sleep schedules; (ii) stimulus control, the value of only going to sleep when tired, and
23
24 strategies that can be used at night when having trouble sleeping; (iii) basic sleep hygiene
25
26 such as avoiding caffeine and stimulating activity in the evenings, suggestions for daytime
27
28 activities to promote night-time sleep (e.g., exercise, no napping); (iv) identifying and
29
30 planning for high-risk situations, how to get back on track after a late-night or sleep in; (v)
31
32 cognitive therapy including how to deal with unhelpful thoughts that can prevent falling
33
34 asleep as well as sleep-related cognitive distortions cognitions and; (vi) a final review
35
36 session which summarises all of the material contained in the app. The user interacts with the
37
38 app through a forced choice chat-bot format which is responsive to the input of the user,
39
40 meaning it personalises information and recommendations based on the selections and sleep
41
42 profile recorded by the participant. Users level up and reach their next “belt” by completing
43
44 one training session and tracking their sleep for 3 nights (out of a 7-night period). As there
45
46 were six training sessions to complete, the app was made available for six weeks (42 days)
47
48 before it locked. Users finish the program with a black belt in sleep.
49
50
51
52
53
54

55
56 It is notable that evidence suggests that the use of screens at bedtime interferes with
57
58 sleep [41]. Drawing on data from large epidemiological studies suggesting that refraining
59
60

1
2
3 from screen use for the hour prior to bedtime alleviates potential interference from screens
4
5 [42], this app has been designed to be used during the day. Users receive a prompt one hour
6
7 before bed (calculated according to sleep guidelines and their wake-up time) to commence
8
9 their pre-bed routine and are encouraged to stop using electronic devices after this time. In
10
11 fact, part of the cognitive component of the app is to educate and challenge beliefs about the
12
13 importance of night time phone use in order to promote healthy sleep habits. We expect these
14
15 factors to mitigate the risk that smartphone use in this context will contribute to poor sleep.
16
17

18 19 **Procedure**

20
21 All procedures were approved by the University of New South Wales Human
22
23 Research Ethics Committee (HC#16702). Participants were encouraged to download consent
24
25 forms if they met the eligibility criteria listed on the study website and submit this directly to
26
27 the research team, once completed. Those who provided written informed consent and that of
28
29 a parent or guardian were then enrolled in the trial and invited to complete the screening
30
31 questions to verify study eligibility, before completing baseline questionnaires which
32
33 included: demographics, ISI, PSQI, PHQ-A, GAD-7 and Expectation of Success. Participants
34
35 could then access the first day of the online seven-day sleep diary. Another diary entry
36
37 became available each day for the following six days and participants were reminded to
38
39 complete entries via text-message. At the completion of seven consecutive entries in the
40
41 diary, participants were given access to the Sleep Ninja app on their personal smartphone
42
43 devices. Participants could use the app for six weeks before the post study questionnaire was
44
45 made available, which included the same battery as baseline with the omission of the
46
47 Expectations of Success questionnaire and with the addition of the Acceptability and Reasons
48
49 for Non-Adherence questionnaires. Participants then completed another seven-day sleep
50
51 diary, which was delivered in the same format and schedule as baseline. After the study had
52
53 finished, participants were invited to participate in a face-to-face or telephone interview to
54
55
56
57
58
59
60

1
2
3 provide feedback on their experience of participating in the study. Participants were
4
5 reimbursed for their time with giftcards to the value of \$10 each for completing baseline and
6
7 post-study assessment schedules; \$20 for a telephone interview and \$30 for a face-to-face
8
9 interview.
10

11 **Participant Involvement and Consultation**

12
13
14
15 Prior to this study, a separate group of young people were consulted in a series of
16
17 focus groups to inform the design, features and structure of the app (for more information
18
19 please see [39]). As a feasibility and acceptability study, participants were asked to report on
20
21 their experiences with respect to both the app and the study procedures, via questionnaires
22
23 and an in-depth semi-structured interview. Given that a key objective of this study was to
24
25 assess the acceptability of the Sleep Ninja app, participants' perspectives were of critical
26
27 importance. A one-page lay summary of the study results has been sent to all participants.
28
29

30 **Statistical Analyses**

31
32
33 Statistical significance was set at $\alpha=.05$. Summary scores for sleep diary variables at
34
35 baseline and post-study were obtained by averaging sleep diary entries at baseline, and
36
37 averaging sleep diary entries at post-study, respectively. All questionnaire and sleep diary
38
39 variables were initially screened for excessive skew (>3) or kurtosis (>8 ; [43]). Six sleep
40
41 diary variables did not pass screening and were further scrutinised (baseline: WASO entry;
42
43 post-study: Time fallen asleep, Time in bed after final morning wake up, SOL, TST, SE
44
45 variables). Examination of these six variables revealed each included an entry that was of an
46
47 extreme value (z -scores ranged from $|4.11|$ to $|6.05|$) and a decision was made to remove these
48
49 six values (hence, $n = 47$ for Baseline WASO; $n_s = 28$ for Post-intervention Time fallen
50
51 asleep, Time in bed after final morning wake up, SOL, TST, and SE). Subsequently all
52
53 variables had satisfactory skew and kurtosis.
54
55
56
57
58
59
60

Questionnaire and sleep diary variables were examined using multilevel modelling. This modelling approach handles missing data by incorporating all available data from each subject into the analysis. Given the aims of our study, our interest centred on the main effect of time (i.e., change from baseline- to post). Random effects were modelled for intercept and time. Models were respecified with a random effect for intercept only in cases where there was no variation in individual baseline- to post-study changes. Within-group effect sizes were computed as the modelled mean difference between baseline and post-study divided by the sample standard deviation at baseline.

Results

Baseline Characteristics

See Table 1 for characteristics of the study sample. Participants had a mean age of 13.71 ($SD=1.35$), were spread across school grade, and nearly all were born in Australia and living in the city. Most participants reported difficulty falling asleep, with about half also reporting problems staying asleep or waking up too early, and about a quarter of the sample were receiving treatment for sleep or a mental health problem.

Table 1. Demographic Variables

Characteristics	Sample ($N = 50$)
Age in years, mean (SD , range)	13.71 (1.35, 12-16)
Age in years, n (%)	
12	10 (20.4%)
13	15 (30.6%)
14	7 (14.3%)
15	12 (24.5%)
16	5 (10.2%)
Female, n (%)	33 (66%)
Born in Australia, n (%)	47 (94%)
Live in the city, n (%)	44 (88%)
Sleep problems, n (%)	
Difficulty falling asleep	47 (94%)
Difficulty staying asleep	28 (56%)
Problems waking up too early	28 (56%)
Receiving treatment for sleep or mental health problem, n (%)	13 (26%)

1
2
3 *Note.* One participant did not indicate their age, so $n = 49$ for age.
4
5
6
7

8 9 **Recruitment Rate**

10
11 There were more than 300 enquires made to the research team about participation in
12 this trial. Of these, 60 individuals indicated eligibility and returned consent forms. Ten of
13 these participants were not enrolled in the trial; four did not meet inclusion criteria and six
14 withdrew prior to the trial. Reasons for withdrawing were: a change of mind ($n=1$), a lack of
15 time ($n=2$), considering participation a chore ($n=1$), confidentiality concerns ($n=1$). One
16 participant did not provide a reason. Therefore, 89% ($n=50$) of the 56 young people who
17 provided consent and met screening criteria continued to trial. See Figure 2 CONSORT
18 diagram for details.
19
20
21
22
23
24
25
26
27
28

29 **Expectation of Success**

30
31 Overall, participants were optimistic about using the app, with a mean score of 12.90
32 ($SD=2.09$) out of a possible 16 points. Every single participant agreed that in principle,
33 people could learn skills for improving sleep from an app, and indicated that they felt that
34 study participation was important. All participants reported that improving their sleep habits
35 were important, with 49% indicating it was 'very important'. Finally, the sample
36 demonstrated their readiness for change with 100% of the sample indicating that they were
37 either moderately ready (16%), ready (43%) or completely ready (41%) to improve their
38 sleep patterns using an app.
39
40
41
42
43
44
45
46
47
48
49

50 **Retention**

51
52 Of the 50 participants in the study sample, 47 (94%) completed the baseline
53 questionnaire and sleep diaries and were invited to download the Sleep Ninja. At post-study,
54 34 participants completed the post-study battery (72% retention). Participants who had
55
56
57
58
59
60

1
2
3 available data at both time-points did not differ significantly from those who only had
4
5 baseline data on any of the questionnaires or sleep diary measures (all F s < 2.58, p s > .115).
6
7

8 **Uptake and Adherence**

9
10 Forty-five participants (96%) who completed the baseline assessment downloaded the
11
12 Sleep Ninja. Program usage data indicated that of these, 82% completed the first lesson, 51%
13
14 completed four of the six lessons, and 33% completed all six. Participants were accurate in
15
16 their reporting of app use, with approximately 80% of participants indicating that they
17
18 completed ‘most’ or ‘almost all’ of the app.
19
20

21 **Acceptability**

22
23 Survey responses on the Acceptability of the Intervention questionnaire indicated that
24
25 young people reported that the app was ‘easy’ or ‘very easy’ to use (97%). The majority of
26
27 participants (59%) indicated that they learnt ‘a fair bit’ from the app, and 28% reported that
28
29 they learnt ‘a great deal’, while 12% did not learn very much or almost nothing. Participants
30
31 found the app to be either ‘useful’ or ‘very useful’ (78%), while the remainder (22%) did not
32
33 find it useful. Most respondents (72%) reported changing their behaviour after using the app,
34
35 and examples of behaviour change included changes to their pre-bedtime routine (22%),
36
37 keeping more consistent sleep-wake cycles (65%), getting up earlier in the morning (22%),
38
39 and restricting the use of their bed for sleep (30%). More than half of the participants
40
41 reported that they would use this kind of app in the future (56%), and encouragingly, 91%
42
43 would recommend the Sleep Ninja app to a friend. The degree to which participants found the
44
45 app useful was positively correlated with module completion, ($r=.35$, $p=.047$), as was the
46
47 degree to which participants reported learning from the app ($r=.49$, $p=.004$).
48
49
50
51
52
53

54 The interview mirrored the findings of the questionnaire in terms of acceptability and
55
56 usefulness. However, there were some aspects of the app that users felt could be improved.
57
58 Specifically, interviewees expressed a desire for improved explanation of the different app
59
60

1
2
3 sections and what they needed to do each time they opened the app. Participants commonly
4
5 reported wanting to be able to personalise their user experience more, including skipping
6
7 information they knew, seeking more information around difficult or unfamiliar topics,
8
9 accessing information in different formats (e.g. video/audio), being able to speed up or slow
10
11 down the Sleep Ninja's speech, and being able to update their wind-down activity choices
12
13 and the time the wind-down reminder appeared. Participants expressed a range of views
14
15 about the tone of the Sleep Ninja, with nearly half of the interviewed participants
16
17 commenting favourably on the Sleep Ninja's jokes, with several participants commenting that
18
19 the Sleep Ninja's language was annoying and too childish. There was consensus that the
20
21 Sleep Ninja's language was repetitive and could be improved by cutting out superfluous
22
23 dialogue that was not delivering core intervention strategies. Nearly half of the interviewed
24
25 participants expressed some difficulty in implementing at least one of the Sleep Ninja's
26
27 recommended strategies due to conflicting parental bedtime rules. For instance, the strategies
28
29 to delay bedtime until sleepy and leave the bed/bedroom if unable to get to sleep after more
30
31 than 30 minutes most commonly encountered parental resistance or required modification.
32
33 Numerous interviewees commented on the usefulness of receiving feedback and summaries
34
35 of their logged sleep, however several commented that this could be improved by displaying
36
37 the information in graphs and over time, so that change and improvement is clearer. Overall,
38
39 participants reported that the number of notifications in the app were acceptable, and that
40
41 additional reminders should be sent to notify them of available lessons and after periods of
42
43 inactivity and that these reminders should contain motivational and encouraging messages.
44
45 While most interviewees considered three nights of sleep tracking per belt acceptable, there
46
47 were others who felt fewer nights of tracking would have been better, and others who
48
49 expressed willingness to track more than three nights before levelling up.
50
51
52
53
54
55
56
57
58
59
60

Reasons for Non-Adherence

Results from this questionnaire indicated that young people were very happy to use an app to receive help for their sleep issues (84%), did not have technical issues with its use (75%), and felt they had the technical skills to use the app (90%). Participants all reported that the material was relevant and conceptually easy to understand. The main reasons participants reported not using the app was that they felt it took too long to work through (53%), there was too much text to read (47%), and that it was too repetitive (59%).

Preliminary Effects

Sleep and Mental Health Questionnaire Outcomes. Table 2 shows the results for the questionnaire measures. As predicted, from baseline- to post-study, there was a significant decrease in insomnia severity measured by the ISI, $\beta=-4.29$, $p<.001$, $d=-0.90$, and sleep quality on the PSQI, $\beta=-1.88$, $p<.001$; $d=-0.46$. Similarly, mental health measures showed a decrease in both depression on the PHQ-A, $\beta=-2.60$, $p<.001$, $d=-0.36$ and anxiety on the GAD-7, $\beta=-2.56$, $p<.001$, $d=-0.41$.

Table 2. Questionnaire Measures

Outcome	Pre-intervention		Post-intervention		Modelled change from pre- to post-intervention	<i>d</i>
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	β [95% CI]	
ISI	50	14.12 (4.75)	34	9.62 (5.23)	-4.29 [-5.63, -2.95]***	-0.90
PSQI	50	10.43 (4.12)	33	8.03 (4.08)	-1.88 [-2.85, -0.90]***	-0.46
PHQ-A	49	13.04 (7.24)	32	9.88 (7.53)	-2.60 [-3.99, -1.22]***	-0.36
GAD-7	49	9.92 (6.19)	32	7.09 (6.13)	-2.56 [-3.59, -1.52]***	-0.41

Note. Raw means (SDs) are presented. Cohen's *d* values are time effects for pre-intervention to post-intervention using the modelled mean difference divided by the sample pre-intervention SD. ISI = Insomnia Severity Index; PSQI = Pittsburgh Sleep Quality Index; PHQ-A = Patient Health Questionnaire modified for Adolescents; GAD-7 = Generalised Anxiety Disorder 7-item.

*** $p < .001$.

Sleep Diary Outcomes. Results for the sleep diary entries are shown in Table 3. At baseline, participants went to sleep, on average at 11:29pm, reported taking an average of one hour and 12 minutes to fall asleep, spent an average of 9 hours and 39 minutes in bed, woke

1
2
3 up an average of 1.47 times, slept for a total of seven hours and 40 minutes, woke up at
4
5 7:27am, and spent approximately 29 minutes awake in bed before getting up. Overall sleep
6
7 efficiency was just above 80%. Results from the analysis at post-intervention indicated that as
8
9 predicted, participants went to bed 35 minutes earlier than at baseline ($\beta=-0.58, p=.003$),
10
11 there was a significant decrease of 21 minutes in how long participants took to fall asleep
12
13 (SOL; $\beta=-0.37, p=.032$), participants spent significantly less time in bed after waking than
14
15 they did at baseline ($\beta=-0.27, p<.001$), and woke significantly less frequently during the night
16
17 reducing to an average of 0.87 times (NWAK; $\beta=-0.46, p=.011$). There were also
18
19 improvements in total sleep time of 33 minutes (TST; $\beta=0.53, p=.005$), SE ($\beta=5.25, p=.016$),
20
21 how refreshing sleep was ($\beta=0.43, p<.001$) and sleep quality ($\beta=0.31, p=.018$). There were no
22
23 significant differences in the time participants woke up in the morning (on average, at
24
25 7:20am), TIB, WASO or medication use (all $ps>.05$). Within-group Cohen's *d* effect sizes
26
27 ranged from small to medium (0.31-0.68).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Sleep Diary Measures

Outcome	Baseline		Post-intervention		Modelled change from baseline- to post- intervention	<i>d</i>
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	β [95% CI]	
Time woken up	48	07:27 (0:46)	29	07:20 (0:54)	-0.10 [-0.41-0.21]	0.13
Time in bed after final morning wake up (h:min)	48	0:29 (0:23)	28	0:13 (0:10)	-0.27 [-0.41, -0.12]***	0.70
Time fallen asleep	48	23:29 (1:19)	28	22:54 (1:03)	-0.58 [-0.95, -0.21]**	0.44
TIB (h:min)	48	9:39 (1:11)	29	9:38 (0:55)	-0.01 [-0.42, 0.41]	0.01
SOL (h:min)	48	1:12 (1:01)	28	0:51 (0:47)	-0.37 [-0.70, -0.03]*	0.36
WASO (h:min)	47	0:14 (0:14)	29	0:12 (0:14)	-0.04 [-0.12, 0.05]	0.17
TST (h:min)	48	7:40 (1:09)	28	8:13 (1:06)	0.53 [0.17, 0.90]**	0.46
NWAK	48	1.47 (1.50)	29	0.87 (1.35)	-0.46 [-0.81, -0.11]*	0.31
SE (%)	48	80.12 (12.01)	28	85.64 (11.30)	5.25 [1.03, 9.47]*	0.44
Sleep refreshingness	48	2.37 (0.63)	29	2.78 (0.78)	0.43 [0.19, 0.68]***	0.68
Sleep quality	48	2.84 (0.74)	29	3.10 (0.82)	0.31 [0.06, 0.56]*	0.42
Use of medication	48	0.12 (0.30)	29	0.11 (0.30)	-0.01 [-0.02, 0.01]	0.03

Note. Raw means (SDs) are presented. Cohen's *d* values are time effects for pre-intervention to post-intervention using the modelled mean difference divided by the sample pre-intervention SD. Time fallen asleep and Time woken up are expressed as times in 24-hour time. Time variables (TIB, SOL, WASO, TST) are expressed in hours:minutes. TIB = Time in bed; SOL = Sleep onset latency; WASO = Wake after sleep onset; TST = Total sleep time; NWAK = Number of awakenings; SE = Sleep efficiency. SE is expressed as a percentage. Refreshingness of sleep is rated on a Likert scale from 1 = *Exhausted* to 5 = *Very refreshed*. Quality of sleep is rated on a Likert scale from 1 = *Very Poor* to 5 = *Very Good*. Use of medication is expressed as a proportion of days medication was used to help with sleep.

* $p < .05$, ** $p < .01$, *** $p < .001$.

Discussion

The purpose of this pilot study was to evaluate the feasibility, acceptability and preliminary effects of the Sleep Ninja app on sleep and mental health symptoms, for use among adolescents with sleep difficulties. A secondary objective of the study was to gather information in order to refine aspects of the app before evaluating it in a larger trial. Our findings confirmed that young people with sleep difficulties were optimistic about using the app and could complete baseline questionnaires and sleep diaries using an automated digital format without assistance. Feasibility was confirmed based on uptake, completion and retention rates with young people volunteering for the study, downloading the app and completing most of the lessons. This provides evidence that the Sleep Ninja app is a feasible intervention to deliver to young people experiencing sleep difficulties.

Intervention adherence levels suggested that while more than half of the participants completed more than half of the app, only one third of participants completed all six lessons. While this is within the range of adherence rates reported in the literature for technology-mediated insomnia programs [44], reasons for non-adherence require consideration as there is room for improvement. The main reason participants had difficulty or stopped using the app was reportedly due to the amount of text presented in the app and the repetitive nature of the material. Participants also requested a more tailored experience. Those who completed more of the app also reported it to be more acceptable. Therefore, it is likely that refining the app by taking these points into account is likely to increase engagement with the content and overall adherence to the intervention.

Efficacy outcomes showed that insomnia symptoms improved significantly from baseline to post-study, effectively moving participants from the lower cut off for clinical insomnia, firmly into the sub-threshold symptom level. There was an improvement in self-reported sleep quality, with a medium effect size ($d=-0.46$) suggesting app use improves

1
2
3 quality of sleep. The improvements detected on the two sleep questionnaire measures (ISI,
4 PQSI) were corroborated by those found in the sleep diaries, with participants going to bed
5 earlier, falling asleep more quickly at night, waking less frequently, sleeping for longer,
6
7
8
9
10 spending less time in bed in the morning after waking, and reporting improved sleep
11
12 efficiency and quality.
13

14
15 The time that participants reported waking each morning did not shift. This finding
16
17 needs to be considered in the context of young people having to get up at a specific time for
18
19 school each morning. School start times are a modifiable contributing factor to insufficient
20
21 sleep, with evidence suggesting that delaying school start times by even 30 minutes is
22
23 associated with higher levels of school attendance, lower daytime sleepiness, and improved
24
25 attention and concentration [45]. While there have been calls to delay school start times to
26
27 improve sleep duration, health and functioning (e.g., [46]), this is unlikely to change in the
28
29 Australian context. Accordingly, a focus on bed times rather than rise times is likely to be
30
31 most useful for treating sleep difficulties in this age group, at least for the time being.
32
33
34

35
36 Findings on these sleep outcomes are consistent with the two studies which have
37
38 evaluated web-delivered CBT-I for adolescents with insomnia [15, 16, 17], and show for the
39
40 first time that CBT-I, when delivered to adolescents by smartphone-app, confer benefits. It is
41
42 encouraging that the within-group effect size obtained in the current study for insomnia ($d=$
43
44 0.90) is comparable to within-group effect size found for digitally-delivered CBT-I in a
45
46 randomised trial ($d=-0.92$; [16]). In this randomised study, the intervention group was found
47
48 to be superior to the waitlist control, suggesting that a within-group effect size of this
49
50 magnitude is likely to reflect improvement over and above what would be expected based on
51
52 a standard placebo effect [16].
53
54

55
56 Beyond the sleep outcomes, we also found that there were decreases in depression and
57
58 anxiety symptoms following the completion of the intervention suggesting that there may be
59
60

1
2
3 value in using this app to address mental illness. The magnitude of the decrease in depression
4 scores is notable, with individuals moving from the moderate range into the mild range, and a
5 within-group effect size that is comparable to other adolescent depression prevention trials
6 [47]. These findings provide proof-of-principle evidence that the Sleep Ninja app may be
7
8 useful in addressing depression. As a pilot study, we did not test the specific hypothesis that
9
10 targeting insomnia will decrease depression risk. Given our encouraging findings, a follow-
11
12 up randomised controlled trial which follows participants over time, which can determine
13
14 causality between insomnia and depression, that can assess the mechanisms of change, as
15
16 well as the impact of intervention on depression risk is now warranted.
17
18
19
20
21
22
23

24 This study makes a unique contribution to the literature by showing that smartphone
25 delivery of CBT-I is a promising format in which to deliver this gold-standard intervention.
26
27 This is the first study that we are aware of that has evaluated app-delivered CBT-I in young
28
29 people [21], and only the second study that has tested mobile phone delivery of CBT-I, the
30
31 first study being conducted in adults, with positive effects on sleep outcomes [48]. Using
32
33 smartphones to deliver interventions such as this offer a myriad of advantages, including
34
35 immediate connectivity to automated interactive applications that can be accessed anytime,
36
37 anywhere. Sleep Ninja has been developed so that it does not rely on internet access for use,
38
39 which is likely to be important to young people who may have limited data plans, and
40
41 individuals who do not have optimal internet coverage. Not requiring internet coverage to
42
43 access digital programs represents a new wave of flexibility in the delivery of health
44
45 interventions and the automated nature of the intervention means it can be delivered without
46
47 professional support. It is notable that we had a 72% retention rate, which is at the upper level
48
49 of that detected for digital interventions which has been shown in a meta-analysis to range
50
51 between 43-99% [49], with retention rates typically lower in non-supported interventions
52
53
54
55
56
57
58 [50].
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

There are several study limitations that need to be considered. First, participants in this trial were required to have relatively minor levels of insomnia symptoms for study entry. This decision was made by balancing inclusiveness against the requirement of sleep disturbance to ensure participants were motivated to use the app. Moreover, given the study focus was on feasibility and acceptability, we felt it would be prudent to establish these factors before targeting a more severe participant group. That said, while we set the threshold for entry relatively low, both the mean and the median converged on an ISI score of just above 14, indicating that participants were at the junction between having subthreshold and clinical levels of insomnia symptoms (cut-off score is 15). Therefore, it is relatively unlikely that there would have been a floor effect as the data showed there was sufficient room to detect symptom-improvement. The high mean symptom level also suggests that the results may generalise to a group with clinical levels of insomnia. Second, we did not include a control group. Again, as the study goal was to establish feasibility and acceptability, it was not necessary to include a control group for this purpose. However, this design is not able to attribute causality to the intervention and a controlled study is now needed. Finally, this study relied exclusively on subjectively reported sleep outcomes. Although objective measures such as polysomnography provide the most accurate way to assess sleep, there is evidence that subjectively measured sleep variables could be more closely associated with functional outcomes [51]. Moreover, subjectively experienced sleep quality and parameters have consistently shown to be strongly associated with psychological wellbeing [52, 53], suggesting that perception of sleep is as important, if not more so, than objective measures. We are currently investigating how inbuilt smartphone sensors such as the accelerometer might be used to provide a more objective estimate of sleep.

56
57
58
59
60

This study provides preliminary evidence supporting the feasibility, acceptability and effects of a fully-automated app that targets adolescent sleep difficulties. The Sleep Ninja

1
2
3 intervention shows promise both as a sleep-focused intervention, but also potentially to
4
5 reduce risk for depression.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. Example screens from the Sleep Ninja app. From left: Homescreen, Training Session Access and Progress Record, Tracking and Bedtime Setting, More Information

Figure 2. Participant Flow

For peer review only

Acknowledgements

Sincere thanks to Tobias Reid, Jacinto Santamaria and Cesar Anonuevo for their tireless IT support on this project. Thanks also to Dean Winder for assistance in recruitment. The authors would also like to thank the participants and their families for their enthusiasm in this project. AW-S is supported by a NSW Health Early-to-Mid Career Fellowship, MT is supported by an early-career NHMRC Fellowship #1138710 and HC is supported by NHMRC Fellowship #1056964.

Author Statement

AW-S, BO, MT and HC conceived of the study and the trial design. AW-S designed the study with input from all authors, and oversaw the management of the trial. LJ led trial recruitment, managed the day-to-day running of the trial and conducted the participant interviews. QW conducted the analyses with assistance from AW-S and LJ. All authors contributed to the preparation of the manuscript.

Competing Interests Statement

No competing interests to declare.

Funding Statement

This work was supported by a grant awarded to the Black Dog Institute by the Corella Foundation.

Data Sharing Statement: No additional unpublished data from this study is publicly available.

References

1. American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed. ed.). Washington DC: Author.
2. Ohayon, M. M., Roberts, R. E., Zulley, J., Smirne, S., & Priest, R. G. (2000). Prevalence and patterns of problematic sleep among older adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39(12), 1549-1556. doi:10.1097/00004583-200012000-00019
3. Carskadon, M. A. (1990). Patterns of sleep and sleepiness in adolescents. *Pediatrician*, 17(1), 5-12.
4. Liu, X., Buysse, D. J., Gentzler, A. L., Kiss, E., Mayer, L., Kapornai, K., ... Kovacs, M. (2007). Insomnia and hypersomnia associated with depressive phenomenology and comorbidity in childhood depression. *Sleep*, 30(1), 83-90.
5. Baglioni, C., Battagliese, G., Feige, B., Spiegelhalder, K., Nissen, C., Voderholzer, U., ... Riemann, D. (2011). Insomnia as a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *Journal of Affective Disorders*, 135(1-3), 10-19. doi:10.1016/j.jad.2011.01.011
6. Franzen, P. L., & Buysse, D. J. (2008). Sleep disturbances and depression: Risk relationships for subsequent depression and therapeutic implications. *Dialogues in Clinical Neuroscience*, 10(4), 473-481.
7. Riemann, D., & Voderholzer, U. (2003). Primary insomnia: A risk factor to develop depression? *Journal of Affective Disorders*, 76(1-3), 255-259.
8. Ashworth, D. K., Sletten, T. L., Junge, M., Simpson, K., Clarke, D., Cunnington, D., & Rajaratnam, S. M. (2015). A randomized controlled trial of cognitive behavioral therapy for insomnia: An effective treatment for comorbid insomnia and depression. *Journal of Counselling Psychology*, 62(2), 115-123. doi:10.1037/cou0000059

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
9. Manber, R., Edinger, J. D., Gress, J. L., San Pedro-Salcedo, M. G., Kuo, T. F., & Kalista, T. (2008). Cognitive behavioral therapy for insomnia enhances depression outcome in patients with comorbid major depressive disorder and insomnia. *Sleep*, *31*(4), 489-495.
 10. Wagley, J. N., Rybarczyk, B., Nay, W. T., Danish, S., & Lund, H. G. (2013). Effectiveness of abbreviated CBT for insomnia in psychiatric outpatients: Sleep and depression outcomes. *Journal of Clinical Psychology*, *69*(10), 1043-1055. doi:10.1002/jclp.21927
 11. Batterham, P. J., Christensen, H., Mackinnon, A. J., Gosling, J. A., Thorndike, F. P., Ritterband, L. M., ... Griffiths, K. M. (2017). Trajectories of change and long-term outcomes in a randomised controlled trial of internet-based insomnia treatment to prevent depression. *BJPsych Open*, *3*(5), 228-235. doi:10.1192/bjpo.bp.117.005231
 12. Christensen, H., Batterham, P. J., Gosling, J. A., Ritterband, L. M., Griffiths, K. M., Thorndike, F. P., ... Mackinnon, A. J. (2016). Effectiveness of an online insomnia program (SHUTi) for prevention of depressive episodes (the GoodNight Study): A randomised controlled trial. *Lancet Psychiatry*, *3*(4), 333-341. doi:10.1016/s2215-0366(15)00536-2
 13. Blake, M., Waloszek, J. M., Schwartz, O., Raniti, M., Simmons, J. G., Blake, L., ... Allen, N. B. (2016). The SENSE study: Post intervention effects of a randomized controlled trial of a cognitive-behavioral and mindfulness-based group sleep improvement intervention among at-risk adolescents. *Journal of Consulting and Clinical Psychology*, *84*(12), 1039-1051. doi:10.1037/ccp0000142
 14. Waloszek, J. M., Schwartz, O., Simmons, J. G., Blake, M., Blake, L., Murray, G., ... Allen, N. B. (2015). The SENSE Study (Sleep and Education: learning New Skills Early): A community cognitive-behavioural therapy and mindfulness-based sleep

- 1
2
3 intervention to prevent depression and improve cardiac health in adolescence. *BMC*
4
5 *Psychology*, 3, 39. doi:10.1186/s40359-015-0096-x
6
7
- 8 15. de Bruin, E. J., Bogels, S. M., Oort, F. J., & Meijer, A. M. (2018). Improvement of
9
10 adolescent psychopathology after insomnia treatment: Results from a randomized
11
12 controlled trial over 1 year. *The Journal of Child Psychology and Psychiatry*, 59(5),
13
14 509-522. doi:10.1111/jcpp.12834.
15
16
- 17 16. de Bruin, E. J., Bogels, S. M., Oort, F. J., & Meijer, A. M. (2015). Efficacy of
18
19 cognitive behavioral therapy for insomnia in adolescents: A randomized controlled
20
21 trial with internet therapy, group therapy and a waiting list condition. *Sleep*, 38(12),
22
23 1913-1926. doi:10.5665/sleep.5240
24
25
- 26 17. de Bruin, E. J., Oort, F. J., Bogels, S. M., & Meijer, A. M. (2014). Efficacy of internet
27
28 and group-administered cognitive behavioral therapy for insomnia in adolescents: A
29
30 pilot study. *Behavioral Sleep Medicine*, 12(3), 235-254.
31
32 doi:10.1080/15402002.2013.784703
33
34
- 35 18. Okajima, I., Komada, Y., & Inoue, Y. (2011). A meta-analysis on the treatment
36
37 effectiveness of cognitive behavioral therapy for primary insomnia. *Sleep and*
38
39 *Biological Rhythms*, 9(1), 24-34. doi:10.1111/j.1479-8425.2010.00481.x
40
41
- 42 19. Zachariae, R., Lyby, M. S., Ritterband, L. M., & O'Toole, M. S. (2016). Efficacy of
43
44 internet-delivered cognitive-behavioral therapy for insomnia - A systematic review
45
46 and meta-analysis of randomized controlled trials. *Sleep Medicine Reviews*, 30, 1-10.
47
48 doi:10.1016/j.smrv.2015.10.004
49
50
- 51 20. Gulliver, A., Griffiths, K. M., & Christensen, H. (2010). Perceived barriers and
52
53 facilitators to mental health help-seeking in young people: A systematic review. *BMC*
54
55 *Psychiatry*, 10, 113-113. doi:10.1186/1471-244X-10-113
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
21. Werner-Seidler, A., Johnston, L., & Christensen, H. (2018). Digitally-delivered cognitive-behavioural therapy for youth insomnia: A systematic review. *Internet Interventions, 11*, 71-78. doi:10.1016/j.invent.2018.01.007
22. Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2008). Developing and evaluating complex interventions: The new Medical Research Council guidance. *BMJ, 337*, a1655. doi:10.1136/bmj.a1655
23. Gitlin, L. N. (2013). Introducing a new intervention: An overview of research phases and common challenges. *American Journal of Occupational Therapy, 67*(2), 177-184. doi:10.5014/ajot.2013.006742
24. Bastien, C. H., Vallieres, A., & Morin, C. M. (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine, 2*(4), 297-307.
25. Chung, K. F., Kan, K. K.-K., & Yeung, W.-F. (2011). Assessing insomnia in adolescents: Comparison of Insomnia Severity Index, Athens Insomnia Scale and Sleep Quality Index. *Sleep Medicine, 12*(5), 463-470.
26. Chahoud, M., Chahine, R., Salameh, P., & Sauleau, E. A. (2017). Reliability, factor analysis and internal consistency calculation of the Insomnia Severity Index (ISI) in French and in English among Lebanese adolescents. *eNeurologicalSci, 7*, 9-14. doi:10.1016/j.ensci.2017.03.003
27. Alvaro, P.K., Roberts, R.M., & Harris, J.K. (2014). The independent relationships between insomnia, depression, subtypes of anxiety and chronotype during adolescence. *Sleep Medicine, 15*, 934-941. doi: 10.1016/j.sleep.2014.03.019
28. de la Vega, R., Tome-Pires, C., Sole, E., Racine, M., Castarlenas, E., Jensen, M. P., & Miro, J. (2015). The Pittsburgh Sleep Quality Index: Validity and factor structure in young people. *Psychological Assessment, 27*(4), e22-27. doi:10.1037/pas0000128

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
29. Cole, J. C., Motivala, S. J., Buysse, D. J., Oxman, M. N., Levin, M. J., & Irwin, M. R. (2006). Validation of a 3-factor scoring model for the Pittsburgh Sleep Quality Index in older adults. *Sleep*, *29*(1), 112.
30. Guo, S., Sun, W., Liu, C., & Wu, S. (2016). Structural validity of the Pittsburgh Sleep Quality Index in Chinese undergraduate students. *Frontiers in Psychology*, *7*, 1126. doi:10.3389/fpsyg.2016.01126
31. Johnson, J.G., Harris, E.S., Spitzer, R.L. & Williams, J.B.W. (2002). The Patient Health Questionnaire for Adolescents: Validation of an instrument for the assessment of mental disorders among adolescent primary care patients. *Journal of Adolescent Health*, *30*, 196–204.
32. Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, *16*(9), 606-613. doi:10.1046/j.1525-1497.2001.016009606.x
33. Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B. W., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, *114*(1), 163-173. doi:10.1016/j.jad.2008.06.026
34. Spitzer, R. L., Kroenke, K., Williams, J. B., & Lowe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, *166*(10), 1092-1097. doi:10.1001/archinte.166.10.1092
35. Tiirikainen, K., Haravouri, H., Ranta, K., Kaltiala-Heino, R. & Marttunen, M. (2019). Psychometric properties of the 7-item Generalized Anxiety Disorder Scale (GAD-7) in a large representative sample of Finnish adolescents. *Psychiatry Research*, *272*, 30-35.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
36. Thorndike, F. P., Saylor, D. K., Bailey, E. T., Gonder-Frederick, L., Morin, C. M., & Ritterband, L. M. (2008). Development and perceived utility and impact of an internet intervention for insomnia. *E-journal of Applied Psychology : Clinical and Social Issues*, 4(2), 32-42.
37. Ritterband, L. M., Ardan, K., Thorndike, F. P., Magee, J. C., Cox, D. J., Sutphen, J. L., & Borowitz, S. M. (2008). Real world use of an Internet intervention for pediatric encopresis. *Journal of Medical Internet Research*, 10(2), e16. doi:10.2196/jmir.1081.
38. Carney, C. E., Buysse, D. J., Ancoli-Israel, S., Edinger, J. D., Krystal, A. D., Lichstein, K. L., & Morin, C. M. (2012). The Consensus Sleep Diary: Standardizing prospective sleep self-monitoring. *Sleep*, 35(2), 287-302. doi:10.5665/sleep.1642
39. Werner-Seidler, A., O'Dea, B., Shand, F., Johnston, L., Frayne, A., Fogarty, A. S., & Christensen, H. (2017). A smartphone app for adolescents with sleep disturbance: Development of the Sleep Ninja. *JMIR Mental Health*, 4(3), e28. doi:10.2196/mental.7614
40. Harvey, A. G. (2009). A transdiagnostic approach to treating sleep disturbance in psychiatric disorders. *Cognitive Behaviour Therapy*, 38 Suppl 1, 35-42. doi:10.1080/16506070903033825
41. Cain, N. & Gradisar, M. (2010). Electronic media use and sleep in school-aged children and adolescents: A review. *Sleep Medicine*, 11, 735-742. doi:10.1016/j.sleep.2010.02.006
42. Hysing, M., Pallesen, S., Stormark, K.M., Jakobsen, R., Lundervold, A.J., & Sivertsen, B. (2014). Sleep and use of electronic devices in adolescence: Results from a large population-based study. *BMJ Open*, 5, e006748. doi:10.1136/bmjopen-2014-006748
43. Kline, R. B. (2011). *Methodology in the Social Sciences. Principles and practice of structural equation modeling* (3rd ed.). New York, NY, US: Guilford Press.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
44. Horsch, C. H., Lancee, J., Jan Beun, R., Neerincx, M. A. & Brinkman, W. P. (2015). Adherence to technology-mediated insomnia treatment: a meta-analysis, interviews and focus groups. *Journal of Medical Internet Research*, *17*(9), e214. doi: 10.2196/jmir.4115
45. Wheaton, A. G., Chapman, D. P. & Croft. (2016). School start times, sleep, behavioral, health, and academic outcomes: a review of the literature. *Journal of School Health*, *86*(5), 363-381. doi: 10.1111/josh.12388
46. Watson, N. F., Martine, J. L., Wise, K. A., Kirsch, D. B. Kristo, D.B.... & Chervin, R. D. (2017). Delaying middle school and high school start times promotes student health and performance. An American Academy of Sleep Medicine Position Statement, *Journal of Clinical Sleep Medicine*, *13*(4), 623-623. doi: 10.5664/jcsm.6558
47. Horowitz, J. L., Garber, J., Ciesla, J. A., Young, J. F., & Mufson, L. (2007). Prevention of depressive symptoms in adolescents: A randomized trial of cognitive-behavioral and interpersonal prevention programs. *Journal of Consulting and Clinical Psychology*, *75*(5), 693-706. doi:10.1037/0022-006x.75.5.693
48. Horsch, C. H., Lancee, J., Griffioen-Both, F., Spruit, S., Fitrianie, S., Neerincx, M. A., ... Brinkman, W. P. (2017). Mobile phone-delivered cognitive behavioral therapy for insomnia: A randomized waitlist controlled trial. *Journal of Medical Internet Research*, *19*(4), e70. doi:10.2196/jmir.6524
49. Christensen, H., Griffiths, K. M., & Farrer, L. (2009). Adherence in internet interventions for anxiety and depression: Systematic review. *Journal of Medical Internet Research*, *11*(2), e13.

- 1
2
3 50. Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized
4
5 psychological treatments for adult depression: A meta-analysis. *Cognitive Behaviour*
6
7 *Therapy, 38*(4), 196-205. doi:10.1080/16506070903318960
8
9
- 10 51. Wilson, D. L., Fung, A., Walker, S. P., & Barnes, M. (2013). Subjective reports versus
11
12 objective measurement of sleep latency and sleep duration in pregnancy. *Behavioral*
13
14 *Sleep Medicine, 11*(3), 207-221. doi:10.1080/15402002.2012.670674
15
16
- 17 52. Fuligni, A. J., & Hardway, C. (2006). Daily variation in adolescents' sleep, activities,
18
19 and psychological well-being. *Journal of Research on Adolescence, 16*(3), 353-378.
20
21 doi:10.1111/j.1532-7795.2006.00498.x
22
23
- 24 53. Lund, H. G., Reider, B. D., Whiting, A. B., & Prichard, J. R. (2010). Sleep patterns
25
26 and predictors of disturbed sleep in a large population of college students. *Journal of*
27
28 *Adolescent Health, 46*(2), 124-132. doi:10.1016/j.jadohealth.2009.06.016
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Example screens from the Sleep Ninja app. From left: Homescreen, Training Session Access and Progress Record, Tracking and Bedtime Setting, More Information

139x238mm (300 x 300 DPI)




Figure 2. Participant Flow

187x238mm (300 x 300 DPI)

BMJ Open

A pilot evaluation of the Sleep Ninja – a smartphone-application for adolescent insomnia symptoms

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-026502.R2
Article Type:	Research
Date Submitted by the Author:	21-Mar-2019
Complete List of Authors:	Werner-Seidler, Aliza ; University of New South Wales, Black Dog Institute Wong, Quincy; University of New South Wales, Black Dog Institute; Western Sydney University School of Science and Health, School of Social Sciences and Psychology Johnston, Lara; University of New South Wales, Black Dog Institute O'Dea, B; University of New South Wales, Black Dog Institute Torok, Michelle; University of New South Wales, Black Dog Institute Christensen, Helen; University of New South Wales - Randwick Campus, Black Dog Institute
Primary Subject Heading:	Mental health
Secondary Subject Heading:	Mental health
Keywords:	insomnia, adolescent mental health, cognitive-behaviour therapy for insomnia, ehealth, Depression & mood disorders < PSYCHIATRY

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

A pilot evaluation of the Sleep Ninja – a smartphone-application for adolescent insomnia symptoms

Aliza Werner-Seidler¹; a.werner-seidler@blackdog.org.au

Quincy Wong^{1, 2}; quincy.wong@blackdog.org.au

Lara Johnston¹; l.johnston@blackdog.org.au

Bridianne O’Dea¹; b.odea@blackdog.org.au

Michelle Torok¹; m.torok@blackdog.org.au

Helen Christensen¹; h.christensen@blackdog.org.au

35
36
37
38
39
40
41
42
43
44
45

Affiliations:

¹Black Dog Institute, University of New South Wales, Sydney, Australia

²School of Social Sciences and Psychology, Western Sydney University, Sydney, NSW, Australia.

46
47
48
49
50

Corresponding author: Dr Aliza Werner-Seidler; Black Dog Institute, Hospital Road, Randwick, New South Wales, Australia; a.werner-seidler@blackdog.org.au; 02 9382 380.

51
52
53

Word Count: 5102

54
55
56
57
58
59
60

Keywords: Insomnia, adolescent mental health, cognitive-behaviour therapy for insomnia, eHealth

Abstract

Objectives: The aim of this study was to test the feasibility, acceptability and preliminary effects of a recently developed smartphone application, Sleep Ninja, for adolescent sleep difficulties.

Setting: The study was conducted online with Australian individuals recruited through the community.

Participants: Participants were 50 young people aged 12-16 years with sleep difficulties.

Design: A single-arm pre-post design was used to evaluate feasibility, acceptability and sleep and mental health variables at baseline and post-intervention.

Intervention: Cognitive Behaviour Therapy for Insomnia (CBT-I) informed the development of the Sleep Ninja. The core strategies covered by the app are psychoeducation, stimulus-control, sleep hygiene, and sleep-related cognitive therapy. It includes six training sessions (lessons), a sleep tracking function, recommended bedtimes based on sleep guidelines, reminders to start a wind-down routine each night, a series of sleep tips, and general information about sleep. Users progress through each training session and conclude the six-week program with a black belt in sleep.

Outcome measures: Feasibility was evaluated based on consent rates, adherence and attrition, acceptability was assessed using questionnaires and a post-study interview, and sleep, depression and anxiety variables were assessed at baseline and post-intervention.

Results: Data indicated that the Sleep Ninja is a feasible intervention and is acceptable to young people. Findings showed there were significant improvements on sleep variables including insomnia (within-group effect size $d=-0.90$), sleep quality ($d=-0.46$), depression ($d=-0.36$) and anxiety ($d=-0.41$).

Conclusions: The Sleep Ninja is a promising intervention that could assist adolescents who experience sleep difficulties. A follow-up randomised controlled trial is now warranted.

Trial registration: Australian New Zealand Clinical Trials Registry

(#ACTRN12617000141347).

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Article Summary

Strengths and limitations of this study

- This is the first study to evaluate app-delivered Cognitive Behavioural Therapy for Insomnia in adolescents with sleep difficulties.
- The intervention being tested, Sleep Ninja, was developed with input from young people, is fully automated and does not require internet coverage to function.
- The evaluation included measures of feasibility and acceptability as well as detailed semi-structured interviews about participants' experience with the app.
- As a preliminary study, this study did not include a control group.

1
2
3 Clinical insomnia is a sleep disturbance characterised by difficulty falling asleep,
4 staying asleep or waking up too early, with associated daytime impairment [1]. It effects
5 approximately 4% of adolescents [2], however sub-threshold symptoms are common, with
6 approximately 25% of young people reporting some degree of sleep disturbance [2, 3].
7
8
9

10
11
12 Depression and insomnia are closely linked, with comorbidity levels as high as 73%
13 in young people [4]. Insomnia is not only a symptom of depression, but is a common
14 precursor, with high quality longitudinal data having established insomnia as an independent
15 risk factor for depression onset [5, 6, 7]. For example, a recent meta-analysis found that
16 insomnia was associated with a greater than two-fold increase in depression risk [5].
17
18 Although depression has multiple causes and maintaining factors that go beyond the presence
19 of sleep problems, the literature suggests that sleep plays an important role, and targeting
20 sleep in the context of depression may have wide-reaching benefits [6].
21
22
23
24
25
26
27
28
29

30
31 There is emerging evidence that addressing insomnia in individuals with concurrent
32 insomnia and depression improves both sleep and depression outcomes [8, 9, 10]. This
33 suggests there may be value in targeting sleep to improve insomnia symptoms, with potential
34 downstream effects on depression. To our knowledge, there have been three studies testing
35 the hypothesis that targeting insomnia can prevent depressive symptoms. In an adult study,
36 insomnia treatment led to a reduction in depression following the intervention and at 6 and
37 18-month follow-up, relative to an active control group [11, 12]. In a youth study, a face-to-
38 face insomnia intervention was delivered to secondary school students and results showed
39 improvements on sleep and anxiety outcomes, but not symptoms of depression [13]. Data
40 from the two-year follow-up from this study has not yet been published [14]. In a second
41 youth study, a sleep intervention delivered either in group format or digitally led to decreased
42 depressive symptoms at both 2 and 12-month follow-up, an effect that was mediated by
43 improvements in insomnia [15].
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The gold standard treatment for insomnia is cognitive behaviour therapy for insomnia
4 (CBT-I; Australasian Sleep Association, American College of Physicians), and there is
5
6 accumulating evidence to support the use of digitally delivered CBT-I in both adults and
7
8 young people [16, 17, 18, 19]. Delivering sleep interventions via digital formats may be
9
10 particularly well-suited to young people, with adolescents showing a strong preference (97%)
11
12 for digital delivery when given the choice between face-to-face and digital CBT-I [17]. This
13
14 preference may in part be explained by the fact that young people are reluctant to seek help
15
16 for psychological issues, for reasons that include stigma and a preference to manage the
17
18 problem themselves [20]. Sleep is typically less stigmatised than disorders like depression,
19
20 suggesting that it may be more appealing to adolescents. Currently, there are no digital CBT-I
21
22 programs that are commercially available for youth [21]. To overall objective of this study
23
24 was to evaluate a newly-developed digital CBT-I program for adolescents with sleep
25
26 difficulties.
27
28
29
30
31
32

33 The aim of this pilot study was to examine the acceptability, feasibility and
34
35 preliminary effects of an intervention (Sleep Ninja) delivered to adolescents via smartphones.
36
37 In line with the guidelines on the development of behavioural interventions [22, 23], the
38
39 primary purpose of this study was to investigate recruitment rates, uptake, intervention
40
41 completion, reasons for non-adherence, and participant retention. The secondary aim was to
42
43 use both quantitative and qualitative methods to determine the acceptability of the app among
44
45 young people with sleep difficulties and allow for the refinement of the intervention prior to a
46
47 formal randomised evaluation. A final aim was to examine the impact of the Sleep Ninja app
48
49 on sleep outcomes and mental health symptoms. We used a single-arm, pre-post design to
50
51 address these aims. It was hypothesised that the app would be a feasible modality in which to
52
53 deliver the automated sleep intervention, as measured by uptake, completion and retention
54
55
56
57
58
59
60

1
2
3 rates, that the app would be acceptable to young people, and that its use would be associated
4
5 with improvement in sleep and mental health symptoms.
6
7
8
9

10 **Method**

11
12 This trial was prospectively registered on the Australian New Zealand Clinical Trials
13 Registry (#ACTRN12617000141347).
14
15

16 **Participants**

17
18 Fifty participants were recruited via media and social media channels, including the
19 Black Dog Institute's website and paid Facebook advertisements that targeted potential
20 participants and their parents between April-June, 2017. A sample size of 50 was selected in
21 order to successfully meet the study's feasibility and acceptability aims. Inclusion criteria
22 were: aged 12-16 years, presence of at least mild insomnia, operationalised by endorsement
23 of at least one of the following symptoms over the preceding two-week period: difficulty
24 falling asleep, difficulty staying asleep or waking up too early. These items are the first three
25 questions on the Insomnia Severity Index [24], and were chosen to include a participant
26 group with at least mild levels of insomnia. For study inclusion, participants also needed to
27 own a smartphone running iOS or Android, have a valid email address, access to the internet,
28 and be able to provide personal and parental consent.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 **Measures**

48
49 **Insomnia Severity Index (ISI).** The ISI is a psychometrically sound, seven-item self-
50 report measure of insomnia symptoms over the previous two weeks [24, 25]. Responses are
51 reported on a Likert scale from 0 to 4, producing total scores of 0 to 28 [24, 25]. Cut-off
52 scores are as follows: 0-7 reflects no clinically significant insomnia, 8-14 indicates
53 subthreshold insomnia, 15-21 suggests moderate severity insomnia, and 22-28 indicates
54
55
56
57
58
59
60

1
2
3 severe insomnia [24]. The ISI was designed for use in adults but has been widely
4 administered to, and validated in, adolescent samples [25, 26, 27]. In one adolescent
5 validation study, reliability was strong (Cronbach's $\alpha=0.83$), and test-retest reliability was
6 acceptable, $r=.79$ [25].
7
8
9

10
11
12 **Pittsburgh Sleep Quality Index (PSQI).** The PSQI is a widely used self-report 19-
13 item scale that assesses usual sleep habits and experiences over the preceding month and has
14 been validated in adolescent samples, with strong internal consistency ($\alpha=.72$) and test-retest
15 reliability over a 6-week period ($r=.81$) [28]. There are seven sub-scales which are sleep
16 quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of
17 sleeping medications, daytime dysfunction [29]. Each component is scored from 0 (no
18 difficulty) to 3 (severe difficulty), which are summed to obtain a Global PSQI score ranging
19 from 0 to 21 [30].
20
21
22
23
24
25
26
27
28
29

30
31 **Patient Health Questionnaire – Adolescent Version (PHQ-A).** The PHQ-A
32 assesses depressive symptoms in the preceding two weeks in adolescents, and has been
33 adapted from the widely used PHQ-9 designed for adults [31]. This measure has excellent
34 psychometric properties, including $\alpha=0.89$, and test-retest reliability of $r=0.84$ [32]. In this
35 study, we used the 8-item version in which the questions are identical to those asked in the
36 PHQ-9 with the exclusion of the last item which asks about suicide and is comparable to the
37 PHQ-9 in terms of diagnosing depressive disorders [32, 33]. Each item is scored on a 4-point
38 scale and summed together to form a total depression score ranging from 0 to 24. Scores
39 correspond to the following cut-offs: 0-9 indicates minimal symptoms, 10-14 indicates mild
40 symptoms, 15-19 reflects moderate symptoms, and 20-24 is indicative of severe depression
41 [31]. The PHQ-A has demonstrated good sensitivity (73%) and high specificity (94%) for
42 major depressive disorder [31].
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Generalised Anxiety Disorder 7-item (GAD-7).** The GAD-7 evaluates symptoms of
4 generalised anxiety disorder [34]. All items are scored on a scale from 0 (not at all) to 3
5 (nearly every day). The scores on each item are summed together to derive a total score,
6 ranging from 0 to 21 of which 0-4 indicates minimal anxiety, 5-9 mild anxiety, 10-14
7 moderate anxiety, and 15-21 severe anxiety [34]. The GAD-7 has good sensitivity (89%) and
8 specificity (82%) for GAD scores >10 [34]. The measure has also been validated in
9 adolescent populations with Cronbach's $\alpha=0.90$ and high convergent and discriminant
10 validity [35].
11
12
13
14
15
16
17
18
19
20

21 **Expectations of Success.** A four-item scale was developed for this study to assess
22 participants' motivation and expectations for improving their sleep with an app (e.g., *I am*
23 *confident that people could learn skills for improving sleep from an app*). The four items
24 assessed perceived confidence, importance, usefulness and readiness to change. The
25 Expectation of Success measure was scored on a five-point scale and total scores were
26 computed by summing each item, ranging from 0 to 16. Higher scores on this scale indicate
27 greater confidence and readiness to target sleep using a smartphone app.
28
29
30
31
32
33
34
35
36

37 **Acceptability of the Intervention.** The Acceptability of the Intervention scale is a
38 seven-item measure that was developed by the research team to assess participants' attitudes
39 and behaviours associated with using the app (e.g., *How much did you learn from the app* and
40 *Would you recommend this program to others?*). This measure was informed by similar
41 acceptability measures commonly used in the field [36]. Each item was designed to assess a
42 different domain. The first four domains to be assessed were app completion, ease of use,
43 amount learnt and usefulness, with each being scored on an ordinal scale from 0-3. The final
44 three items assessed behaviour change, whether the participant would use an app like this in
45 the future, and whether they would recommend it to a friend, and were scored dichotomously
46 as either yes or no, with an option for participants to describe the nature of their behaviour
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 change if yes. As each question assessed a different domain, item scores were considered
4
5 separately.
6

7
8 **Reasons for Non-Adherence.** The Reasons for Non-Adherence measure is a 23-item
9
10 scale that was adapted from a previous measure [37] to assess the degree to which different
11
12 reasons impacted on participants' use of the app. There are four domains assessed:
13
14 phone/internet/technical issues (*e.g., My phone wasn't working or was having problems*);
15
16 personal issues (*e.g., I didn't think I deserved help*); intervention-general issues (*e.g., I wasn't*
17
18 *convinced the app would be helpful*); and intervention-specific issues (*e.g., There was too*
19
20 *much text to read*). Participants responded on an ordinal scale indicating whether each item
21
22 played no, a little or major part in why they stopped or had difficulty using the app as
23
24 intended. The scores on each item were considered separately.
25
26
27

28
29 **Sleep Diary.** The ten-item Sleep Diary was developed by the research team
30
31 incorporating the questions from the Consensus Sleep Diary [38], with the addition of two
32
33 questions regarding daytime naps and use of sleep medication. Participants answered 10
34
35 questions which included bedtime, time taken to fall asleep (sleep onset latency; SOL),
36
37 number and duration of night-time awakenings (number of awakenings; NWAK, duration of
38
39 wakefulness after sleep onset; WASO), time of final awakening, time participants got out of
40
41 bed for the day, subjective sleep quality, how refreshed participants felt upon awakening,
42
43 duration of any daytime naps and use of sleep medication. Sleep diaries were completed
44
45 electronically with pre-set categories from which users selected responses from a drop-down
46
47 menu. A clock scroller was used to enter the time and/or duration of all sleep-related
48
49 activities and all times were entered in 12-hour format to minimise errors associated with 24-
50
51 hour time. Restrictions were set to ensure participants could not enter a wake time earlier than
52
53 bedtime and visa versa. All questions required answers for the sleep diary to be submitted.
54
55
56 From the sleep diary we calculated time between waking in the morning and getting out of
57
58
59
60

1
2
3 bed, time in bed (TIB), total sleep time (TST; calculated by subtracting SOL, WASO and
4 time between waking and getting up in the morning, from TIB) and sleep efficiency (SE;
5 calculated by taking the percentage of TST/TIB).
6
7
8
9

10 **Post-Study Interview.** After study completion, participants were invited to attend a
11 face-to-face or telephone interview to provide feedback on their experience. Interviews were
12 semi-structured and explored participants' opinions about the study in general, and
13 specifically in relation to the intervention. Questions were open-ended, and flexible enough to
14 explore ideas that were raised during each interview. Interviews were audio recorded and
15 then transcribed verbatim by the interviewer. The interview content was pragmatically coded
16 into relevant themes by the same researcher, with oversight and guidance provided by the
17 research team.
18
19
20
21
22
23
24
25
26
27

28 **Intervention – ‘Sleep Ninja’**

29
30 The Sleep Ninja app was derived from CBT-I and developed by our team, as a fully-
31 automated smartphone app. A participatory design process was used whereby young people
32 contributed to the content, functionality and accessibility/user experience of the app through a
33 series of focus groups [39]. The core strategies included in the app were: psychoeducation,
34 stimulus-control, sleep hygiene and sleep-focused cognitive therapy. Sleep restriction, which
35 aims to increase sleep efficiency by reducing the amount of time spent in bed, was
36 deliberately omitted because some support (parental and/or professional) is likely to be
37 required to successfully implement sleep restriction, particularly in young people. Although
38 sleep restriction did not comprise part of the app, there was instead a focus on the importance
39 or regular sleep-wake cycles. The app teaches users about the importance of consistent sleep
40 and wake times, and recommended bedtimes are calculated based on the time they need to
41 wake up (according to sleep guidelines). This strategy draws from transdiagnostic approaches
42 to target sleep difficulties that go beyond insomnia (e.g., delayed sleep phase and irregular
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 sleep presentations; [40]) and may therefore be useful to adolescents experiencing a broad
4
5 range of sleep difficulties.
6

7
8 The structure of the Sleep Ninja app includes six training sessions (lessons), a sleep
9
10 tracking function, recommended bedtimes based on sleep guidelines, reminders to start a
11
12 wind-down routine each night, a series of sleep tips, and general information about sleep. The
13
14 home screen has 3 options: Train, Track, and More (see Figure 1). Users complete training
15
16 sessions which are delivered through a chat-bot format where the sleep ninja essentially acts
17
18 as a sleep coach. Training sessions take approximately 5-10 minutes to complete, and cover:
19
20 (i) psychoeducation, information about circadian rhythms and the importance of keeping
21
22 regular sleep schedules; (ii) stimulus control, the value of only going to sleep when tired, and
23
24 strategies that can be used at night when having trouble sleeping; (iii) basic sleep hygiene
25
26 such as avoiding caffeine and stimulating activity in the evenings, suggestions for daytime
27
28 activities to promote night-time sleep (e.g., exercise, no napping); (iv) identifying and
29
30 planning for high-risk situations, how to get back on track after a late-night or sleep in; (v)
31
32 cognitive therapy including how to deal with unhelpful thoughts that can prevent falling
33
34 asleep as well as sleep-related cognitive distortions cognitions and; (vi) a final review
35
36 session which summarises all of the material contained in the app. The user interacts with the
37
38 app through a forced choice chat-bot format which is responsive to the input of the user,
39
40 meaning it personalises information and recommendations based on the selections and sleep
41
42 profile recorded by the participant. Users level up and reach their next “belt” by completing
43
44 one training session and tracking their sleep for 3 nights (out of a 7-night period). As there
45
46 were six training sessions to complete, the app was made available for six weeks (42 days)
47
48 before it locked. Users finish the program with a black belt in sleep.
49
50
51
52
53
54

55
56 It is notable that evidence suggests that the use of screens at bedtime interferes with
57
58 sleep [41]. Drawing on data from large epidemiological studies suggesting that refraining
59
60

1
2
3 from screen use for the hour prior to bedtime alleviates potential interference from screens
4
5 [42], this app has been designed to be used during the day. Users receive a prompt one hour
6
7 before bed (calculated according to sleep guidelines and their wake-up time) to commence
8
9 their pre-bed routine and are encouraged to stop using electronic devices after this time. In
10
11 fact, part of the cognitive component of the app is to educate and challenge beliefs about the
12
13 importance of night time phone use in order to promote healthy sleep habits. We expect these
14
15 factors to mitigate the risk that smartphone use in this context will contribute to poor sleep.
16
17

18 19 **Procedure**

20
21 All procedures were approved by the University of New South Wales Human
22
23 Research Ethics Committee (HC#16702). Participants were encouraged to download consent
24
25 forms if they met the eligibility criteria listed on the study website and submit this directly to
26
27 the research team, once completed. Those who provided written informed consent and that of
28
29 a parent or guardian were then enrolled in the trial and invited to complete the screening
30
31 questions to verify study eligibility, before completing baseline questionnaires which
32
33 included: demographics, ISI, PSQI, PHQ-A, GAD-7 and Expectation of Success. Participants
34
35 could then access the first day of the online seven-day sleep diary. Another diary entry
36
37 became available each day for the following six days and participants were reminded to
38
39 complete entries via text-message. At the completion of seven consecutive entries in the
40
41 diary, participants were given access to the Sleep Ninja app on their personal smartphone
42
43 devices. Participants could use the app for six weeks before the post study questionnaire was
44
45 made available, which included the same battery as baseline with the omission of the
46
47 Expectations of Success questionnaire and with the addition of the Acceptability and Reasons
48
49 for Non-Adherence questionnaires. Participants then completed another seven-day sleep
50
51 diary, which was delivered in the same format and schedule as baseline. After the study had
52
53 finished, participants were invited to participate in a face-to-face or telephone interview to
54
55
56
57
58
59
60

1
2
3 provide feedback on their experience of participating in the study. Participants were
4
5 reimbursed for their time with giftcards to the value of \$10 each for completing baseline and
6
7 post-study assessment schedules; \$20 for a telephone interview and \$30 for a face-to-face
8
9 interview.
10

11 **Participant Involvement and Consultation**

12
13
14
15 Prior to this study, a separate group of young people were consulted in a series of
16
17 focus groups to inform the design, features and structure of the app (for more information
18
19 please see [39]). As a feasibility and acceptability study, participants were asked to report on
20
21 their experiences with respect to both the app and the study procedures, via questionnaires
22
23 and an in-depth semi-structured interview. Given that a key objective of this study was to
24
25 assess the acceptability of the Sleep Ninja app, participants' perspectives were of critical
26
27 importance. A one-page lay summary of the study results has been sent to all participants.
28
29

30 **Statistical Analyses**

31
32
33 Statistical significance was set at $\alpha=.05$. Summary scores for sleep diary variables at
34
35 baseline and post-study were obtained by averaging sleep diary entries at baseline, and
36
37 averaging sleep diary entries at post-study, respectively. All questionnaire and sleep diary
38
39 variables were initially screened for excessive skew (>3) or kurtosis (>8 ; [43]). Six sleep
40
41 diary variables did not pass screening and were further scrutinised (baseline: WASO entry;
42
43 post-study: Time fallen asleep, Time in bed after final morning wake up, SOL, TST, SE
44
45 variables). Examination of these six variables revealed each included an entry that was of an
46
47 extreme value (z -scores ranged from $|4.11|$ to $|6.05|$) and a decision was made to remove these
48
49 six values (hence, $n = 47$ for Baseline WASO; $n_s = 28$ for Post-intervention Time fallen
50
51 asleep, Time in bed after final morning wake up, SOL, TST, and SE). Subsequently all
52
53 variables had satisfactory skew and kurtosis.
54
55
56
57
58
59
60

Questionnaire and sleep diary variables were examined using multilevel modelling. This modelling approach handles missing data by incorporating all available data from each subject into the analysis. Given the aims of our study, our interest centred on the main effect of time (i.e., change from baseline- to post). Random effects were modelled for intercept and time. Models were respecified with a random effect for intercept only in cases where there was no variation in individual baseline- to post-study changes. Within-group effect sizes were computed as the modelled mean difference between baseline and post-study divided by the sample standard deviation at baseline.

Results

Baseline Characteristics

See Table 1 for characteristics of the study sample. Participants had a mean age of 13.71 ($SD=1.35$), were spread across school grade, and nearly all were born in Australia and living in the city. Most participants reported difficulty falling asleep, with about half also reporting problems staying asleep or waking up too early, and about a quarter of the sample were receiving treatment for sleep or a mental health problem.

Table 1. Demographic Variables

Characteristics	Sample ($N = 50$)
Age in years, mean (SD , range)	13.71 (1.35, 12-16)
Age in years, n (%)	
12	10 (20.4%)
13	15 (30.6%)
14	7 (14.3%)
15	12 (24.5%)
16	5 (10.2%)
Female, n (%)	33 (66%)
Born in Australia, n (%)	47 (94%)
Live in the city, n (%)	44 (88%)
Sleep problems, n (%)	
Difficulty falling asleep	47 (94%)
Difficulty staying asleep	28 (56%)
Problems waking up too early	28 (56%)
Receiving treatment for sleep or mental health problem, n (%)	13 (26%)

Note. One participant did not indicate their age, so $n = 49$ for age.

Recruitment Rate

There were more than 300 enquires made to the research team about participation in this trial. Of these, 60 individuals indicated eligibility and returned consent forms. Ten of these participants were not enrolled in the trial; four did not meet inclusion criteria and six withdrew prior to the trial. Reasons for withdrawing were: a change of mind ($n=1$), a lack of time ($n=2$), considering participation a chore ($n=1$), confidentiality concerns ($n=1$). One participant did not provide a reason. Therefore, 89% ($n=50$) of the 56 young people who provided consent and met screening criteria continued to trial. See Figure 2 CONSORT diagram for details.

Expectation of Success

Overall, participants were optimistic about using the app, with a mean score of 12.90 ($SD=2.09$) out of a possible 16 points. Every single participant agreed that in principle, people could learn skills for improving sleep from an app, and indicated that they felt that study participation was important. All participants reported that improving their sleep habits were important, with 49% indicating it was 'very important'. Finally, the sample demonstrated their readiness for change with 100% of the sample indicating that they were either moderately ready (16%), ready (43%) or completely ready (41%) to improve their sleep patterns using an app.

Retention

Of the 50 participants in the study sample, 47 (94%) completed the baseline questionnaire and sleep diaries and were invited to download the Sleep Ninja. At post-study, 34 participants completed the post-study battery (72% retention). Participants who had available data at both time-points did not differ significantly from those who only had baseline data on any of the questionnaires or sleep diary measures (all F s < 2.58 , p s $> .115$).

Uptake and Adherence

Forty-five participants (96%) who completed the baseline assessment downloaded the Sleep Ninja. Program usage data indicated that of these, 82% completed the first lesson, 51% completed four of the six lessons, and 33% completed all six. Participants were accurate in their reporting of app use, with approximately 80% of participants indicating that they completed 'most' or 'almost all' of the app.

Acceptability

Survey responses on the Acceptability of the Intervention questionnaire indicated that young people reported that the app was 'easy' or 'very easy' to use (97%). The majority of participants (59%) indicated that they learnt 'a fair bit' from the app, and 28% reported that they learnt 'a great deal', while 12% did not learn very much or almost nothing. Participants found the app to be either 'useful' or 'very useful' (78%), while the remainder (22%) did not find it useful. Most respondents (72%) reported changing their behaviour after using the app, and examples of behaviour change included changes to their pre-bedtime routine (22%), keeping more consistent sleep-wake cycles (65%), getting up earlier in the morning (22%), and restricting the use of their bed for sleep (30%). More than half of the participants reported that they would use this kind of app in the future (56%), and encouragingly, 91% would recommend the Sleep Ninja app to a friend. The degree to which participants found the app useful was positively correlated with module completion, ($r=.35$, $p=.047$), as was the degree to which participants reported learning from the app ($r=.49$, $p=.004$).

The interview mirrored the findings of the questionnaire in terms of acceptability and usefulness. However, there were some aspects of the app that users felt could be improved. Specifically, interviewees expressed a desire for improved explanation of the different app sections and what they needed to do each time they opened the app. Participants commonly reported wanting to be able to personalise their user experience more, including skipping

1
2
3 information they knew, seeking more information around difficult or unfamiliar topics,
4
5 accessing information in different formats (e.g. video/audio), being able to speed up or slow
6
7 down the Sleep Ninja's speech, and being able to update their wind-down activity choices
8
9 and the time the wind-down reminder appeared. Participants expressed a range of views
10
11 about the tone of the Sleep Ninja, with nearly half of the interviewed participants
12
13 commenting favourably on the Sleep Ninja's jokes, with several participants commenting that
14
15 the Sleep Ninja's language was annoying and too childish. There was consensus that the
16
17 Sleep Ninja's language was repetitive and could be improved by cutting out superfluous
18
19 dialogue that was not delivering core intervention strategies. Nearly half of the interviewed
20
21 participants expressed some difficulty in implementing at least one of the Sleep Ninja's
22
23 recommended strategies due to conflicting parental bedtime rules. For instance, the strategies
24
25 to delay bedtime until sleepy and leave the bed/bedroom if unable to get to sleep after more
26
27 than 30 minutes most commonly encountered parental resistance or required modification.
28
29 Numerous interviewees commented on the usefulness of receiving feedback and summaries
30
31 of their logged sleep, however several commented that this could be improved by displaying
32
33 the information in graphs and over time, so that change and improvement is clearer. Overall,
34
35 participants reported that the number of notifications in the app were acceptable, and that
36
37 additional reminders should be sent to notify them of available lessons and after periods of
38
39 inactivity and that these reminders should contain motivational and encouraging messages.
40
41 While most interviewees considered three nights of sleep tracking per belt acceptable, there
42
43 were others who felt fewer nights of tracking would have been better, and others who
44
45 expressed willingness to track more than three nights before levelling up.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reasons for Non-Adherence

Results from this questionnaire indicated that young people were very happy to use an app to receive help for their sleep issues (84%), did not have technical issues with its use (75%), and felt they had the technical skills to use the app (90%). Participants all reported that the material was relevant and conceptually easy to understand. The main reasons participants reported not using the app was that they felt it took too long to work through (53%), there was too much text to read (47%), and that it was too repetitive (59%).

Preliminary Effects

Sleep and Mental Health Questionnaire Outcomes. Table 2 shows the results for the questionnaire measures. As predicted, from baseline- to post-study, there was a significant decrease in insomnia severity measured by the ISI, $\beta=-4.29$, $p<.001$, $d=-0.90$, and sleep quality on the PSQI, $\beta=-1.88$, $p<.001$; $d=-0.46$. Similarly, mental health measures showed a decrease in both depression on the PHQ-A, $\beta=-2.60$, $p<.001$, $d=-0.36$ and anxiety on the GAD-7, $\beta=-2.56$, $p<.001$, $d=-0.41$.

Table 2. Questionnaire Measures

Outcome	Pre-intervention		Post-intervention		Modelled change from pre- to post-intervention	<i>d</i>
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	β [95% CI]	
ISI	50	14.12 (4.75)	34	9.62 (5.23)	-4.29 [-5.63, -2.95]***	-0.90
PSQI	50	10.43 (4.12)	33	8.03 (4.08)	-1.88 [-2.85, -0.90]***	-0.46
PHQ-A	49	13.04 (7.24)	32	9.88 (7.53)	-2.60 [-3.99, -1.22]***	-0.36
GAD-7	49	9.92 (6.19)	32	7.09 (6.13)	-2.56 [-3.59, -1.52]***	-0.41

Note. Raw means (SDs) are presented. Cohen's *d* values are time effects for pre-intervention to post-intervention using the modelled mean difference divided by the sample pre-intervention SD. ISI = Insomnia Severity Index; PSQI = Pittsburgh Sleep Quality Index; PHQ-A = Patient Health Questionnaire modified for Adolescents; GAD-7 = Generalised Anxiety Disorder 7-item.

*** $p < .001$.

Sleep Diary Outcomes. Results for the sleep diary entries are shown in Table 3. At baseline, participants went to sleep, on average at 11:29pm, reported taking an average of one hour and 12 minutes to fall asleep, spent an average of 9 hours and 39 minutes in bed, woke

1
2
3 up an average of 1.47 times, slept for a total of seven hours and 40 minutes, woke up at
4
5 7:27am, and spent approximately 29 minutes awake in bed before getting up. Overall sleep
6
7 efficiency was just above 80%. Results from the analysis at post-intervention indicated that as
8
9 predicted, participants went to bed 35 minutes earlier than at baseline ($\beta=-0.58, p=.003$),
10
11 there was a significant decrease of 21 minutes in how long participants took to fall asleep
12
13 (SOL; $\beta=-0.37, p=.032$), participants spent significantly less time in bed after waking than
14
15 they did at baseline ($\beta=-0.27, p<.001$), and woke significantly less frequently during the night
16
17 reducing to an average of 0.87 times (NWAK; $\beta=-0.46, p=.011$). There were also
18
19 improvements in total sleep time of 33 minutes (TST; $\beta=0.53, p=.005$), SE ($\beta=5.25, p=.016$),
20
21 how refreshing sleep was ($\beta=0.43, p<.001$) and sleep quality ($\beta=0.31, p=.018$). There were no
22
23 significant differences in the time participants woke up in the morning (on average, at
24
25 7:20am), TIB, WASO or medication use (all $ps>.05$). Within-group Cohen's d effect sizes
26
27 ranged from small to medium (0.31-0.68).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. Sleep Diary Measures

Outcome	Baseline		Post-intervention		Modelled change from baseline- to post- intervention	<i>d</i>
	<i>n</i>	<i>M (SD)</i>	<i>n</i>	<i>M (SD)</i>	β [95% CI]	
Time woken up	48	07:27 (0:46)	29	07:20 (0:54)	-0.10 [-0.41-0.21]	0.13
Time in bed after final morning wake up (h:min)	48	0:29 (0:23)	28	0:13 (0:10)	-0.27 [-0.41, -0.12]***	0.70
Time fallen asleep	48	23:29 (1:19)	28	22:54 (1:03)	-0.58 [-0.95, -0.21]**	0.44
TIB (h:min)	48	9:39 (1:11)	29	9:38 (0:55)	-0.01 [-0.42, 0.41]	0.01
SOL (h:min)	48	1:12 (1:01)	28	0:51 (0:47)	-0.37 [-0.70, -0.03]*	0.36
WASO (h:min)	47	0:14 (0:14)	29	0:12 (0:14)	-0.04 [-0.12, 0.05]	0.17
TST (h:min)	48	7:40 (1:09)	28	8:13 (1:06)	0.53 [0.17, 0.90]**	0.46
NWAK	48	1.47 (1.50)	29	0.87 (1.35)	-0.46 [-0.81, -0.11]*	0.31
SE (%)	48	80.12 (12.01)	28	85.64 (11.30)	5.25 [1.03, 9.47]*	0.44
Sleep refreshingness	48	2.37 (0.63)	29	2.78 (0.78)	0.43 [0.19, 0.68]***	0.68
Sleep quality	48	2.84 (0.74)	29	3.10 (0.82)	0.31 [0.06, 0.56]*	0.42
Use of medication	48	0.12 (0.30)	29	0.11 (0.30)	-0.01 [-0.02, 0.01]	0.03

Note. Raw means (SDs) are presented. Cohen's *d* values are time effects for pre-intervention to post-intervention using the modelled mean difference divided by the sample pre-intervention SD. Time fallen asleep and Time woken up are expressed as times in 24-hour time. Time variables (TIB, SOL, WASO, TST) are expressed in hours:minutes. TIB = Time in bed; SOL = Sleep onset latency; WASO = Wake after sleep onset; TST = Total sleep time; NWAK = Number of awakenings; SE = Sleep efficiency. SE is expressed as a percentage. Refreshingness of sleep is rated on a Likert scale from 1 = *Exhausted* to 5 = *Very refreshed*. Quality of sleep is rated on a Likert scale from 1 = *Very Poor* to 5 = *Very Good*. Use of medication is expressed as a proportion of days medication was used to help with sleep.

* $p < .05$, ** $p < .01$, *** $p < .001$.

Discussion

The purpose of this pilot study was to evaluate the feasibility, acceptability and preliminary effects of the Sleep Ninja app on sleep and mental health symptoms, for use among adolescents with sleep difficulties. A secondary objective of the study was to gather information in order to refine aspects of the app before evaluating it in a larger trial. Our findings confirmed that young people with sleep difficulties were optimistic about using the app and could complete baseline questionnaires and sleep diaries using an automated digital format without assistance. Feasibility was confirmed based on uptake, completion and retention rates with young people volunteering for the study, downloading the app and completing most of the lessons. This provides evidence that the Sleep Ninja app is a feasible intervention to deliver to young people experiencing sleep difficulties.

Intervention adherence levels suggested that while more than half of the participants completed more than half of the app, only one third of participants completed all six lessons. While this is within the range of adherence rates reported in the literature for technology-mediated insomnia programs [44], reasons for non-adherence require consideration as there is room for improvement. The main reason participants had difficulty or stopped using the app was reportedly due to the amount of text presented in the app and the repetitive nature of the material. Participants also requested a more tailored experience. Those who completed more of the app also reported it to be more acceptable. Therefore, it is likely that refining the app by taking these points into account is likely to increase engagement with the content and overall adherence to the intervention.

Efficacy outcomes showed that insomnia symptoms improved significantly from baseline to post-study, effectively moving participants from the lower cut off for clinical insomnia, firmly into the sub-threshold symptom level. There was an improvement in self-reported sleep quality, with a medium effect size ($d=-0.46$) suggesting app use improves

1
2
3 quality of sleep. The improvements detected on the two sleep questionnaire measures (ISI,
4 PQSI) were corroborated by those found in the sleep diaries, with participants going to bed
5 earlier, falling asleep more quickly at night, waking less frequently, sleeping for longer,
6
7
8
9
10 spending less time in bed in the morning after waking, and reporting improved sleep
11
12
13 efficiency and quality.

14
15 The time that participants reported waking each morning did not shift. This finding
16
17 needs to be considered in the context of young people having to get up at a specific time for
18
19 school each morning. School start times are a modifiable contributing factor to insufficient
20
21 sleep, with evidence suggesting that delaying school start times by even 30 minutes is
22
23 associated with higher levels of school attendance, lower daytime sleepiness, and improved
24
25 attention and concentration [45]. While there have been calls to delay school start times to
26
27 improve sleep duration, health and functioning (e.g., [46]), this is unlikely to change in the
28
29 Australian context. Accordingly, a focus on bed times rather than rise times is likely to be
30
31 most useful for treating sleep difficulties in this age group, at least for the time being.
32
33
34

35
36 Findings on these sleep outcomes are consistent with the two studies which have
37
38 evaluated web-delivered CBT-I for adolescents with insomnia [15, 16, 17], and show for the
39
40 first time that CBT-I, when delivered to adolescents by smartphone-app, confer benefits. It is
41
42 encouraging that the within-group effect size obtained in the current study for insomnia ($d=$
43
44 0.90) is comparable to within-group effect size found for digitally-delivered CBT-I in a
45
46 randomised trial ($d=-0.92$; [16]). In this randomised study, the intervention group was found
47
48 to be superior to the waitlist control, suggesting that a within-group effect size of this
49
50 magnitude is likely to reflect improvement over and above what would be expected based on
51
52 a standard placebo effect [16].
53
54

55
56 Beyond the sleep outcomes, we also found that there were decreases in depression and
57
58 anxiety symptoms following the completion of the intervention suggesting that there may be
59
60

1
2
3 value in using this app to address mental illness. The magnitude of the decrease in depression
4 scores is notable, with individuals moving from the moderate range into the mild range, and a
5 within-group effect size that is comparable to other adolescent depression prevention trials
6 [47]. These findings provide proof-of-principle evidence that the Sleep Ninja app may be
7 useful in addressing depression. As a pilot study, we did not test the specific hypothesis that
8 targeting insomnia will decrease depression risk. Given our encouraging findings, a follow-
9 up randomised controlled trial which follows participants over time, which can determine
10 causality between insomnia and depression, that can assess the mechanisms of change, as
11 well as the impact of intervention on depression risk is now warranted.
12
13
14
15
16
17
18
19
20
21
22
23

24 This study makes a unique contribution to the literature by showing that smartphone
25 delivery of CBT-I is a promising format in which to deliver this gold-standard intervention.
26 This is the first study that we are aware of that has evaluated app-delivered CBT-I in young
27 people [21], and only the second study that has tested mobile phone delivery of CBT-I, the
28 first study being conducted in adults, with positive effects on sleep outcomes [48]. Using
29 smartphones to deliver interventions such as this offer a myriad of advantages, including
30 immediate connectivity to automated interactive applications that can be accessed anytime,
31 anywhere. Sleep Ninja has been developed so that it does not rely on internet access for use,
32 which is likely to be important to young people who may have limited data plans, and
33 individuals who do not have optimal internet coverage. Not requiring internet coverage to
34 access digital programs represents a new wave of flexibility in the delivery of health
35 interventions and the automated nature of the intervention means it can be delivered without
36 professional support. It is notable that we had a 72% retention rate, which is at the upper level
37 of that detected for digital interventions which has been shown in a meta-analysis to range
38 between 43-99% [49], with retention rates typically lower in non-supported interventions
39 [50].
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

There are several study limitations that need to be considered. First, participants in this trial were required to have relatively minor levels of insomnia symptoms for study entry. This decision was made by balancing inclusiveness against the requirement of sleep disturbance to ensure participants were motivated to use the app. Moreover, given the study focus was on feasibility and acceptability, we felt it would be prudent to establish these factors before targeting a more severe participant group. That said, while we set the threshold for entry relatively low, both the mean and the median converged on an ISI score of just above 14, indicating that participants were at the junction between having subthreshold and clinical levels of insomnia symptoms (cut-off score is 15). Therefore, it is relatively unlikely that there would have been a floor effect as the data showed there was sufficient room to detect symptom-improvement. The high mean symptom level also suggests that the results may generalise to a group with clinical levels of insomnia. Second, we did not include a control group. Again, as the study goal was to establish feasibility and acceptability, it was not necessary to include a control group for this purpose. However, this design is not able to attribute causality to the intervention and a controlled study is now needed. Finally, this study relied exclusively on subjectively reported sleep outcomes. Although objective measures such as polysomnography provide the most accurate way to assess sleep, there is evidence that subjectively measured sleep variables could be more closely associated with functional outcomes [51]. Moreover, subjectively experienced sleep quality and parameters have consistently shown to be strongly associated with psychological wellbeing [52, 53], suggesting that perception of sleep is as important, if not more so, than objective measures. We are currently investigating how inbuilt smartphone sensors such as the accelerometer might be used to provide a more objective estimate of sleep.

56
57
58
59
60

This study provides preliminary evidence supporting the feasibility, acceptability and effects of a fully-automated app that targets adolescent sleep difficulties. The Sleep Ninja

1
2
3 intervention shows promise both as a sleep-focused intervention, but also potentially to
4
5 reduce risk for depression.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. Example screens from the Sleep Ninja app. From left: Homescreen, Training Session Access and Progress Record, Tracking and Bedtime Setting, More Information

Figure 2. Participant Flow

For peer review only

Acknowledgements

Sincere thanks to Tobias Reid, Jacinto Santamaria and Cesar Anonuevo for their tireless IT support on this project. Thanks also to Dean Winder for assistance in recruitment. The authors would also like to thank the participants and their families for their enthusiasm in this project. AW-S is supported by a NSW Health Early-to-Mid Career Fellowship, MT is supported by an early-career NHMRC Fellowship #1138710 and HC is supported by NHMRC Fellowship #1056964.

Author Statement

AW-S, BO, MT and HC conceived of the study and the trial design. AW-S designed the study with input from all authors, and oversaw the management of the trial. LJ led trial recruitment, managed the day-to-day running of the trial and conducted the participant interviews. QW conducted the analyses with assistance from AW-S and LJ. All authors contributed to the preparation of the manuscript.

Competing Interests Statement

No competing interests to declare.

Funding Statement

This work was supported by a grant awarded to Aliza Werner-Seidler and the Black Dog Institute by the Corella Foundation. Funders had no role in the design, implementation or reporting of this study.

Data Sharing Statement

No additional unpublished data from this study is publicly available.

References

1. American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed. ed.). Washington DC: Author.
2. Ohayon, M. M., Roberts, R. E., Zulley, J., Smirne, S., & Priest, R. G. (2000). Prevalence and patterns of problematic sleep among older adolescents. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39(12), 1549-1556. doi:10.1097/00004583-200012000-00019
3. Carskadon, M. A. (1990). Patterns of sleep and sleepiness in adolescents. *Pediatrician*, 17(1), 5-12.
4. Liu, X., Buysse, D. J., Gentzler, A. L., Kiss, E., Mayer, L., Kapornai, K., ... Kovacs, M. (2007). Insomnia and hypersomnia associated with depressive phenomenology and comorbidity in childhood depression. *Sleep*, 30(1), 83-90.
5. Baglioni, C., Battagliese, G., Feige, B., Spiegelhalder, K., Nissen, C., Voderholzer, U., ... Riemann, D. (2011). Insomnia as a predictor of depression: A meta-analytic evaluation of longitudinal epidemiological studies. *Journal of Affective Disorders*, 135(1-3), 10-19. doi:10.1016/j.jad.2011.01.011
6. Franzen, P. L., & Buysse, D. J. (2008). Sleep disturbances and depression: Risk relationships for subsequent depression and therapeutic implications. *Dialogues in Clinical Neuroscience*, 10(4), 473-481.
7. Riemann, D., & Voderholzer, U. (2003). Primary insomnia: A risk factor to develop depression? *Journal of Affective Disorders*, 76(1-3), 255-259.
8. Ashworth, D. K., Sletten, T. L., Junge, M., Simpson, K., Clarke, D., Cunnington, D., & Rajaratnam, S. M. (2015). A randomized controlled trial of cognitive behavioral therapy for insomnia: An effective treatment for comorbid insomnia and depression. *Journal of Counselling Psychology*, 62(2), 115-123. doi:10.1037/cou0000059

- 1
2
3 9. Manber, R., Edinger, J. D., Gress, J. L., San Pedro-Salcedo, M. G., Kuo, T. F., &
4
5 Kalista, T. (2008). Cognitive behavioral therapy for insomnia enhances depression
6
7 outcome in patients with comorbid major depressive disorder and insomnia. *Sleep*,
8
9 *31*(4), 489-495.
10
11
- 12 10. Wagley, J. N., Rybarczyk, B., Nay, W. T., Danish, S., & Lund, H. G. (2013).
13
14 Effectiveness of abbreviated CBT for insomnia in psychiatric outpatients: Sleep and
15
16 depression outcomes. *Journal of Clinical Psychology*, *69*(10), 1043-1055.
17
18 doi:10.1002/jclp.21927
19
- 20 21 11. Batterham, P. J., Christensen, H., Mackinnon, A. J., Gosling, J. A., Thorndike, F. P.,
22
23 Ritterband, L. M., ... Griffiths, K. M. (2017). Trajectories of change and long-term
24
25 outcomes in a randomised controlled trial of internet-based insomnia treatment to
26
27 prevent depression. *BJPsych Open*, *3*(5), 228-235. doi:10.1192/bjpo.bp.117.005231
28
29
- 30 31 12. Christensen, H., Batterham, P. J., Gosling, J. A., Ritterband, L. M., Griffiths, K. M.,
32
33 Thorndike, F. P., ... Mackinnon, A. J. (2016). Effectiveness of an online insomnia
34
35 program (SHUTi) for prevention of depressive episodes (the GoodNight Study): A
36
37 randomised controlled trial. *Lancet Psychiatry*, *3*(4), 333-341. doi:10.1016/s2215-
38
39 0366(15)00536-2
40
41
- 42 43 13. Blake, M., Waloszek, J. M., Schwartz, O., Raniti, M., Simmons, J. G., Blake, L., ...
44
45 Allen, N. B. (2016). The SENSE study: Post intervention effects of a randomized
46
47 controlled trial of a cognitive-behavioral and mindfulness-based group sleep
48
49 improvement intervention among at-risk adolescents. *Journal of Consulting and*
50
51 *Clinical Psychology*, *84*(12), 1039-1051. doi:10.1037/ccp0000142
52
53
- 54 55 14. Waloszek, J. M., Schwartz, O., Simmons, J. G., Blake, M., Blake, L., Murray, G., ...
56
57 Allen, N. B. (2015). The SENSE Study (Sleep and Education: learning New Skills
58
59 Early): A community cognitive-behavioural therapy and mindfulness-based sleep
60

- 1
2
3 intervention to prevent depression and improve cardiac health in adolescence. *BMC*
4
5 *Psychology*, 3, 39. doi:10.1186/s40359-015-0096-x
6
7
- 8 15. de Bruin, E. J., Bogels, S. M., Oort, F. J., & Meijer, A. M. (2018). Improvement of
9
10 adolescent psychopathology after insomnia treatment: Results from a randomized
11
12 controlled trial over 1 year. *The Journal of Child Psychology and Psychiatry*, 59(5),
13
14 509-522. doi:10.1111/jcpp.12834.
15
16
- 17 16. de Bruin, E. J., Bogels, S. M., Oort, F. J., & Meijer, A. M. (2015). Efficacy of
18
19 cognitive behavioral therapy for insomnia in adolescents: A randomized controlled
20
21 trial with internet therapy, group therapy and a waiting list condition. *Sleep*, 38(12),
22
23 1913-1926. doi:10.5665/sleep.5240
24
25
- 26 17. de Bruin, E. J., Oort, F. J., Bogels, S. M., & Meijer, A. M. (2014). Efficacy of internet
27
28 and group-administered cognitive behavioral therapy for insomnia in adolescents: A
29
30 pilot study. *Behavioral Sleep Medicine*, 12(3), 235-254.
31
32 doi:10.1080/15402002.2013.784703
33
34
- 35 18. Okajima, I., Komada, Y., & Inoue, Y. (2011). A meta-analysis on the treatment
36
37 effectiveness of cognitive behavioral therapy for primary insomnia. *Sleep and*
38
39 *Biological Rhythms*, 9(1), 24-34. doi:10.1111/j.1479-8425.2010.00481.x
40
41
- 42 19. Zachariae, R., Lyby, M. S., Ritterband, L. M., & O'Toole, M. S. (2016). Efficacy of
43
44 internet-delivered cognitive-behavioral therapy for insomnia - A systematic review
45
46 and meta-analysis of randomized controlled trials. *Sleep Medicine Reviews*, 30, 1-10.
47
48 doi:10.1016/j.smrv.2015.10.004
49
50
- 51 20. Gulliver, A., Griffiths, K. M., & Christensen, H. (2010). Perceived barriers and
52
53 facilitators to mental health help-seeking in young people: A systematic review. *BMC*
54
55 *Psychiatry*, 10, 113-113. doi:10.1186/1471-244X-10-113
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
21. Werner-Seidler, A., Johnston, L., & Christensen, H. (2018). Digitally-delivered cognitive-behavioural therapy for youth insomnia: A systematic review. *Internet Interventions, 11*, 71-78. doi:10.1016/j.invent.2018.01.007
22. Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2008). Developing and evaluating complex interventions: The new Medical Research Council guidance. *BMJ, 337*, a1655. doi:10.1136/bmj.a1655
23. Gitlin, L. N. (2013). Introducing a new intervention: An overview of research phases and common challenges. *American Journal of Occupational Therapy, 67*(2), 177-184. doi:10.5014/ajot.2013.006742
24. Bastien, C. H., Vallieres, A., & Morin, C. M. (2001). Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine, 2*(4), 297-307.
25. Chung, K. F., Kan, K. K.-K., & Yeung, W.-F. (2011). Assessing insomnia in adolescents: Comparison of Insomnia Severity Index, Athens Insomnia Scale and Sleep Quality Index. *Sleep Medicine, 12*(5), 463-470.
26. Chahoud, M., Chahine, R., Salameh, P., & Sauleau, E. A. (2017). Reliability, factor analysis and internal consistency calculation of the Insomnia Severity Index (ISI) in French and in English among Lebanese adolescents. *eNeurologicalSci, 7*, 9-14. doi:10.1016/j.ensci.2017.03.003
27. Alvaro, P.K., Roberts, R.M., & Harris, J.K. (2014). The independent relationships between insomnia, depression, subtypes of anxiety and chronotype during adolescence. *Sleep Medicine, 15*, 934-941. doi: 10.1016/j.sleep.2014.03.019
28. de la Vega, R., Tome-Pires, C., Sole, E., Racine, M., Castarlenas, E., Jensen, M. P., & Miro, J. (2015). The Pittsburgh Sleep Quality Index: Validity and factor structure in young people. *Psychological Assessment, 27*(4), e22-27. doi:10.1037/pas0000128

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
29. Cole, J. C., Motivala, S. J., Buysse, D. J., Oxman, M. N., Levin, M. J., & Irwin, M. R. (2006). Validation of a 3-factor scoring model for the Pittsburgh Sleep Quality Index in older adults. *Sleep*, *29*(1), 112.
30. Guo, S., Sun, W., Liu, C., & Wu, S. (2016). Structural validity of the Pittsburgh Sleep Quality Index in Chinese undergraduate students. *Frontiers in Psychology*, *7*, 1126. doi:10.3389/fpsyg.2016.01126
31. Johnson, J.G., Harris, E.S., Spitzer, R.L. & Williams, J.B.W. (2002). The Patient Health Questionnaire for Adolescents: Validation of an instrument for the assessment of mental disorders among adolescent primary care patients. *Journal of Adolescent Health*, *30*, 196–204.
32. Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, *16*(9), 606-613. doi:10.1046/j.1525-1497.2001.016009606.x
33. Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B. W., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, *114*(1), 163-173. doi:10.1016/j.jad.2008.06.026
34. Spitzer, R. L., Kroenke, K., Williams, J. B., & Lowe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7. *Archives of Internal Medicine*, *166*(10), 1092-1097. doi:10.1001/archinte.166.10.1092
35. Tiirikainen, K., Haravouri, H., Ranta, K., Kaltiala-Heino, R. & Marttunen, M. (2019). Psychometric properties of the 7-item Generalized Anxiety Disorder Scale (GAD-7) in a large representative sample of Finnish adolescents. *Psychiatry Research*, *272*, 30-35.

- 1
2
3 36. Thorndike, F. P., Saylor, D. K., Bailey, E. T., Gonder-Frederick, L., Morin, C. M., &
4
5 Ritterband, L. M. (2008). Development and perceived utility and impact of an internet
6
7 intervention for insomnia. *E-journal of Applied Psychology : Clinical and Social*
8
9 *Issues, 4(2)*, 32-42.
- 10
11
12 37. Ritterband, L. M., Ardan, K., Thorndike, F. P., Magee, J. C., Cox, D. J., Sutphen, J.
13
14 L., & Borowitz, S. M. (2008). Real world use of an Internet intervention for pediatric
15
16 encopresis. *Journal of Medical Internet Research, 10(2)*, e16. doi:10.2196/jmir.1081.
- 17
18
19 38. Carney, C. E., Buysse, D. J., Ancoli-Israel, S., Edinger, J. D., Krystal, A. D., Lichstein,
20
21 K. L., & Morin, C. M. (2012). The Consensus Sleep Diary: Standardizing prospective
22
23 sleep self-monitoring. *Sleep, 35(2)*, 287-302. doi:10.5665/sleep.1642
- 24
25
26 39. Werner-Seidler, A., O'Dea, B., Shand, F., Johnston, L., Frayne, A., Fogarty, A. S., &
27
28 Christensen, H. (2017). A smartphone app for adolescents with sleep disturbance:
29
30 Development of the Sleep Ninja. *JMIR Mental Health, 4(3)*, e28.
31
32 doi:10.2196/mental.7614
- 33
34
35 40. Harvey, A. G. (2009). A transdiagnostic approach to treating sleep disturbance in
36
37 psychiatric disorders. *Cognitive Behaviour Therapy, 38 Suppl 1*, 35-42.
38
39 doi:10.1080/16506070903033825
- 40
41
42 41. Cain, N. & Gradisar, M. (2010). Electronic media use and sleep in school-aged
43
44 children and adolescents: A review. *Sleep Medicine, 11*, 735-742
45
46 doi:10.1016/j.sleep.2010.02.006
- 47
48
49 42. Hysing, M., Pallesen, S., Stormark, K.M., Jakobsen, R., Lundervold, A.J., & Sivertsen,
50
51 B. (2014). Sleep and use of electronic devices in adolescence: Results from a large
52
53 population-based study. *BMJ Open, 5*, e006748. doi:10.1136/bmjopen-2014-006748
- 54
55
56 43. Kline, R. B. (2011). *Methodology in the Social Sciences. Principles and practice of*
57
58 *structural equation modeling (3rd ed.)*. New York, NY, US: Guilford Press.
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
44. Horsch, C. H., Lancee, J., Jan Beun, R., Neerincx, M. A. & Brinkman, W. P. (2015). Adherence to technology-mediated insomnia treatment: a meta-analysis, interviews and focus groups. *Journal of Medical Internet Research*, 17(9), e214. doi: 10.2196/jmir.4115
45. Wheaton, A. G., Chapman, D. P. & Croft. (2016). School start times, sleep, behavioral, health, and academic outcomes: a review of the literature. *Journal of School Health*, 86(5), 363-381. doi: 10.1111/josh.12388
46. Watson, N. F., Martine, J. L., Wise, K. A., Kirsch, D. B. Kristo, D.B.... & Chervin, R. D. (2017). Delaying middle school and high school start times promotes student health and performance. An American Academy of Sleep Medicine Position Statement, *Journal of Clinical Sleep Medicine*, 13(4), 623-623. doi: 10.5664/jcsm.6558
47. Horowitz, J. L., Garber, J., Ciesla, J. A., Young, J. F., & Mufson, L. (2007). Prevention of depressive symptoms in adolescents: A randomized trial of cognitive-behavioral and interpersonal prevention programs. *Journal of Consulting and Clinical Psychology*, 75(5), 693-706. doi:10.1037/0022-006x.75.5.693
48. Horsch, C. H., Lancee, J., Griffioen-Both, F., Spruit, S., Fitrianie, S., Neerincx, M. A., ... Brinkman, W. P. (2017). Mobile phone-delivered cognitive behavioral therapy for insomnia: A randomized waitlist controlled trial. *Journal of Medical Internet Research*, 19(4), e70. doi:10.2196/jmir.6524
49. Christensen, H., Griffiths, K. M., & Farrer, L. (2009). Adherence in internet interventions for anxiety and depression: Systematic review. *Journal of Medical Internet Research*, 11(2), e13.

- 1
2
3 50. Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized
4
5 psychological treatments for adult depression: A meta-analysis. *Cognitive Behaviour*
6
7 *Therapy*, 38(4), 196-205. doi:10.1080/16506070903318960
8
9
10 51. Wilson, D. L., Fung, A., Walker, S. P., & Barnes, M. (2013). Subjective reports versus
11
12 objective measurement of sleep latency and sleep duration in pregnancy. *Behavioral*
13
14 *Sleep Medicine*, 11(3), 207-221. doi:10.1080/15402002.2012.670674
15
16
17 52. Fuligni, A. J., & Hardway, C. (2006). Daily variation in adolescents' sleep, activities,
18
19 and psychological well-being. *Journal of Research on Adolescence*, 16(3), 353-378.
20
21 doi:10.1111/j.1532-7795.2006.00498.x
22
23
24 53. Lund, H. G., Reider, B. D., Whiting, A. B., & Prichard, J. R. (2010). Sleep patterns
25
26 and predictors of disturbed sleep in a large population of college students. *Journal of*
27
28 *Adolescent Health*, 46(2), 124-132. doi:10.1016/j.jadohealth.2009.06.016
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Example screens from the Sleep Ninja app. From left: Homescreen, Training Session Access and Progress Record, Tracking and Bedtime Setting, More Information

139x238mm (300 x 300 DPI)




Figure 2. Participant Flow

187x238mm (300 x 300 DPI)



CONSORT 2010 checklist of information to include when reporting a pilot or feasibility trial*

Section/Topic	Item No	Checklist item	Reported on page No
Title and abstract			
	1a	Identification as a pilot or feasibility randomised trial in the title	1
	1b	Structured summary of pilot trial design, methods, results, and conclusions (for specific guidance see CONSORT abstract extension for pilot trials)	2
Introduction			
Background and objectives	2a	Scientific background and explanation of rationale for future definitive trial, and reasons for randomised pilot trial	5-6
	2b	Specific objectives or research questions for pilot trial	6
Methods			
Trial design	3a	Description of pilot trial design (such as parallel, factorial) including allocation ratio	6
	3b	Important changes to methods after pilot trial commencement (such as eligibility criteria), with reasons	N/A
Participants	4a	Eligibility criteria for participants	7
	4b	Settings and locations where the data were collected	13
	4c	How participants were identified and consented	13
Interventions	5	The interventions for each group with sufficient details to allow replication, including how and when they were actually administered	11-12
Outcomes	6a	Completely defined prespecified assessments or measurements to address each pilot trial objective specified in 2b, including how and when they were assessed	7-11, 13
	6b	Any changes to pilot trial assessments or measurements after the pilot trial commenced, with reasons	N/A
	6c	If applicable, prespecified criteria used to judge whether, or how, to proceed with future definitive trial	N/A
Sample size	7a	Rationale for numbers in the pilot trial	7
	7b	When applicable, explanation of any interim analyses and stopping guidelines	N/A
Randomisation:			
Sequence generation	8a	Method used to generate the random allocation sequence	N/A
	8b	Type of randomisation(s); details of any restriction (such as blocking and block size)	N/A
Allocation concealment mechanism	9	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned	N/A

Implementation	10	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions	N/A
Blinding	11a	If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	N/A
	11b	If relevant, description of the similarity of interventions	N/A
Statistical methods	12	Methods used to address each pilot trial objective whether qualitative or quantitative	14-15
Results			
Participant flow (a diagram is strongly recommended)	13a	For each group, the numbers of participants who were approached and/or assessed for eligibility, randomly assigned, received intended treatment, and were assessed for each objective	Figure 2
	13b	For each group, losses and exclusions after randomisation, together with reasons	16
Recruitment	14a	Dates defining the periods of recruitment and follow-up	7
	14b	Why the pilot trial ended or was stopped	N/A
Baseline data	15	A table showing baseline demographic and clinical characteristics for each group	Table 1, p. 15
Numbers analysed	16	For each objective, number of participants (denominator) included in each analysis. If relevant, these numbers should be by randomised group	Table 2, p. 19 Table 3, p. 21
Outcomes and estimation	17	For each objective, results including expressions of uncertainty (such as 95% confidence interval) for any estimates. If relevant, these results should be by randomised group	Table 2, p. 19 Table 3, p. 21
Ancillary analyses	18	Results of any other analyses performed that could be used to inform the future definitive trial	16-19
Harms	19	All important harms or unintended effects in each group (for specific guidance see CONSORT for harms)	N/A
	19a	If relevant, other important unintended consequences	N/A
Discussion			
Limitations	20	Pilot trial limitations, addressing sources of potential bias and remaining uncertainty about feasibility	25
Generalisability	21	Generalisability (applicability) of pilot trial methods and findings to future definitive trial and other studies	23-24
Interpretation	22	Interpretation consistent with pilot trial objectives and findings, balancing potential benefits and harms, and considering other relevant evidence	22-23
	22a	Implications for progression from pilot to future definitive trial, including any proposed amendments	24
Other information			
Registration	23	Registration number for pilot trial and name of trial registry	7
Protocol	24	Where the pilot trial protocol can be accessed, if available	N/A
Funding	25	Sources of funding and other support (such as supply of drugs), role of funders	28
	26	Ethical approval or approval by research review committee, confirmed with reference number	13

1 Citation: Eldridge SM, Chan CL, Campbell MJ, Bond CM, Hopewell S, Thabane L, et al. CONSORT 2010 statement: extension to randomised pilot and feasibility trials. BMJ. 2016;355.
 2 *We strongly recommend reading this statement in conjunction with the CONSORT 2010, extension to randomised pilot and feasibility trials, Explanation and Elaboration for important
 3 clarifications on all the items. If relevant, we also recommend reading CONSORT extensions for cluster randomised trials, non-inferiority and equivalence trials, non-pharmacological
 4 treatments, herbal interventions, and pragmatic trials. Additional extensions are forthcoming: for those and for up to date references relevant to this checklist, see www.consort-statement.org.
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42

For peer review only

6/bmjopen-2016-026502 on 27 May 2019. Downloaded from http://bmjopen.bmj.com/ on April 19, 2024 by guest. Protected by copyright.