BMJ Open Cross-sectional study about the use of telemedicine for type 2 diabetes mellitus management in Spain: patient's perspective. The EnREDa2 Study

Patricia Rodríguez-Fortúnez,¹ Josep Franch-Nadal,^{2,3} José A Fornos-Pérez,⁴ Fernando Martínez-Martínez,⁵ Hector David de Paz,⁶ María Luisa Orera-Peña¹

To cite: Rodríguez-Fortúnez P, Franch-Nadal J, Fornos-Pérez JA, *et al.* Cross-sectional study about the use of telemedicine for type 2 diabetes mellitus management in Spain: patient's perspective. The EnREDa2 Study. *BMJ Open* 2019;**9**:e028467. doi:10.1136/bmjopen-2018-028467

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-028467>).

Received 14 December 2018

Revised 3 June 2019

Accepted 4 June 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Patricia Rodríguez-Fortúnez; patricia.rodriguez@scren.es

ABSTRACT

Objectives The usefulness of telemedicine (TM) in type 2 diabetes mellitus (T2DM) has been discussed in recent years. The aim of this study is to describe patients' perceptions about TM and to identify preferences on TM resources, in Spain.

Design An observational, cross-sectional study was conducted using a structured questionnaire.

Participants 1036 patients with T2DM accepted to participate in the study (response rate: 68%).

Results Blood glucose values were recorded by 85.9% of the patients while data such as lifestyle habits were only recorded by 14.4% of the patients. Previous experience in TM was reported by 9.8% of the patients, out of which 70.5% were satisfied with its service and 73.5% considered that the use of TM had optimised their T2DM management. However, most of these patients noted aspects to be improved such as user-friendliness (81.4%), interaction with the medical team (78.4%) and time required for recording/transferring data (78.4%). Experienced patients had better perception about TM usefulness than naïve patients for all listed aspects ($p < 0.05$). Among naïve patients, 38.2% expressed their willingness to participate in TM programmes, but only 4.7% were invited to do so. Patients considered that physicians' (77.5%) and pharmacists' (75.5%) encouragement can boost the use of TM.

Conclusions In Spain, nearly 10% of patients with T2DM have experience with TM and it is well accepted, especially one based on glucometers. Nevertheless, in order to promote TM use, easier and time-saving programmes for patient-physician interaction should be optimised.

INTRODUCTION

In Spain, the prevalence of type two diabetes mellitus (T2DM) is 13.8% (2010).¹ T2DM is one of the most common global metabolic disorders and its prevalence is expected to increase in the coming years.² The rise in patients with T2DM will contribute to an increase in the need for healthcare.³ New information and communication technologies (ICTs) can allow the exchange of data between physicians and patients to be

Strengths and limitations of this study

- This is one of the few recent studies that evaluate the prevalence and patient's preferences about the use of telemedicine TM in Spain with an important sample size.
- This study presents a number of limitations inherent in its observational design, including susceptibility to bias and confounding, which restrict the ability to define causality.
- In addition, the results must be interpreted while keeping in mind that all of the issues evaluated, including those related to the patient's clinical situation, are always expressed from the patient's point of view, which increases the possibility that the issues raised here may have different interpretations.
- Furthermore, snowball technique could lead to potential sampling bias and low sample representativeness.
- Nevertheless, the wide sample variability observed (clinical and sociodemographic) suggests a very limited bias.

faster and more fluid, and increase patients' involvement in the self-management of their disease.⁴ Similarly, they can make exchange information easier for medical specialists, avoiding the need for referral and reducing waiting lists.^{5–7} Thus, the use of health telematics to manage these conditions could improve the quality of healthcare for these patients. A number of studies support the use of telemedicine (TM) for chronic disease management, demonstrating how its use improves compliance with treatment recommendations, and reduces the time spent per visit and the generated costs.^{8–12}

T2DM is one of the chronic diseases which can benefit the most from TM. The use of TM contributes to optimisation of glycaemic control. Thus, patients who use TM have shown a greater reduction of glycated haemoglobin (HbA1c) levels than those

receiving usual care.^{13 14} Moreover, the use of TM, at least in specific interventions, helps to optimise improvement of weight control,¹⁵ body mass index,¹⁵ blood pressure,^{16 17} low-density lipoprotein values,^{17 18} pain¹⁹ and to improve health-related quality of life.^{15 19 20} At the same time, TM is a useful tool for reducing the number of visits to the doctor by patients with T2DM, thereby saving time and money on commutes.^{21–23}

Although the use of TM has shown to have multiple benefits and policy makers and payers consider it to be of interest,²⁴ there are factors related to patients' trust²⁵ and sociodemographic characteristics²⁶ that can make its implementation for T2DM management difficult. The aim of this study is to describe and clarify the current situation in Spain regarding the use of TM by patients with T2DM, patients' perceptions about the benefits of using TM and their preferences for TM resources.

SUBJECTS AND METHODS

Participants

Patients ≥18 years of age with T2DM of at least 1 year's duration were included. Patients from every Spanish region were selected using snowball sampling techniques.²⁷ The initial participants were recruited from primary care of the Spanish Healthcare System from a diverse set of primary care offices. The physicians involved in the study had at least 2 years of clinical experience in T2DM management.

The population size was estimated assuming maximum indetermination, with a 99% CI and a 4% precision error margin. The adult Spanish population for 2015 (37 007 319)²⁸ and T2DM prevalence in Spain (13.8%)¹ were considered. This resulted in a final planned sample size of 1036 patients.

Study design

A descriptive observational study was conducted from 18 April 2016 to 5 August 2016. An ad hoc questionnaire was designed after literature review about the use of TM for T2DM management and about patients' preferences for TM resources in Spain. A scientific committee of two expert physicians was involved in the questionnaire composition, which consisted of 48 questions divided into four sections: (1) sociodemographic variables, (2) patient-reported clinical variables, (3) matters related to the use of ICTs and TM resources and (4) patients' perceptions about the use of TM. This latter section distinguished between patients who had reported prior experience using TM and those who had none (naïve). TM was previously defined as the exchange of medical information to healthcare professionals via electronic communication. Open, closed-ended (dichotomous or multichoice) and 5-point Likert-scale questions comprised the survey (see online supplementary file). The survey was conducted online and by phone.

Statistical analysis

Statistical analysis was performed using STATA statistical software, V.14. In all statistical tests, p value <0.05

was considered to be statistically significant. No methods for handling with missing data were applied due to the high sample size and the small amount of missing data. Absolute and relative frequencies were calculated for qualitative variables. Means, SD and percentile distributions were used to describe quantitative variables. The χ^2 test was used for comparing qualitative variables between groups while quantitative variables were compared using Student's t-test or the equivalent method for non-parametric variables (Mann-Whitney U test).

In addition, since the technology skills of the elderly tend to be less developed,²⁹ a subanalysis of questions related to the use of ICTs based on patient's age (<65 vs ≥65 years) was performed using the χ^2 test.

For clarity, responses to Likert-scale questions have been grouped in three categories (clustering the extreme categories). However, the statistical analysis has been calculated with the disaggregated data (5-point Likert scale).

Ethical considerations

All the patients received information relevant to the study and gave their informed consent to participate. All documents were coded in order to ensure the confidentiality of the data. The results of this study will be published by an open access scientific journal and by a doctoral thesis.

Patient and public involvement

No patients were included in the design and planning of the study. Including patient and public involvement (PPI) statements aligns closely with *BMJ Open*'s values of transparency and inclusiveness. We hope that including PPI statements in all articles is the first step of many for *BMJ Open* in encouraging patient's involvement.

RESULTS

Description of the patients

A total number of 1523 patients with T2DM were invited to participate in the study. Of them, 1036 participants, representative of all the autonomous communities, accepted to participate (rate of response: 68%). When the minimum estimated sample was reached, recruitment was concluded. The mean age of the patients was 60.3 years, 50.5% were men and the majority resided in urban areas (table 1). The overwhelming majority of patients (95.6%) were receiving some type of treatment for their T2DM, the most common being a combination of oral antidiabetic drugs (64.2%) (table 2).

Use of ICTs by patients with T2DM

The vast majority of patients reported having access to ICTs in their homes: mobile or landline phone (97.5%), internet access (71.6%) and personal computers (PCs) (67.4%).

Searching for information

Only a 14.3% of the patients searched for information about their T2DM on the internet frequently or very

Table 1 Sociodemographic characteristics of the patients with T2DM

	Total (n=1036)	E (n=102)	N (n=934)	P value (E vs N)	NI (n=577)	I (n=357)	P value (ND vs N)
Age (median (SD)), years	60.3 (15.0)	42.7 (13.5)	62.3 (13.9)	<0.001	66.6 (11.9)	55.2 (13.9)	<0.001
Men (% (n))	50.5 (523)	58.8 (60)	49.6 (463)	χ^2 (df=1)=3.14, p=0.076	44.5 (257)	57.7 (206)	χ^2 (df=1)=15.28; p<0.001
Location (% (n))							
Rural	23.4 (242)	15.7 (16)	24.2 (226)	χ^2 (df=1)=4.38; p=112	26.5 (153)	20.4 (73)	χ^2 (df=1)=5.21; p=0.074
Urban	76.6 (794)	84.3 (86)	75.8 (708)		73.5 (424)	79.6 (254)	
Marital status (n (%))							
Single	6.2 (64)	5.9 (6)	6.2 (58)	χ^2 (df=3)=16.81; p=0.001	4.5 (26)	9.0 (32)	χ^2 (df=3)=26.47; p<0.001
Partnership/Married	75.6 (783)	89.2 (91)	74.1 (692)		71.6 (413)	78.2 (279)	
Separated/Divorced	6.1 (63)	4.9 (5)	6.2 (58)		6.4 (37)	5.9 (21)	
Widowed	12.2 (126)	0.0 (0)	13.5 (126)		17.5 (101)	7.0 (25)	
Level of education (% (n))							
No education	10.2 (106)	0.9 (1)	11.2 (105)	χ^2 (df=3)=50.03; p<0.001	16.8 (97)	2.2 (8)	χ^2 (df=3)=111.91; p<0.001
Primary school	20.2 (209)	3.9 (4)	21.9 (205)		26.5 (153)	14.6 (52)	
High school	39.8 (412)	39.2 (40)	39.2 (372)		39.7 (229)	40.1 (143)	
University	29.8 (309)	55.9 (57)	26.9 (252)		17.0 (98)	43.1 (154)	
Employment status (% (n))							
Student	0.3 (3)	0.9 (1)	0.2 (2)	χ^2 (df=5)=50.03; p<0.001	0.2 (1)	0.3 (1)	χ^2 (df=5)=165.38; p<0.001
Currently employed	35.8 (371)	81.4 (83)	30.8 (288)		16.8 (97)	53.5 (191)	
Disabled	1.8 (19)	2.9 (3)	1.7 (16)		1.2 (7)	2.5 (9)	
Unemployed	3.9 (41)	2.9 (2)	4.1 (38)		2.8 (16)	6.2 (22)	
Retired	39.0 (404)	7.8 (8)	42.4 (396)		52.5 (303)	26.1 (93)	
Housework	19.1 (198)	3.9 (4)	20.8 (194)		26.5 (153)	11.5 (41)	

E, experienced; I, interested in TM; N, naïve; NI, not interested in TM; T2DM, type 2 diabetes mellitus; TM, telemedicine.

frequently, 32.3% did so once in a while and 53.4% never looked for information on internet. The most common searches were for T2DM-related complications (51.6%), adverse reactions to treatment (31.1%), characteristics of new treatments (27.5%), appearance of hypoglycaemia (21.5%) and changes to the treatment regimen (20.1%). The devices that were most widely used for searching for this information were the computer (41.60%), mobile phone (17.47%) and personal digital assistant (PDA)/tablet (8.30%).

The most highly valued informational content on T2DM-related apps were information about medication (54.4%), dose reminders (47.4%), diet menu plans (41.0%), calorie calculators (35.4%) and physical activity metres (34.8%).

Recording blood glucose levels and other variables

The majority of patients (85.9%) recorded their blood glucose levels. Of the total of patients, the most commonly used formats for record keeping were paper (33.6%), glucometer (27.9%), PC (10.3%) and mobile phone/PDA (4.9%). The results of the subanalysis according to age showed significant differences. Patients aged <65 years used digital formats (PC: χ^2 (df=1)=61.64, p<0.001; mobile phone/PDA: χ^2 (df=1)=33.24, p<0.001;

glucometer: χ^2 (df=1)=15.59; p<0.001) more often than did patients aged ≥65 years (figure 1).

Other T2DM-related data, different from blood glucose levels, were recorded by 14.4% of patients. Of these, 51.0% kept record of their exercise sessions, 45.6% their calorie intake, 32.2% water intake, 21.4% HbA1c levels and 20.8% the number of hypoglycaemia episodes. Of the total patients, the format used to record these data were paper (6.7%), followed by PC (6.2%), glucometer (4.9%) and mobile phone/PDA (2.9%). Again, the use of digital formats was significantly greater among those who were <65 years of age (PC: χ^2 (df=1)=42.89, p<0.001; mobile phone/PDA: χ^2 (df=1)=9.43, p=0.002; glucometer: χ^2 (df=1)=0.53, p=0.463) (figure 1).

The data recorded were shared with the physician by 58.7% of patients, with the nurse (23.3%) and the pharmacist (5.9%), especially by patients under the age of 65 years (physicians: χ^2 (df=1)=5.20, p=0.023; nurse: χ^2 (df=1)=0.03, p=0.842; pharmacists: χ^2 (df=1)=13.81, p<0.001). In addition, 3.0% of the patients shared their data with other patients.

Use of TM in Spain

Small percentage of participants (9.8%) had experience with TM. Compared with naïve patients, those with TM

Table 2 Clinical characteristics of the patients with T2DM

	Total (n=1.036)
Time for diagnosis (% (n))	
<5 years	28.4 (294)
5–10 years	37.6 (390)
>10 years	33.9 (352)
Time from start of treatment (% (n))	
<5 years	33.1 (328)
5–10 years	31.2 (368)
>10 years	29.7 (294)
Missing data	46
Type of T2DM treatment (% (n))	
Oral	87.4 (906)
Oral (monotherapy)	4.5 (4)
Oral (combined treatment)	64.2 (582)
Oral (fixed combination)	31.2 (283)
Injectable (insulin)	38.2 (396)
Injectable (monotherapy)	36.87 (146)
Injectable (combined treatment)	63.13 (250)
Treatment adherence (% (n))	64.00 (663)
T2DM-related health problems (% (n))	
Diabetic foot	8.49 (88)
Diabetic retinopathy	14.67 (152)
Diabetic nephropathy	11.39 (118)
Diabetic neuropathy	19.88 (206)
Diabetic heart disease	12.55 (130)
Comorbidities (% (n))	
Arterial hypertension	46.81 (485)
Hypercholesterolaemia	41.60 (431)
Hypertriglyceridaemia	25.48 (264)
Obesity	45.46 (471)
Repeated cystitis	12.07 (125)

T2DM, type 2 diabetes mellitus.

experience were significantly younger, had a higher level of education, were predominantly in active employment and lived with a partner (table 1). In addition, patients with insulin treatment had more experience in TM than those without insulin (17.7% vs 5.0%; X^2 (df=1)=44.28; $p<0.001$).

Among patients who had no experience in TM, 38.2% expressed a willingness to participate in TM programmes but only 4.7% had been invited to do so. Subanalysis by age (<65 vs ≥65 years) showed that most patients invited to participate were under age of 65 years (3.2% vs 1.5%; X^2 (df=1)=12.56; $p<0.001$). Comparing patient who expressed interest in participating and those who did not, males (table 1) and patients <65 years (60.4% vs 21.9%; X^2 (df=1)=142.59; $p<0.001$) were more interested in TM.

The TM resources that were most used by the patients were SMS/WhatsApp messages (45.1%), telephone calls (42.9%), glucometres with data transmission (41.2%), online platform (37.95%), mobile apps (32.4%) and videoconferences (6.9%) and those with a higher rate of satisfaction were glucometer (100%), videoconferences (100%) SMS/WhatsApp (97%) and online platform (97%).

Perceptions of the benefits of using TM

Patients felt that TM could reduce (quite a bit or a lot) the number of visits to the doctor (47.7%), nurse (43.9%), emergency room (39.9%), pharmacy (35.9%) and the time per visit (43.2%). Compared with naïve patients, the perceptions of patients with prior experience using TM was significantly more positive: visits to the doctor (61.8% vs 46.1%; X^2 (df=4)=14.83; $p=0.005$), visits to the nurse (59.8% vs 42.2%; X^2 (df=4)=15.88; $p=0.003$), visits to the emergency room (59.8% vs 37.7%; X^2 (df=4)=21.39; $p<0.001$), visits to the pharmacy (45.1% vs 34.9%; X^2 (df=4)=16.13; $p=0.003$) and time required for the doctor's visits (63.7% vs 41.0%; X^2 (df=4)=38.51; $p<0.001$) (figure 2).

The majority of patients thought that TM could improve (quite a bit or a lot) all proposed aspects related to T2DM management, except for absences from work (figure 3). In comparison with naïve patients, those with prior TM experience perceived TM provided greater benefits in the improvement of: knowledge about the disease (78.4% vs 60.9%; $p=0.001$; X^2 (df=4)=18.64; $p=0.001$), knowledge of the consequences of not adhering to treatment (77.5% vs 54.7%; X^2 (df=4)=25.47; $p<0.001$), adherence to treatment (74.5% vs 54.6%; X^2 (df=4)=19.76; $p=0.001$), observance of dietary recommendations (67.6% vs 55.6%; X^2 (df=4)=14.38; $p=0.006$) and physical exercise recommendations (76.5% vs 53.1%; X^2 (df=4)=25.47; $p<0.001$), glycaemic control (73.5% vs 58.7%; $p<0.001$; X^2 (df=4)=20.15; $p<0.001$), prevention of hypoglycaemic episodes (77.5% vs 56.9%; X^2 (df=4)=22.18; $p<0.001$), health-related quality of life (78.4% vs 61.3%; X^2 (df=4)=16.36; $p=0.003$) and the reduction of absences from work (65.7% vs 33.4%; $p<0.001$; X^2 (df=4)=57.62; $p<0.001$).

Most patients with TM experience (73.6%) considered that the use of TM had optimised (quite a bit or a lot) the management of their T2DM.

Preferences and satisfaction with TM

With regard to their preferences for TM resources, the majority of patients with experience using TM showed quite a bit or a lot of interest in online platforms with content supervised by professionals (70.6%), online platforms that allow communication with the medical team (70.6%), mobile apps (70.6%), SMS/WhatsApp reminders sent by the healthcare centre or pharmacy (66.7%) and online platforms that allow them to communicate with other patients with similar health problems (56.9%) (figure 4).

Figure 1 Percentage of patients who record their blood glucose levels and other type 2 diabetes mellitus-related data according to technology format and age group. * $P<0.05$ (χ^2 test). PC, personal computer; PDA, personal digital assistant.

The percentage of patients who were satisfied or very satisfied with the use of TM for the management of their T2DM was 70.5%, and 72.5% would participate again in another TM programme.

Improving and promoting TM

Participants who had prior TM experience were asked about possible improvement features. They indicated that it would be necessary or very necessary to improve certain issues, such as explanations about the programmes to help make them easier to use (81.4%), interaction with the medical team (78.4%), time required for recording/transferring data (78.4%), content (76.5%), devices available (74.5%), access to devices at the healthcare centre (72.6%), interaction with other patients (62.8%) and app design (64.7%) (figure 5).

Furthermore, 80.4% of patients would recommend the use of TM. However, in order to boost its use, they considered the following issues to be important: support from

a health professional or experienced patient (79.4%), simple instructions (78.4%), recommendations for its use by the physician (77.5%) or pharmacist (75.5%), availability of the information that is recorded (77.5%), recommendations for its use by friends or relatives (69.6%) and television advertising (58.8%).

DISCUSSION

There is a widespread adoption and use of ICT devices by Spanish households. In 2016, according to the Spanish National Statistics Institute, 81.9% of Spanish households had access to the internet, 96.7% to a mobile phone and 77.1% had a computer or similar device.³⁰ These data are similar to those observed in our study, with the exception of the availability of computers, which in this case was 67.4%.

Figure 2 Distribution of patients according to their perceptions about how telemedicine may reduce the use of healthcare resources. E, experienced; N, naïve; T, total. * $P<0.05$ (χ^2 test; 4 df).

Figure 3 Distribution of patients according to their perceptions about how telemedicine may improve different aspects related to the management of type 2 diabetes mellitus. E, experienced; HRQoL, health-related quality of life; N, naïve; T, total. * $P < 0.05$ (χ^2 test; 4 df).

Internet usage by patients to search for information about their disease has risen over the last decade, increasing from 23.9% in 2004³¹ to 66.9% in 2009.³² For T2DM, a study conducted in 203 patients showed that 28.1% of these patients searched for information on the internet about their disease.³³ In contrast, in our study 46.6% of the patients reported to search information on the internet. In any case, it is apparent that to a greater or lesser extent, many patients with T2DM are interested in expanding their knowledge about their condition and one way to do so is by turning to the internet for information. However, in general, quality of the information about the T2DM available on the internet is poor taken in account the dissemination of patient-oriented evidence that matters,³⁴ which could induce to follow mistaken health habits as in other diseases.³⁵

Another fact to consider when assessing the potential use of TM for the management of T2DM is the level of patient's use of ICTs. A recent study evaluated the relationship between HbA1c levels and the frequency with which patients with T2DM record and share their blood glucose levels with their physician. This study suggested that higher frequency in the recording of these levels did not correlate with a significant decrease in HbA1c levels, but recording and sharing with the physician did.³⁶ In keeping with these findings, 85.9% and 14.4% of the patients who participated in our study recorded their blood glucose levels and other T2DM-related values, respectively. However, the percentage who shared these data with their doctor was 58.7%. Fuji *et al*⁴ found that many patients felt that their doctor already had all the information necessary for the management of their

Figure 4 Patient's point of view: telemedicine (TM) resource preferences (patients with TM experience).

Figure 5 Patient's point of view: items to be improved in telemedicine.

disease, which may explain why many of the patients who record these data do not share the information. It would therefore be important to stress how useful recording and sharing this type of data is and to promote and facilitate the use of the tools needed in order to do so. This may be particularly helpful in patients with a recent diagnosis of T2DM and in patients treated with insulin or oral antidiabetic drugs.

With regard to the clinical benefits of TM use, two recent meta-analyses have shown that, compared with usual care, TM helps to optimise glycaemic control by lowering HbA1c levels.^{13 14}

For all that, beyond the objective results, adherence to this type of programme may be conditioned by patients' perceptions about the utility of TM. In this respect, consistent with our results, a European study conducted with >2000 patients in nine countries suggested that patients perceived TM to be a useful tool for improving management of their disease by saving time in unnecessary commutes.³⁷ In young patients, where hospital attendance may be hampered by working time constraints, TM may be particularly advantageous. With regard to TM resources, Faruque *et al* found that TM programmes that used online platforms and text messages that allowed patient-physician communication were significantly more effective at glycaemic control (HbA1c levels) by allowing treatment to be adjusted.¹³ This was one of the types of platform that aroused most interest in the patients surveyed in our study. These results are in line with the results obtained in a qualitative study that evaluated the implementation of a TM programme for patients with T2DM, which found that interaction with the physician was highly valued by the patients.³⁸

The level of satisfaction with TM that was observed in our study is comparable to that seen in previous studies.

In one of these, the level of satisfaction of patients who had participated in TM programmes conducted in six health centres in the Albacete healthcare region, was high or very high in 96% of cases.²⁰ Similarly, another study found that patients' acceptance was 5.8 (IQR 5.2–6) on a scale from 1 to 6.³⁷

Despite the high level of satisfaction reported in our study, the majority of the patients indicated that in order to promote TM use, some aspects such as providing programmes with simple instructions, quickly record and transfer data and the availability of the information recorded should be improved. In line with these results, Koopman *et al* also identified the demand for simple strategies that are technically stable and easily adapted to daily routines.³⁸ Similarly, availability of the data recorded is in high demand by patients who record their disease-related data.^{4 37}

The profile of patients who are interested in participating in TM programmes is that of a middle-aged male patient, who is in a partnership, has a higher education and is in active employment. From an economic point of view, it has been suggested that the benefit of TM interventions is especially relevant to patients who reside in areas that are far from their healthcare centre^{39 40} and who are typically elderly. However, whereas older patients are less accustomed to recording information using digital formats, investment in especially simple and/or automated technological resources together with greater training in their use could contribute to improve the willingness of elderly patients to participate in TM programmes.

Finally, the study data reveal how low the implementation and promotion of the use of TM for the management of T2DM are in Spain. In this sense, physicians and pharmacists would have a very important role in promoting its

use and identifying which TM resources are best for each patient.

This study shows limitations inherent in its observational design, including susceptibility to bias and confounding, which restrict the ability to define causality.⁴¹ In addition, the results must be interpreted while keeping in mind that all of the issues evaluated, including those related to the patient's clinical situation, are always expressed from the patient's point of view, which increases the possibility that the issues raised here may have different interpretations. Furthermore, snowball technique could lead to potential sampling bias and low sample representativeness. Nevertheless, the wide sample variability observed (clinical and sociodemographic) suggests a limited bias.

CONCLUSION

In Spain, nearly 10% of patients with T2DM have experience with TM and it is well accepted, especially the one based on glucometers. Although the use of TM has been found to offer numerous benefits in the management of T2DM, its implementation and promotion is scant, especially among the elderly. Simple resources, allowing treatment-related data to be recorded as well as providing hygiene and nutrition tips, enabling the adjustment of treatment and the exchange of information with the medical team, stand out as the most suitable for the management of T2DM.

Author affiliations

¹Medical Department, Mylan, Madrid, Spain

²DAP-Cat group. Unitat de Suport a la Recerca Barcelona Ciutat, Institut Universitari d'Investigació en Atenció Primària Jordi Gol (IDIAP Jordi Gol), Barcelona, Spain

³Primary Health Care Center Raval Sud, Gerència d'Àmbit d'Atenció Primària Barcelona Ciutat, Institut Català de la Salut, Barcelona, Spain

⁴Coordinator Working Group on Diabetes SEFAC Pontevedra, Spain

⁵Universidad de Granada, Granada, Spain

⁶Outcomes'10, Castellon, Spain

Acknowledgements The authors would like to thank the patients who participated in this study. The authors would also like to thank Ana López (Assistant from Mylan Medical Department) and Miriam Prades (HE&OR Manager from Outcomes'10) for their assistance in the study.

Contributors This article is part of a Doctoral Thesis of PR-F on behalf of the Doctoral Programme in Pharmacy, University of Granada (Spain). All authors have contributed in all the tasks of the study (design, start-up, analysis of the results and manuscript elaboration).

Funding The study was sponsored by BGP Products Operations SL (Mylan). JF-N, JAF-P and FM-M state they have no conflict of interest. PR-F and MLO-P work at BGP Products Operations SL (Mylan). HDdP works for an independent research organisation (Outcomes'10), which has received fees for its contribution to the development and coordination of the project and to the writing of this manuscript.

Disclaimer The funding source had no role in the design, conduct or reporting of the study.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study protocol was approved by the ethics committee of the 'Hospital Universitario Puerta de Hierro' (Madrid, Spain).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No unpublished data for this article are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Soriguer F, Goday A, Bosch-Comas A, *et al*. Prevalence of diabetes mellitus and impaired glucose regulation in Spain: the Di@bet.es Study. *Diabetologia* 2012;55:88–93.
2. Guariguata L, Whiting DR, Hambleton I, *et al*. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract* 2014;103:137–49.
3. Christensen K, Doblhammer G, Rau R, *et al*. Ageing populations: the challenges ahead. *Lancet* 2009;374:1196–208.
4. Fuji KT, Abbott AA, Galt KA. A qualitative study of how patients with type 2 diabetes use an electronic stand-alone personal health record. *Telemed J E Health* 2015;21:296–300.
5. Bashshur RL, Shannon GW, Smith BR, *et al*. The empirical evidence for the telemedicine intervention in diabetes management. *Telemed J E Health* 2015;21:321–54.
6. Lim D, Oakley AM, Rademaker M. Better, sooner, more convenient: a successful teledermatology service. *Australas J Dermatol* 2012;53:22–5.
7. Caffery LJ, Farjian M, Smith AC. Telehealth interventions for reducing waiting lists and waiting times for specialist outpatient services: A scoping review. *J Telemed Telecare* 2016;22:504–12.
8. Capoccia K, Odegard P, Letassy N. Medication adherence with diabetic medication. *Diabetes Educ* 2015. Available: <http://tde.sagepub.com/content/early/2015/12/03/0145721715619038.long>
9. Hamine S, Gerth-Guyette E, Faulx D, *et al*. Impact of mHealth chronic disease management on treatment adherence and patient outcomes: a systematic review. *J Med Internet Res* 2015;17:e52.
10. Anglada-Martínez H, Riu-Viladoms G, Martín-Conde M, *et al*. Does mHealth increase adherence to medication? Results of a systematic review. *Int J Clin Pract* 2015;69:9–32.
11. Di Cerbo A, Morales-Medina JC, Palmieri B, *et al*. Narrative review of telemedicine consultation in medical practice. *Patient Prefer Adherence* 2015;9:65–75.
12. Ciemins E, Coon P, Peck R, *et al*. Using telehealth to provide diabetes care to patients in rural Montana: findings from the promoting realistic individual self-management program. *Telemed J E Health* 2011;17:596–602.
13. Faruque LI, Wiebe N, Ehteshami-Afshar A, *et al*. Effect of telemedicine on glycated hemoglobin in diabetes: a systematic review and meta-analysis of randomized trials. *CMAJ* 2017;189:E341–E364.
14. Su D, Zhou J, Kelley MS, *et al*. Does telemedicine improve treatment outcomes for diabetes? A meta-analysis of results from 55 randomized controlled trials. *Diabetes Res Clin Pract* 2016;116:136–48.
15. Wayne N, Perez DF, Kaplan DM, *et al*. Health Coaching Reduces HbA1c in Type 2 Diabetic Patients From a Lower-Socioeconomic Status Community: A Randomized Controlled Trial. *J Med Internet Res* 2015;17:e224.
16. Crowley MJ, Edelman D, McAndrew AT, *et al*. Practical Telemedicine for Veterans with Persistently Poor Diabetes Control: A Randomized Pilot Trial. *Telemed J E Health* 2016;22:376–84.
17. Kuo A, Dang S. Secure Messaging in Electronic Health Records and Its Impact on Diabetes Clinical Outcomes: A Systematic Review. *Telemed J E Health* 2016;22:769–77.
18. Rasmussen OW, Lauszus FF, Loekke M. Telemedicine compared with standard care in type 2 diabetes mellitus: A randomized trial in an outpatient clinic. *J Telemed Telecare* 2016;22:363–8.
19. Quinn CC, Khokhar B, Weed K, *et al*. Older Adult Self-Efficacy Study of Mobile Phone Diabetes Management. *Diabetes Technol Ther* 2015;17:455–61.
20. López-Torres J, Rabanales J, Fernández R, *et al*. Results of a telemedicine program for primary care patients with type 2 diabetes [Resultados de un programa de telemedicina para pacientes con diabetes tipo 2 en atención primaria]. *Gac Sanit* 2015;29:55–8.
21. Pérez-Ferre N, Galindo M, Fernández MD, *et al*. A Telemedicine system based on Internet and short message service as a new approach in the follow-up of patients with gestational diabetes. *Diabetes Res Clin Pract* 2010;87:e15–e17.
22. Peña NV, Torres M, Cardona JA, *et al*. Impact of telemedicine assessment on glycemic variability in children with type 1 diabetes mellitus. *Diabetes Technol Ther* 2013;15:136–42.

23. Azar M, Gabbay R. Web-based management of diabetes through glucose uploads: has the time come for telemedicine? *Diabetes Res Clin Pract* 2009;83:9–17.
24. Rojahn K, Laplante S, Sloand J, et al. Remote Monitoring of Chronic Diseases: A Landscape Assessment of Policies in Four European Countries. *PLoS One* 2016;11:e0155738.
25. Klonoff DC. Telemedicine for Diabetes. *J Diabetes Sci Technol* 2016;10:3–5.
26. Shahrokni A, Mahmoudzadeh S, Saeedi R, et al. Older People with Access to Hand-Held Devices: Who Are They? *Telemedicine and e-Health* 2015;21:550–6.
27. Goodman LA. Snowball sampling. *Ann Math Stat* 1961;32:148–70.
28. Instituto nacional de estadística. Cifras de población (>18) [Internet]. 2015. Available: <http://www.ine.es/jaxiT3/Datos.htm?t=9663> (cited 1 Nov 2015).
29. Czaja SJ, Lee CC. Information technology and older adults. In: Sears A, Jacko J, eds. *Human-Computer Interaction: Designing for Diverse Users and Domains*, 2009:18–30.
30. Instituto Nacional de Estadística. Encuesta sobre equipamiento y uso de tecnologías de información y comunicación en los hogares. In: INEbase [Internet]. 2016. Available: http://www.ine.es/dyngs/INEbase/es/operacion.htm?c=Estadistica_C&cid=1254736176741&menu=ultiDatos&idp=1254735976608 (cited 1 Feb 2017).
31. Ortego Centeno N, Callejas Rubio JL, Benticuaga Martínez MN, et al. Utilización de Internet como fuente de información por los pacientes de una consulta de Enfermedades Autoinmunes Sistémicas. *Revista Clínica Española* 2005;205:255–8.
32. Ortego Centeno N, Barnosi Marín AC, Simeón Aznar CP, et al. Búsqueda de información en Internet por parte de pacientes de diferentes Comunidades Autónomas con enfermedades autoinmunitarias crónicas. *Medicina Clínica* 2009;133:467–71.
33. Kalantzi S, Kostagiolas P, Kechagias G, et al. Information seeking behavior of patients with diabetes mellitus: a cross-sectional study in an outpatient clinic of a university-affiliated hospital in Athens, Greece. *BMC Res Notes* 2015;8:48.
34. Giménez-Pérez G, Caixàs A, Giménez-Palop O, et al. Dissemination of 'patient-oriented evidence that matters' on the Internet: the case of Type 2 diabetes treatment. *Diabet Med* 2005;22:688–92.
35. Zhang D, Schumacher C, Harris MB, et al. The Quality and Readability of Information Available on the Internet Regarding Lumbar Fusion. *Global Spine J* 2016;6:133–8.
36. Chow N, Shearer D, Aydin Plaa J, et al. Blood glucose self-monitoring and internet diabetes management on A1C outcomes in patients with type 2 diabetes. *BMJ Open Diabetes Res Care* 2016;4:e000134.
37. Dario C, Luisotto E, Dal Pozzo E, et al. Assessment of Patients' Perception of Telemedicine Services Using the Service User Technology Acceptability Questionnaire. *Int J Integr Care* 2016;16:13–14.
38. Koopman RJ, Wakefield BJ, Johanning JL, et al. Implementing home blood glucose and blood pressure telemonitoring in primary care practices for patients with diabetes: lessons learned. *Telemed J E Health* 2014;20:253–60.
39. Fountoulakis S, Papanastasiou L, Gryparis A, et al. Impact and duration effect of telemonitoring on HbA1c, BMI and cost in insulin-treated Diabetes Mellitus patients with inadequate glycemic control: A randomized controlled study. *Hormones* 2015;14:632–43.
40. Chérrez-Ojeda I, Vanegas E, Calero E, et al. What Kind of Information and Communication Technologies Do Patients with Type 2 Diabetes Mellitus Prefer? An Ecuadorian Cross-Sectional Study. *Int J Telemed Appl* 2018;2018:1–8.
41. Yang W, Zilov A, Soewondo P, et al. Observational studies: going beyond the boundaries of randomized controlled trials. *Diabetes Res Clin Pract* 2010;88 Suppl 1:S3–S9.