

## Supplemental Data

**Table 1.** Multivariate analysis: risk factors associated with Stroke/TIA; Excessive PACs defined as >100 PACs/24 hours

<b>Characteristic</b>	<b>Stroke/TIA</b>	
	<i>Odds Ratio (95% CI)</i>	<i>P value</i>
<b>Female</b>	1.54 (1.10 - 2.14)	<0.05
<b>Hypertension</b>	1.46 (1.04 - 2.07)	<0.05
<b>Smoking</b>	1.57 (1.06 - 2.35)	<0.05
<b>Excessive PACs</b>	1.57 (1.10 - 2.25)	<0.05

CI = confidence interval; PAC = premature atrial complexes.

*Variables adjusted in the multivariate model: Age, sex, excessive premature atrial complexes, hypertension, diabetes mellitus, smoking, dyslipidaemia, peripheral vascular disease and sleep apnoea*

**Table 2.** Multivariate analysis: risk factors associated with Stroke/TIA; Excessive PACs defined as >700 PACs/24 hours or an atrial run >20 beats

<b>Characteristic</b>	<b>Stroke/TIA</b>	
	<i>Odds Ratio (95% CI)</i>	<i>P value</i>
<b>Female</b>	1.63 (1.16 - 2.27)	<0.01
<b>Hypertension</b>	1.45 (1.03 - 2.05)	<0.05
<b>Smoking</b>	1.59 (1.06 - 2.37)	<0.05
<b>Excessive PACs</b>	3.22 (1.78 - 5.83)	<0.01

CI = confidence interval; PAC = premature atrial complexes.

*Variables adjusted in the multivariate model: Age, sex, excessive premature atrial complexes, hypertension, diabetes mellitus, smoking, dyslipidaemia, peripheral vascular disease and sleep apnoea*