

BMJ Open Correlates of the double burden of malnutrition among women: an analysis of cross sectional survey data from sub-Saharan Africa

Dickson Abanimi Amugsi,¹ Zacharie Tsala Dimbuene,^{2,3} Catherine Kyobutungi¹

To cite: Amugsi DA, Dimbuene ZT, Kyobutungi C. Correlates of the double burden of malnutrition among women: an analysis of cross sectional survey data from sub-Saharan Africa. *BMJ Open* 2019;**9**:e029545. doi:10.1136/bmjopen-2019-029545

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-029545>).

Received 30 January 2019
Revised 24 May 2019
Accepted 30 May 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Maternal and Child Wellbeing Unit, African Population and Health Research Center, Nairobi, Kenya

²Population Sciences and Development, University of Kinshasa, Kinshasa, The Democratic Republic of the Congo

³Social Analysis and Modeling Division, Statistics Canada, Ottawa, Canada

Correspondence to

Dr Dickson Abanimi Amugsi; damugsi2002@yahoo.com

ABSTRACT

Objective To investigate the correlates of the double burden of malnutrition (DBM) among women in five sub-Saharan African countries.

Design Secondary analysis of Demographic and Health Surveys (DHS). The outcome variable was body mass index (BMI), a measure of DBM. The BMI was classified into underweight (BMI <18.50 kg/m²), normal weight (18.50–24.99 kg/m²), overweight (25.0–29.9 kg/m²) and obesity (≥30.0 kg/m²).

Settings Ghana, Nigeria, Kenya, Mozambique and Democratic Republic of Congo (DRC).

Subjects Women aged 15–49 years (n=64698).

Results Compared with normal weight women, number of years of formal education was associated with the likelihood of being overweight and obese in Ghana, Mozambique and Nigeria, while associated with the likelihood of being underweight in Kenya and Nigeria. Older age was associated with the likelihood of being underweight, overweight and obese in all countries. Positive associations were also observed between living in better-off households and overweight and obesity, while a negative association was observed for underweight. Breastfeeding was associated with less likelihood of underweight in DRC and Nigeria, obesity in DRC and Ghana, overweight in Kenya and overweight and obesity in Mozambique and Nigeria relative to normal weight.

Conclusions Our analysis reveals that in all the countries, women who are breastfeeding are less likely to be underweight, overweight and obese. Education, age and household wealth index tend to associate with a higher likelihood of DBM among women. Interventions to address DBM should take into account the variations in the effects of these correlates.

INTRODUCTION

The double burden of malnutrition (DBM), which is the coexistence of both undernutrition and overnutrition in the same population across the life course,^{1,2} is a global public health problem.^{1,2} Data from the WHO indicate that in 2014, 1.9 billion adults aged 18 years and above were overweight, while over 600 million were obese globally.¹ Similarly, in the same year, 462 million adults were

Strengths and limitations of this study

- Large nationally representative samples used, thereby providing more robust estimates of observed associations.
- The height and weight data used to compute the body mass index were objectively measured, reducing possible misclassification.
- Use of multicountry data helped unmask differences and commonalities in the effects of the correlates on double burden of malnutrition (DBM) across countries.
- The use of cross-sectional surveys may not allow to establish causation.
- Due to data limitation, the DBM was examined only at the population or national level.

underweight and 264 million women of reproductive age were affected by iron-amenable anaemia.^{2,3} These key indicators of DBM are also increasing globally, with the low-income and middle income countries (LMICs) being the most affected.⁴ For example, while globally, obesity has doubled in the last three decades, it has tripled in LMICs in just two decades.⁵ An analysis of survey data from 24 African countries spanning 25 years revealed that overweight and obesity among women are on the rise.⁶ Also, comparative analysis of data on women and men in the LMICs showed that DBM tends to disproportionately affect women than men.^{7,8} The vulnerability of women to DBM may be attributed to their high nutritional requirements for pregnancy and lactation and also because of gender inequalities in poverty.⁸ Further evidence suggests that if micronutrient deficiencies are taken into account, Africa is in fact experiencing a triple burden of malnutrition.^{2,3} It has been estimated that almost 50% of pregnant women in Africa suffer from anaemia, which increases the risk of death for themselves as well as their unborn babies.²

Furthermore, the contribution of DBM to the burden of disease has been documented. The available evidence suggests that underweight and obesity are among the top 10 leading risk factors for the global burden of disease.^{9 10} Furthermore, more recent WHO data have identified underweight among the top four risks factors for the burden of disease in the world, as measured in disability-adjusted life years (DALYs)—contributing up to 6% of global DALYs.¹⁰ The relationship between maternal and child weight and the consequences on disease incidence later in life have also been documented. For example, being overweight as a mother is associated with overweight and obesity in their offspring.^{2 11} Rapid weight gain early in life may predispose to long-term excessive weight or obesity. And as the evidence suggests, obesity is an important underlying cause of many non-communicable diseases (NCDs), including hypertension, diabetes, cancer, stroke and ischaemic heart disease.^{3 5 11} These diseases are responsible for most of the deaths worldwide, with LMICs disproportionately bearing the brunt, where 80% of the NCD deaths occur.²⁻⁴ Experts warned that unless countries in Africa start enacting measures to tackle the DBM affecting the continent, the road towards universal healthcare (UHC) will be marred with obstacles as will the aspiration to achieve health and well-being for all by 2030.¹²

It is important to underscore that DBM can exist at the individual, household and population levels.² For instance, at the individual level, obesity can occur with deficiency of one or various vitamins and minerals, or overweight in an adult who was stunted during childhood. At the household level, a mother may be overweight or anaemic and a child or grandparent is underweight. DBM at the population level occurs when there is a prevalence of both undernutrition and overnutrition in the same community, nation or region.² Since it will be difficult to determine individual and within households DBM using these data, our definition of DBM is at population or country level, whereby underweight and overweight/obesity co-exist in the same country. Undoubtedly, the DBM offers an important opportunity for use of multidimensional approaches in tackling malnutrition in all its forms. Addressing the DBM will be key to achieving the Sustainable Development Goals (in particular Goal 2 and Target 3.4) and the Commitments of the Rome Declaration on Nutrition, within the UN Decade of Action on Nutrition.¹ However, as a starting point, countries need to understand the key correlates of DBM. The present paper attempts to elucidate these correlates.

The factors influencing DBM are complex, ranging from biological to environmental factors.² Some of these factors may include poor water and sanitation systems, weak public health systems thereby thwarting efforts to reduce undernutrition.² Also, increasing urbanisation and changing dietary patterns and sedentary lifestyles, income level, older age, household wealth, higher education, place of residence among others have been identified as key contributing factors to the DBM epidemic.¹³⁻¹⁶

For example, Kamal and colleagues observed in their study that household wealth index and place of residence are key predictors of the DBM among women in Bangladesh.¹⁷ Women from the poorest wealth quintile were more likely to suffer from DBM relative to those from the richest wealth quintile. Besides, marital status, age at first childbirth, parity, household size and food security were also found to play a critical role in the DBM epidemic.^{14 17 18} The above-referenced studies used data from a single country, masking differences and commonalities of the effects of the correlates on DBM across countries. Suffice to add that Neupane and colleagues¹⁹ attempted to investigate the problem of DBM using data sets from 32 sub-Saharan African (SSA) countries. This study, however, fell short of addressing DBM as it focused only on overweight and obesity. Second, the referenced study used only three variables (wealth quintile, place of residence and education) to predict overweight and obesity. The present study filled these gaps by simultaneously analysing underweight, overweight and obesity, using a comprehensive list of predictor variables, to provide a robust picture of the correlates of DBM in SSA.

Given the anticipated long-term effects of DBM, the factors that are associated with being underweight, overweight or obese should be considered while formulating effective interventions to address DBM among women.^{15 16} This stresses the need for prevention strategies targeted at addressing all forms of malnutrition. The present study is well positioned to provide evidence on the key correlates of DBM in SSA, which is currently lacking in the region. Understanding the role of these risk factors is key to developing clear and effective strategies for improving public health in SSA. The overarching aim of our study is to examine the correlates of DBM among women in five SSA countries.

METHODOLOGY

Sources of data and sampling procedure

Design and data sources

The study used the recent Demographic and Health Surveys (DHS)²⁰ data from Ghana,²¹ Kenya (2014), Nigeria (2013), Mozambique (2011) and Democratic Republic of Congo (DRC) (2013–2014). The selection of these five countries was informed by our previous analysis.²² The DHS data are nationally representative, repeated cross-sectional household surveys collected primarily in LMICs approximately every 5 years, using standardised questionnaires to enable cross-country comparisons.^{23 24} The design of the DHS surveys is identical across all participating countries, making possible the comparisons between and across countries. The DHS utilises a two-stage sample design.^{21 25-28} The first stage involves the selection of sample points or clusters from an updated master sampling frame constructed from National Population and Housing Census of the respective countries. The clusters are selected using systematic sampling with probability proportional to size.

Table 1 Characteristics of the BMI samples

Variables	DRC	Ghana	Kenya	Mozambique	Nigeria
	Mean/%	Mean/%	Mean/%	Mean/%	Mean/%
BMI					
BMI=18.50–24.99 (normal weight)	73.50	59.20	62.00	77.60	66.10
BMI <18.50 (underweight)	12.70	5.30	11.80	5.70	8.70
BMI=25–29.99 (overweight)	11.50	23.40	18.90	13.70	18.30
BMI≥30 (obesity)	2.30	12.20	7.30	3.00	7.00

BMI, body mass index; DRC, Democratic Republic of Congo.

Household listing is then conducted in all the selected clusters to provide a sampling frame for the second stage selection of households. The second stage of selection involves the systematic sampling of the households listed in each cluster and households to be included in the survey are randomly selected from the list. The rationale for the second stage selection is to ensure adequate numbers of completed individual interviews to provide estimates for key indicators with an acceptable precision. All men and women aged 15–59 and 15–49, respectively, in the selected households (men in half of the households) were eligible to participate in the surveys if they were either usual residents of the household or visitors present in the household on the night before the survey. We limited our analyses to women aged 15–49 years in all countries and who have complete anthropometry data. The samples for the respective countries are as follows: DRC (9506), Ghana (3012), Kenya (9993), Mozambique (11017) and Nigeria (31 170).

ETHICS STATEMENT

The DHS obtained ethical clearance from Government recognised Ethical Review Committees/Institutional Review Boards of the respective countries as well as the Institutional Review Board of ICF International, USA, before the surveys were conducted. Written informed consent was obtained from the women before participation. The authors of this paper sought and obtained permission from the DHS programme for the use of the data. The data were completely anonymised and therefore the authors did not seek further ethical clearance before their use.

PATIENT AND PUBLIC INVOLVEMENT STATEMENT

We used completely anonymised secondary data for the analysis. Therefore, no patients or public involvement can be reported.

Outcome and predictor variables

Outcome variables

The outcome variable of interest was women body mass index (BMI) derived from results of height and weight measurements. The height and weight were

measured objectively by trained field technicians using standard techniques.²⁴ Weight measurements were taken using electronic Seca scales with a digital screen, while height measurements were taken using a stadiometer produced by Shorr Productions. BMI, also referred to as Quetelet's Index,²⁹ was derived by dividing weight in kg by the squared height in metres. Based on the BMI (kg/m²) estimates, and according to WHO guidelines,³⁰ the participants were classified as underweight (BMI <18.50 kg/m²), normal weight (18.50–24.99 kg/m²), overweight (25.0–29.9 kg/m²) and obese (≥30.0 kg/m²). The normal weight (18.50–24.99 kg/m²) was used as reference category in the analysis.

Predictor variables

The predictor variables of interest used in the analysis included, women's age, education, employment status, breastfeeding status, parity, place of residence, marital status, women decision making autonomy (decision on large household purchases and decision on health), household size, frequency of watching TV and household wealth index. These potential correlates were identified based on literature search and further subjected to bivariate analysis to establish their relationship with the DBM indicators. All statistically significant variables were included in the multi-variable analysis.

Analytical strategy

We utilised multinomial logistic regression (MLR) in the analysis. MLR approach was contemplated to be suitable as the outcome measure is polychotomous by nature. Further, the MLR was considered attractive analytical technique because it does not assume normality, linearity, or homoscedasticity.³¹ In MLR, we observe vectors $Y=(y_1, y_2, \dots, y_{k+1})^T$; $y_i=0$ for all i , and one j with $y_j=1$ and corresponding probability p_j , implying

$$EY=p, \text{Cov } Y=\Lambda_p = \begin{pmatrix} p_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & p_{k+1} \end{pmatrix} \quad (1)$$

The multinomial logistic regression is given by

Table 2 Characteristics of the samples, categorical and continuous variables

Variables	DRC		Ghana		Kenya		Mozambique		Nigeria	
	Mean/%	SD	Mean/%	SD	Mean/%	SD	Mean/%	SD	Mean/%	SD
Women's education (in years)	5.2	3.8	6.0	4.9	7.6	4.0	3.1	3.3	4.7	5.7
Women's age (in years)	29.2	6.8	30.6	6.8	28.6	6.4	28.6	7.3	29.4	7.0
Household wealth index										
Poorest	22.4		22.7		23.4		23.6		23.5	
Poorer	23.0		19.8		20.3		21.4		23.1	
Middle	20.6		20.6		18.5		20.0		18.9	
Rich	18.6		18.7		18.2		19.9		17.8	
Richest	15.5		18.2		19.6		15.1		16.8	
Respondent currently working (yes)	74.9		78.8		63.6		39.3		68.9	
Place of residence (urban)	29.9		45.4		36.1		27.5		35.0	
Household head (female)	21.6		26.0		30.4		30.3		9.7	
Parity	4.4	2.6	3.6	2.1	3.5	2.3	3.9	2.3	4.3	2.6
Household size	6.8	2.9	5.6	2.6	5.6	2.4	6.0	2.6	7.0	3.6
Frequency of watching TV										
Not at all	66.4		19.6		22.0		34.5		39.8	
Less than once a week	14.6		32.7		12.5		23.2		25.0	
At least once a week	19.0		47.7		65.5		42.3		35.2	
Currently breastfeeding (yes)	68.6		57.8		54.0		60.8		54.8	
Marital status										
Never married	4.3		7.1		7.1		4.2		1.6	
Married or cohabiting	87.2		86.5		83.7		84.3		95.9	
Formerly in union	8.5		6.4		9.2		11.5		2.6	
Decision on large household purchases										
Respondent alone	8.7		20.8		29.1		17.2		4.7	
Both respondent and partner	31.0		42.7		34.2		39.1		29.2	
Partner alone	47.3		22.2		20.1		27.5		61.6	
Someone else/other	0.2		0.9		0.4		0.6		0.2	
Decision on respondent's health										
Respondent alone	13.8		16.3		14.7		10.0		4.3	
Both Respondent and partner	37.2		44.8		42.2		38.2		28.4	
Partner alone	35.6		24.4		26.3		35.4		62.8	
Someone else/other	0.3		1.0		0.4		0.7		0.2	

DRC, Democratic Republic of Congo.

$$p_i = \frac{\exp(\pi^{(i)T} x)}{1 + \sum_{j=1}^k \exp(\pi^{(j)T} x)} \text{ for } i=1, \dots, k \quad (2)$$

$$p_{k+1} = \frac{1}{1 + \sum_{j=1}^k \exp(\pi^{(j)T} x)} \quad (3)$$

where $x = (x_1, x_2, \dots, x_m)^T$ is the vector of covariates, and $\Pi^{(i)}$ is the parameter vector corresponding to the i -th response category. In Equation (3), the parameters are

set to zero and allows computing the probability for the base category in the MLR.

Because of the normalisation condition,

$$\sum_{j=1}^k P(y^{(j)} = 1 | x, \pi) = 1, \quad (4)$$

the weight vector of one of the classes need not to be estimated without loss of generality, in this case the $(j+1)$ -th category. To perform maximum likelihood (ML),

one simply maximises the log-likelihood function using Equation (5),

$$\log \prod_{j=1}^{k+1} p_j^{y_j} = \sum_{j=1}^k y_j \pi^{(i)T} x - \log \left[1 + \sum_{j=1}^k \exp \left(\pi^{(i)T} x \right) \right] \quad (5)$$

The MLR model was constructed to investigate the net effects of the correlates on underweight, overweight and obesity. Using a BMI category of 18.5–24.99 kg/m² (normal weight) as the reference, a set of logistic regressions for underweight, overweight and obese

categories was estimated in which, each of the categories was contrasted with the reference category. All covariates were simultaneously entered into the model. Results were presented in the form of coefficients with significance levels and 95% CI.

RESULTS

Descriptive

The tables 1 and 2 present the characteristics of the samples. The results in table 1 show that among the

Table 3 Multivariable analysis of the association between sociodemographic correlates and malnutrition, DRC

Variables	Underweight	Overweight	Obesity
	(BMI<18.50)	(BMI=25–29.99)	(BMI≥30)
Women's education (in years)	0.030* (–0.004 to 0.064)	–0.011 (–0.052 to 0.030)	–0.023 (–0.093 to 0.047)
Women's age (in years)	0.041*** (0.015 to 0.068)	0.034** (0.007 to 0.060)	–0.045 (–0.104 to 0.015)
Household wealth index			
Poorest (reference)			
Poorer	–0.009 (–0.306 to 0.288)	–0.069 (–0.488 to 0.351)	–0.241 (–1.575 to 1.094)
Middle	–0.144 (–0.534 to 0.247)	0.421** (0.039 to 0.804)	0.570 (–0.529 to 1.669)
Rich	–0.480** (–0.942 to –0.018)	0.803* (0.345 to 1.261)	2.365*** (1.225 to 3.505)
Richest	–1.017*** (–1.564 to –0.471)	1.486* (0.985 to 1.987)	4.014*** (2.562 to 5.465)
Woman currently working (yes)	0.226 (–0.052 to 0.504)	–0.462*** (–0.794 to –0.131)	–0.350 (–0.850 to 0.149)
Place of residence (urban)	–0.246 (–0.601 to 0.108)	0.034 (–0.350 to 0.417)	–0.094 (–0.810 to 0.622)
Sex of household head (female)	–0.0453 (–0.377 to 0.286)	0.167 (–0.144 to 0.478)	0.363 (–0.380 to 1.105)
Parity	–0.122*** (–0.199 to –0.045)	0.0335 (–0.040 to 0.107)	0.350*** (0.174 to 0.525)
Household size	–0.019 (–0.066 to 0.028)	–0.008 (–0.048 to 0.031)	–0.052 (–0.158 to 0.056)
Frequency of watching TV			
Not at all (reference)			
Less than once a week	0.089 (–0.251 to 0.429)	0.165 (–0.192 to 0.522)	–0.209 (–0.912 to 0.494)
At least once a week	–0.163 (–0.623 to 0.298)	0.180 (–0.140 to 0.500)	0.149 (–0.373 to 0.671)
Currently breastfeeding	0.327*** (0.082 to 0.572)	–0.010 (–0.361 to 0.164)	–0.512** (–1.020 to –0.005)
Marital status			
Never in union (reference)			
Married or cohabiting	–0.766* (–1.567 to 0.0347)	0.809* (0.132 to 1.486)	0.248 (–1.131 to 1.626)
Formerly in union	–0.357 (–1.126 to 0.413)	0.0680 (–0.583 to 0.718)	–0.549 (–1.956 to 0.859)
Decision on large household purchases			
Respondent alone (reference)			
Both respondent and partner	0.186 (–0.292 to 0.664)	0.155 (–0.291 to 0.601)	0.885* (–0.154 to 1.923)
Partner alone	0.436* (–0.059 to 0.931)	0.098 (–0.325 to 0.522)	0.423 (–0.370 to 1.215)
Someone else/other	–0.787 (–2.567 to 0.993)	–3.160*** (–5.513 to –0.808)	–12.640*** (–14.160 to –11.130)
Decision on respondent's health			
Respondent alone (reference)			
Both respondent and partner	–0.180 (–0.599 to 0.239)	–0.332 (–0.731 to 0.067)	–0.743* (–1.594 to 0.109)
Partner alone	–0.176 (–0.552 to 0.201)	–0.525** (–0.933 to –0.116)	–0.602* (–1.282 to 0.078)
Someone else/other	1.672** (0.073 to 3.272)	0.687 (–1.990 to 3.363)	–14.200*** (–15.970 to –12.430)
Observations	9506	9506	9506

CI in parentheses.

*P<0.1; **P<0.05; ***P<0.01.

BMI, body mass index; DRC, Democratic Republic of Congo.

Table 4 Multivariable analysis of the association between sociodemographic correlates and malnutrition, Ghana

Variables	Underweight	Overweight	Obesity
	BMI<18.50	BMI=25–29.99	BMI≥30
Women's education (in years)	-0.022 (-0.080 to 0.036)	0.074*** (0.037 to 0.110)	0.028 (-0.014 to 0.070)
Women's age (in years)	0.054** (0.005 to 0.103)	0.0405** (0.008 to 0.073)	0.083*** (0.048 to 0.118)
Household wealth index			
Poorest (reference)			
Poorer	-0.398 (-0.972 to 0.177)	0.367 (-0.129 to 0.864)	1.390*** (0.582 to 2.197)
Middle	-1.050*** (-1.768 to -0.331)	0.893* (0.340 to 1.447)	2.804*** (1.947 to 3.662)
Rich	-0.529 (-1.426 to 0.369)	1.436* (0.878 to 1.995)	3.591*** (2.615 to 4.568)
Richest	-1.788*** (-3.052 to -0.523)	1.271* (0.555 to 1.988)	4.121*** (3.125 to 5.117)
Woman currently working (yes)	-0.125 (-0.713 to 0.463)	0.121 (-0.238 to 0.481)	0.0769 (-0.424 to 0.577)
Place of residence (urban)	0.463* (-0.0488 to 0.975)	0.113 (-0.250 to 0.477)	-0.073 (-0.582 to 0.435)
Sex of household head (female)	0.409 (-0.276 to 1.093)	0.0971 (-0.309 to 0.503)	0.293 (-0.173 to 0.759)
Parity	-0.205** (-0.372 to -0.038)	0.0719 (-0.055 to 0.199)	0.0324 (-0.118 to 0.183)
Household size	0.021 (-0.045 to 0.086)	-0.047 (-0.121 to 0.027)	0.035 (-0.077 to 0.147)
Frequency of watching TV			
Not at all (reference)			
Less than once a week	-0.043 (-0.604 to 0.518)	0.231 (-0.240 to 0.703)	0.673* (0.0003 to 1.345)
At least once a week	-0.233 (-0.754 to 0.287)	0.248 (-0.184 to 0.679)	0.756** (0.146 to 1.365)
Currently breastfeeding (yes)	0.136 (-0.364 to 0.637)	-0.177 (-0.525 to 0.171)	-0.471** (-0.832 to -0.110)
Marital status			
Never in union (reference)			
Married or Cohabiting	-0.536 (-1.843 to 0.772)	-0.259 (-0.904 to 0.386)	0.295 (-0.860 to 1.451)
Formerly in union	0.271 (-0.837 to 1.378)	0.420 (-0.286 to 1.126)	1.435* (0.400 to 2.470)
Decision on large household purchases			
Respondent alone (reference)			
Both respondent and partner	0.464 (-0.555 to 1.483)	-0.0143 (-0.523 to 0.495)	-0.0496 (-0.657 to 0.557)
Partner alone	0.664 (-0.217 to 1.546)	-0.0223 (-0.556 to 0.512)	-0.265 (-0.917 to 0.386)
Someone else/other	-1.212 (-3.637 to 1.213)	-0.0290 (-1.485 to 1.426)	0.752 (-1.161 to 2.665)
Decision on respondent's health			
Respondent alone (reference)			
Both respondent and partner	0.0525 (-0.868 to 0.973)	0.785* (0.280 to 1.290)	0.787* (0.166 to 1.408)
Partner alone	0.119 (-0.643 to 0.881)	0.526* (-0.0457 to 1.098)	0.915* (0.156 to 1.675)
Someone else/other	-0.714 (-2.832 to 1.403)	-1.544 (-3.520 to 0.433)	-13.43*** (-15.04 to -11.81)
Observations	3012	3012	3012

CI in parentheses.

*P<0.1; **P<0.05; ***P<0.01.

BMI, body mass index.

five countries, Mozambique had the highest number of normal weight women (78%) followed by DRC (74%), with Ghana having the lowest (59%). Kenya (12%) and DRC (13%) had the highest prevalence of underweight women, while Ghana had the highest number of overweight (23%) and obese (12%) women. In all the countries analysed, the prevalence of overweight and obesity had overtaken underweight. In table 2, women in Kenya had more years of education, while Mozambique had women with the least years of education. The age of study participants was similar across all the five countries and

ranged from 28 years in Mozambique and Kenya to 30 years in Ghana. Further, among all the countries, Ghana had the highest number of women who were working (79%), while Mozambique had the lowest (39%).

Multivariable

The tables 3–7 present the MLR results of the correlates of DBM among women in DRC, Ghana, Kenya, Mozambique and Nigeria. The results showed a significant positive relationship between women's years' of education and overweight in Ghana, both overweight and obesity in Mozambique and

Table 5 Multivariable analysis of the association between sociodemographic correlates and malnutrition, Kenya

Variables	Underweight	Overweight	Obesity
	(BMI<18.50)	(BMI=25.290.99)	(BMI≥30)
Women's education (in years)	-0.113*** (-0.148 to -0.078)	0.013 (-0.016 to 0.042)	0.013 (-0.030 to 0.056)
Women's age (in years)	0.026** (0.001 to 0.052)	0.066*** (0.046 to 0.089)	0.120*** (0.087 to 0.153)
Household wealth index			
Poorest (reference)			
Poorer	-0.488*** (-0.805 to -0.172)	0.279* (-0.025 to 0.582)	0.633* (0.080 to 1.187)
Middle	-0.500*** (-0.869 to -0.130)	0.609*** (0.261 to 0.957)	0.790* (0.268 to 1.311)
Rich	-0.940*** (-1.376 to -0.504)	0.927* (0.578 to 1.276)	1.690*** (1.164 to 2.216)
Richest	-1.307*** (-1.946 to -0.668)	1.427*** (1.042 to 1.813)	2.616*** (2.049 to 3.184)
Woman currently working (yes)	-0.357*** (-0.590 to -0.123)	0.0162 (-0.221 to 0.253)	0.210 (-0.110 to 0.529)
Place of residents (urban)	0.0337 (-0.272 to 0.339)	0.107 (-0.115 to 0.330)	0.497*** (0.177 to 0.817)
Sex of household head (female)	0.230 (-0.0570 to 0.518)	0.0205 (-0.177 to 0.218)	-0.0346 (-0.336 to 0.267)
Parity	-0.164*** (-0.243 to -0.085)	-0.118*** (-0.190 to -0.046)	-0.084 (-0.199 to 0.030)
Household size	-0.035 (-0.089 to 0.019)	0.013 (-0.034 to 0.061)	0.034 (-0.044 to 0.111)
Frequency of watching TV			
Not at all			
Less than once a week	-0.539*** (-0.905 to -0.174)	0.076 (-0.295 to 0.447)	-0.131 (-0.700 to 0.439)
At least once a week	-0.066 (-0.355 to 0.223)	0.043 (-0.194 to 0.281)	0.035 (-0.357 to 0.426)
Currently breastfeeding (yes)	0.102 (-0.112 to 0.316)	-0.156* (-0.338 to 0.026)	-0.204 (-0.510 to 0.102)
Marital status			
Never in union (reference)			
Married or cohabiting	0.299 (-0.197 to 0.794)	0.870* (0.422 to 1.317)	0.898* (0.212 to 1.584)
Formerly in union	0.209 (-0.339 to 0.756)	0.488** (0.047 to 0.929)	0.312 (-0.392 to 1.017)
Decision on household large purchases			
Respondent alone (reference)			
Both respondent and partner	-0.217 (-0.546 to 0.113)	-0.303** (-0.543 to -0.063)	0.0495 (-0.259 to 0.358)
Partner alone	-0.298* (-0.641 to 0.045)	-0.0677 (-0.350 to 0.215)	-0.128 (-0.528 to 0.273)
Someone else/other	-2.309*** (-4.053 to -0.565)	-0.970 (-3.170 to 1.231)	-13.66*** (-14.580 to -12.730)
Decision on respondent's health			
Respondent alone (reference)			
Both respondent and partner	-0.0559 (-0.427 to 0.315)	0.0704 (-0.194 to 0.335)	-0.131 (-0.524 to 0.262)
Partner alone	-0.220 (-0.601 to 0.161)	-0.016 (-0.301 to 0.268)	-0.183 (-0.633 to 0.266)
Someone else/other	-2.446*** (-4.226 to -0.666)	0.092 (-2.677 to 2.862)	-12.920*** (-14.610 to -11.230)
Observations	9993	9993	9993

CI in parentheses.

*P<0.1; **P<0.05; ***P<0.01.

BMI, body mass index.

Nigeria. Thus, compared with normal weight women, an additional year of education was associated with a higher likelihood of overweight and obesity in Ghana, Mozambique and Nigeria. Conversely, number of years of education was associated negatively with underweight in Kenya and Nigeria relative to normal weight. Also, older age was significantly and positively associated with underweight, overweight and obesity compared with normal weight women in all countries included in the analysis. An exception could be made of Nigeria where older age was inversely related to underweight. The results in relation to household wealth index

were mixed. While generally, significant positive associations were observed between wealth index (middle, richer and richest quintiles) and overweight and obesity in all the five countries, inverse relationship was observed for underweight. Compared with normal weight women, higher maternal parity was inversely related to underweight in Ghana and Mozambique, and both underweight and overweight in Kenya. In DRC, maternal parity was inversely related to underweight and positively related to obesity. Breastfeeding was associated with less likelihood of underweight among women in DRC and Nigeria, obesity in DRC and Ghana, overweight in Kenya

Table 6 Multivariable analysis of the association between sociodemographic correlates and malnutrition, Mozambique

Variables	Underweight	Overweight	Obesity
	(BMI<18.50)	(BMI=25–29.99)	(BMI≥30)
Women' education (yes)	0.011 (–0.041 to 0.063)	0.044*** (0.0127 to 0.076)	0.098*** (0.041 to 0.156)
Women's age (in years)	0.018 (–0.013 to 0.049)	0.032*** (0.012 to 0.052)	0.075*** (0.042 to 0.109)
Household wealth index			
Poorest (reference)			
Poorer	–0.146 (–0.479 to 0.187)	0.618* (0.222 to 1.015)	1.061 (–0.529 to 2.650)
Middle	–0.741*** (–1.118 to –0.365)	0.856* (0.430 to 1.282)	2.424*** (1.023 to 3.825)
Rich	–0.618*** (–1.081 to –0.154)	1.235* (0.818 to 1.652)	2.905*** (1.576 to 4.234)
Richest	–0.800*** (–1.408 to –0.192)	1.977*** (1.517 to 2.437)	4.832*** (3.496 to 6.168)
Woman currently working (yes)	0.195 (–0.087 to 0.477)	–0.0795 (–0.271 to 0.112)	0.014 (–0.371 to 0.400)
Place of residence (urban)	0.0766 (–0.306 to 0.459)	–0.0722 (–0.350 to 0.205)	–0.0180 (–0.504 to 0.468)
Sex of household head (female)	–0.342** (–0.649 to –0.036)	–0.056 (–0.264 to 0.153)	–0.142 (–0.570 to 0.286)
Parity	–0.124** (–0.224 to –0.023)	0.0379 (–0.024 to 0.100)	0.160** (0.037 to 0.282)
Household size	0.0161 (–0.037 to 0.070)	–0.010 (–0.042 to 0.022)	–0.037 (–0.108 to 0.035)
Frequency of watching TV			
Not at all (reference)			
Less than once a week	–0.0113 (–0.328 to 0.306)	0.157 (–0.091 to 0.404)	0.630*** (0.214 to 1.047)
At least once a week	–0.214 (–0.518 to 0.0895)	–0.0124 (–0.213–0.188)	0.044 (–0.329 to 0.418)
Currently breastfeeding	0.215 (–0.049 to 0.478)	–0.298*** (–0.493 to –0.102)	–0.516** (–0.938 to –0.0944)
Marital status			
Never in union (reference)			
Married or cohabiting	0.253 (–0.419 to 0.925)	0.196 (–0.223 to 0.615)	0.708* (–0.0469 to 1.462)
Formerly in union	0.351 (–0.288 to 0.989)	–0.029 (–0.462 to 0.404)	0.773** (0.0197 to 1.527)
Decision on large household purchases			
Respondent alone (reference)			
Both respondent and partner	–0.096 (–0.472 to 0.281)	–0.0634 (–0.329 to 0.202)	–0.0414 (–0.647 to 0.564)
Partner alone	–0.049 (–0.420 to 0.323)	0.037 (–0.256 to 0.329)	–0.366 (–1.022 to 0.289)
Someone else/other	0.036 (–1.044 to 1.115)	0.746 (–0.600 to 2.091)	–0.983 (–3.268 to 1.303)
Decision on respondent's health			
Respondent alone (reference)			
Both respondent and partner	–0.184 (–0.633 to 0.265)	–0.134 (–0.425 to 0.158)	–0.143 (–0.777 to 0.491)
Partner alone	–0.317 (–0.786 to 0.151)	–0.148 (–0.440 to 0.144)	0.227 (–0.360 to 0.814)
Someone else/other	–0.480 (–1.477 to 0.516)	–0.646 (–1.649 to 0.357)	–1.232 (–3.319 to 0.855)
Observations	11 017	11 017	11 017

CI in parentheses.

*P<0.1; **P<0.05; ***P<0.01.

BMI, body mass index.

and overweight and obesity in Mozambique and Nigeria relative to normal weight women.

The figures 1 and 2 are pictorial presentations of the probability of women falling into underweight, overweight and obese categories if there were changes in their age and years of education.

DISCUSSION

This study investigated the correlates of DBM among women in five SSA countries. The key indicators of DBM used in the analysis were underweight, overweight and

obesity. The results showed expectedly that there are some variations across countries on how the correlates included in this study are associated with DBM. Our analysis reveals that in Ghana, Mozambique and Nigeria, a higher number of years of formal education is associated with the likelihood of overweight and obesity relative to normal weight women. Thus, a higher number of years of education is a risk factor for women with unhealthy weight. This could be due to lifestyles changes as one achieve more years of education, which may include sedentary lifestyles and poor dietary patterns.³² Further,

Table 7 Multivariable analysis of the association between sociodemographic correlates and malnutrition, Nigeria

Variable	Underweight	Overweight	Obesity
	(BMI<18.50)	(BMI=25–29.99)	(BMI≥30)
Women's education (in years)	-0.042*** (-0.067 to -0.018)	0.034*** (0.019 to 0.049)	0.048*** (0.028 to 0.068)
Women's age (in years)	-0.019** (-0.035 to -0.0021)	0.040*** (0.030 to 0.051)	0.072*** (0.056 to 0.088)
Household wealth index			
Poorest (reference)			
Poorer	-0.200** (-0.366 to -0.034)	0.289*** (0.090 to 0.488)	0.208 (-0.280 to 0.697)
Middle	-0.291** (-0.517 to -0.0650)	0.691*** (0.479 to 0.903)	0.788* (0.298 to 1.278)
Rich	-0.208 (-0.477 to 0.0614)	0.998* (0.764 to 1.232)	1.196* (0.694 to 1.698)
Richest	-0.530*** (-0.921 to -0.139)	1.351*** (1.074 to 1.628)	1.862*** (1.353 to 2.372)
Woman currently working (yes)	-0.0518 (-0.198 to 0.0944)	0.0231 (-0.102 to 0.148)	-0.292*** (-0.497 to -0.0878)
Place of residence (urban)	0.166 (-0.0500 to 0.381)	0.0775 (-0.0643 to 0.219)	0.178* (-0.0244 to 0.379)
Sex of household head (female)	-0.0836 (-0.336 to 0.169)	0.0127 (-0.149 to 0.174)	-0.084 (-0.300 to 0.133)
Parity	-0.011 (-0.0538 to 0.0313)	0.039** (0.009 to 0.0698)	0.0230 (-0.021 to 0.067)
Household size	0.007 (-0.015 to 0.029)	-0.009 (-0.025 to 0.00654)	0.023* (-0.002 to 0.047)
Frequency of watching TV			
Not at all (reference)			
Less than once a week	-0.146 (-0.336 to 0.045)	0.0771 (-0.056 to 0.210)	0.0585 (-0.162 to 0.278)
At least once a week	-0.252*** (-0.438 to -0.066)	0.155** (0.024 to 0.285)	0.205* (-0.009 to 0.420)
Currently breastfeeding	0.178*** (0.043 to 0.314)	-0.176*** (-0.278 to -0.073)	-0.363*** (-0.524 to -0.202)
Marital status			
Never in union			
Married or cohabiting	-0.177 (-0.652 to 0.299)	0.633* (0.188 to 1.078)	0.557* (-0.090 to 1.204)
Formerly in union	0.0721 (-0.400 to 0.545)	0.921* (0.426 to 1.416)	1.071* (0.417 to 1.724)
Decision on large household purchases			
Respondent alone			
Both respondent and partner	-0.206 (-0.641 to 0.230)	0.0218 (-0.209 to 0.252)	0.341* (-0.002 to 0.684)
Partner alone	-0.181 (-0.579 to 0.218)	-0.015 (-0.238 to 0.209)	-0.177 (-0.538 to 0.184)
Someone else/other	1.314* (-0.239 to 2.866)	-0.651 (-1.912 to 0.609)	-0.225 (-1.731 to 1.281)
Decision on respondent's health			
Respondent alone (reference)			
Both respondent and partner	-0.0847 (-0.525 to 0.356)	-0.0141 (-0.251 to 0.223)	-0.192 (-0.547 to 0.163)
Partner alone	0.069 (-0.334 to 0.474)	0.025 (-0.206 to 0.256)	-0.088 (-0.460 to 0.283)
Someone else/other	-2.381*** (-1.417 to -0.423)	-1.369** (-4.688 to -3.669)	-2.604** (-7.160 to -5.311)
Observations	31 170	31 170	31 170

CI in parentheses.

*P<0.1; **P<0.05; ***P<0.01.

BMI, body mass index.

it could be that people who are already overweight or obese have higher propensity of adding more weight relative to those who have normal weight. In contrast, there was an inverse relationship between the number of years of formal education and underweight in Kenya and Nigeria. These findings are in line with previous studies. In Ghana, higher education was associated with overweight and obesity among women.³³ Additionally, high education was associated positively with overweight among women in Indonesia.³⁴ Nevertheless, in the same study, high education was associated with reduced risk of

being underweight by 10%–30%.³⁴ This is contrary to our findings in Kenya and Nigeria.

Furthermore, age was associated positively with all the DBM indicators across the five countries. Thus, older women are more likely to be overweight and obese and less likely to be underweight. This suggests that older age is a protective factor for underweight, while a risk factor for overweight and obesity. This positive correlation between age, overweight and obesity may have a consequential effect on non-communicable diseases (NCDs) among older people, as an unhealthy weight is a major




Figure 1 Graphical Illustration of the results of the probability of falling into the double burden of malnutrition categories when years of education increase.

risk factor for NCDs.^{35–38} An exception to these findings could be made of Nigeria where older age was associated negatively with underweight women. This implies that in this setting, the older women become the more likelihood that they will suffer from underweight. The consequential effect of this may be poor health outcomes, as being underweight exposes women to higher risks of morbidity and mortality during pregnancy and child birth.^{39–41} The relationship between age and DBM has been documented in previous studies. For example, Doku and Neupane observed a significant positive association between age and the key indicators of DBM in Ghana.³³ A study in Bangladesh observed a significant positive relationship between older age and DBM.⁴² These findings together with the findings from our study confirmed the contribution of age to the DBM epidemic.

Our analysis showed that household wealth index had mixed effects on DBM. In general, the three rich quintiles: middle, richer and richest were associated significantly and positively with overweight and obesity among women across all the countries included in the analysis. This may be due to obesogenic effects of increased household wealth as dietary pattern changes,⁴³ and the fact that there is a documented positive correlation between household wealth and unhealthy body weight.^{44 45} In Bangladesh and Nepal, higher household wealth was associated with an increased likelihood of being overweight and obese.⁴⁴ Also, being rich was associated with

overweight and obesity among Ghanaian women.³³ Interestingly, higher household wealth had an inverse effect on underweight. This inverse relationship may be due to the fact that most underweight women are likely to be in the poorer wealth quintiles⁴⁶ and, therefore, may be unaffected by the higher household wealth quintiles. The inverse relationship has been observed previously.⁴⁷ Relatedly, compared with the richest, women from the poorest households were significantly most likely to be underweight and least likely to be overweight over normal weight in Bangladesh.¹⁷ Suggesting that being in the poorest household is protective against overweight but not underweight.⁴⁶

The health benefits of breastfeeding were illuminated in this study. Breastfeeding was found to associate with less likelihood of underweight, overweight and obesity in the five countries analysed. This suggests that mothers who have unhealthy weight should be encouraged to practice breastfeeding as the benefits are not only to their offspring but also for their own health and well-being. These findings confirm the widely recognised benefits of breastfeeding for improved health and developmental outcomes in mothers and their infants.^{48–51} The implication of this may be that interventions to promote breastfeeding may have positive impact on the DBM epidemic at the national level. The benefits of breastfeeding to women health have previously been documented. For example, breastfeeding has been suggested as an efficient means of








Figure 2 Graphical Illustration of results of the probability of falling into the double burden of malnutrition categories when age increases.

promoting postpartum weight loss due to its high energy cost.⁵² Further evidence suggest that at 12-weeks postpartum, exclusively breastfeeding (EBF) mothers had lost more total body weight than mixed feeding mothers. However, mixed feeding mothers lost slightly more per cent body fat than exclusively breastfeeding mothers.⁵² The authors concluded that EBF promotes greater weight loss than mixed feeding among mothers even in the early postpartum period. Several other studies have shown that EBF influences postpartum weight loss.⁵¹⁻⁵⁴ The preceding discussion points to the need for health policymakers to design programmes to encourage mothers to breastfeed, especially, practice exclusive breastfeeding as a means of DBM prevention.

An important strength of our study is the use of large nationally representative samples, thereby providing more robust estimates of observed associations as well as enhancing the generalisability of the findings. The use of multicountry data help unmask differences and commonalities in the effects of the correlates on DBM across countries, which would not have been possible with single country data. Additionally, the height and weight data which were used to compute the BMI were objectively measured, reducing possible misclassification. A limitation worth mentioning is the cross-sectional nature of the data, which does not lends itself to the establishments of causal relationship between the predictor and outcome variables. The conclusions in the paper are

therefore interpreted as mere associations between the predictor variables and the outcome variable. Another limitation is that due to data limitation, we were not able to examine DBM at the individual and within households. The analysis and interpretation in this paper are therefore limited to DBM at population or national level, whereby underweight, overweight and obesity coexist in the same country.

CONCLUSIONS

The study investigated the correlates of the DBM in five SSA countries. The analysis revealed that the effects of the correlates on DBM are largely similar across countries, except in few cases where there were disparities in the effects. The results indicate that higher number of years of education increases the likelihood of overweight and obesity among women in Ghana, Mozambique and Nigeria. Conversely, number of years of education is associated negatively with underweight in Kenya and Nigeria. Living in better-off households increases significantly the likelihood of overweight and obesity among women across all countries, while associated with the likelihood of underweight. Interventions to address DBM should take into account socioeconomic status. This may include providing special programmes for women who have unhealthy weight in wealthy households. A unit change in age is also associated significantly and positively with

underweight, overweight and obesity in all the countries included in the analysis. Breastfeeding is associated with least likelihood of underweight, overweight and obesity in breastfeeding mothers. This implies that interventions to prevent DBM should incorporate breastfeeding to enhance their effectiveness.

Acknowledgements We wish to express our profound gratitude to The DHS Program, USA, for providing us access to the data. We also wish to acknowledge institutions of respective countries that played critical roles in the data collection process.

Contributors DAA conceived and designed the study. DAA and ZTD conducted the data analysis, interpreted the results and drafted the manuscript. CK contributed to study design, data analysis, interpretation and critical revision of the manuscript. All authors take responsibility of any issues that might arise from the publication of this manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement This study was a re-analysis of existing data that are publicly available from The DHS Program at <http://dhsprogram.com/publications/publication-fr221-dhs-final-reports.cfm>. Data are accessible free of charge upon a registration with the Demographic and Health Survey program (The DHS Program). The registration is done on the DHS website indicated above.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- World Health Organization. Double burden of malnutrition. 2017 <http://www.who.int/nutrition/double-burden-malnutrition/en/>
- Shrimpton R, Rokx C. *The Double Burden of Malnutrition: A Review of Global Evidence. Health, Nutrition and Population Discussion Paper*. World Bank, Washington, DC: World Bank, 2012. <https://openknowledge.worldbank.org/handle/10986/27417>
- NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 2016;387:1377–96.
- Kolčić I. Double burden of malnutrition: A silent driver of double burden of disease in low- and middle-income countries. *J Glob Health* 2012;2:020303.
- World Health Organization. *Global Status Report on Non-communicable diseases 2010*. Geneva: WHO, 2011.
- Amugsi DA, Dimbuene ZT, Mberu B, et al. Prevalence and time trends in overweight and obesity among urban women: an analysis of demographic and health surveys data from 24 African countries, 1991–2014. *BMJ Open* 2017;7:e017344.
- Mendez MA, Monteiro CA, Popkin BM. Overweight exceeds underweight among women in most developing countries. *Am J Clin Nutr* 2005;81:714–21.
- Delisle HF. Poverty: the double burden of malnutrition in mothers and the intergenerational impact. *Ann N Y Acad Sci* 2008;1136:172–84.
- Guilbert JJ. The world health report 2002 - reducing risks, promoting healthy life. *Educ Health* 2003;16:230.
- World Health Organization. Global health risks: mortality and burden of disease attributable to selected major risks. 2009 http://www.who.int/healthinfo/global_burden_disease/GlobalHealthRisks_report_full.pdf
- Delisle H. Early nutritional influences on obesity, diabetes and cardiovascular disease risk. International Workshop, 6–9 June 2004, Montreal University, Quebec, Canada. *Maternal and Child Nutrition* 2005;1:128–9.
- World Health Organization. Urgent action needed to tackle the double burden of malnutrition and to achieve Universal Health Coverage and Sustainable Development Goals in Africa. 2018 <https://www.afro.who.int/news/urgent-action-needed-tackle-double-burden-malnutrition-and-achieve-universal-health-coverage>
- Kennedy G, Nantel G, Shetty P. Assessment of the double burden of malnutrition in six case study countries. <http://www.fao.org/docrep/009/a0442e/a0442e03.htm#TopOfPage>
- Hasan M, Sutradhar I, Shahabuddin A, et al. *Double Burden of Malnutrition among Bangladeshi Women: A Literature Review*. Muacevic A, Adler JR, eds. 9, 2017. Cureus.
- Sarma H, Saquib N, Hasan MM, et al. Determinants of overweight or obesity among ever-married adult women in Bangladesh. *BMC Obes* 2016;3:13.
- Zeba AN, Delisle HF, Renier G, et al. The double burden of malnutrition and cardiometabolic risk widens the gender and socio-economic health gap: a study among adults in Burkina Faso (West Africa). *Public Health Nutr* 2012;15:2210–9.
- Kamal SM, Hassan CH, Alam GM. Dual burden of underweight and overweight among women in Bangladesh: patterns, prevalence, and sociodemographic correlates. *J Health Popul Nutr* 2015;33:92–105.
- Kimani-Murage EW, Muthuri SK, Oti SO, et al. Evidence of a Double Burden of Malnutrition in Urban Poor Settings in Nairobi, Kenya. *PLoS One* 2015;10:e0129943.
- Neupane S, Prakash KC, Doku DT. Overweight and obesity among women: analysis of demographic and health survey data from 32 Sub-Saharan African Countries. *BMC Public Health* 2016;16:30.
- The DHS Program. Demographic and Health Surveys. <http://dhsprogram.com/data/available-datasets.cfm>
- Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF International. *Ghana Demographic and Health Survey 2014*. Rockville, Maryland, USA: GSS, GHS, and ICF International, 2015.
- Amugsi DA, Dimbuene ZT, Kimani-Murage EW, et al. Differential effects of dietary diversity and maternal characteristics on linear growth of children aged 6–59 months in sub-Saharan Africa: a multi-country analysis. *Public Health Nutr* 2017;20:1029–45.
- The DHS Program. Demographic and Health Surveys. https://dhsprogram.com/data/dataset_admin/download-datasets.cfm (cited 9 Jun 2018).
- The DHS Program. DHS Methodology. <http://dhsprogram.com/What-We-Do/Survey-Types/DHS-Methodology.cfm>
- Ghana Statistical Service (GSS), Ghana Health Service (GHS), ICF Macro. *Ghana Demographic and Health Survey 2008 Accra, Ghana: GSS, GHS, and ICF Macro*, 2009.
- Ministry of Health and Social Services (MoHSS), Macro International Inc. *Namibia Demographic and Health Survey 2006–07 Windhoek, Namibia and Calverton*. Maryland, USA: MoHSS and Macro International Inc, 2008.
- National Population Commission (NPC) [Nigeria], ICF International. *Nigeria Demographic and Health Survey*. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International, 2014.
- National Bureau of Statistics-Kenya, ICF International. *2014 KDHS Key Findings*. Rockville, Maryland, USA: KNBS and ICF International, 2015.
- Eknoyan G. Adolphe Quetelet (1796–1874)—the average man and indices of obesity. *Nephrol Dial Transplant* 2008;23:47–51.
- World Health Organization. BMI classification: WHO, 2006. Available from: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html
- Starkweather J, Moske KA. Multinomial Logistic Regression, 2011. Available from: https://it.unt.edu/sites/default/files/mlr_jds_aug2011.pdf
- Kautzky-Willer A, Dorner T, Jensby A, et al. Women show a closer association between educational level and hypertension or diabetes mellitus than males: a secondary analysis from the Austrian HIS. *BMC Public Health* 2012;12:392.
- Doku DT, Neupane S. Double burden of malnutrition: increasing overweight and obesity and stall underweight trends among Ghanaian women. *BMC Public Health* 2015;15:670.
- Hanandita W, Tampubolon G. The double burden of malnutrition in Indonesia: Social determinants and geographical variations. *SSM Popul Health* 2015;1:16–25.
- Kunutsor S, Powles J. Descriptive epidemiology of blood pressure in a rural adult population in Northern Ghana Rural Remote Health. 2009;9:1095.
- Agyemang C. Rural and urban differences in blood pressure and hypertension in Ghana, West Africa. *Public Health* 2006;120:525–33.
- Van Middendorp D, Asbroek A, Bio FY, et al. Rural and urban differences in blood pressure and pregnancy-induced hypertension among pregnant women in Ghana Globalization and Health. 2013;9:59.
- Nichols SR, Hypertension EP. In: Murray CJL, Lopez AD, eds. *Quantifying Global Health Risks: the Burden of Disease Attributable*

- to *Selected Risk Factors*. Cambridge, Massachusetts: Harvard University Press, 1996.
39. Negoro H, Sugino Y, Nishizawa K, *et al*. Underweight body mass index is a risk factor of mortality in outpatients with nocturia in Japan. *BMC Res Notes* 2015;8:490.
 40. Park D, Lee JH, Han S. Underweight: another risk factor for cardiovascular disease?: A cross-sectional 2013 Behavioral Risk Factor Surveillance System (BRFSS) study of 491,773 individuals in the USA. *Medicine* 2017;96:e8769.
 41. Victora CG, Adair L, Fall C, *et al*. Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* 2008;371:340–57.
 42. Oddo VM, Rah JH, Semba RD, *et al*. Predictors of maternal and child double burden of malnutrition in rural Indonesia and Bangladesh. *Am J Clin Nutr* 2012;95:951–8.
 43. Aitsi-Selmi A, Bell R, Shipley MJ, *et al*. Education modifies the association of wealth with obesity in women in middle-income but not low-income countries: an interaction study using seven national datasets, 2005–2010. *PLoS One* 2014;9:e90403.
 44. Bishwajit G. Household wealth status and overweight and obesity among adult women in Bangladesh and Nepal. *Obes Sci Pract* 2017;3:185–92.
 45. Balarajan Y, Villamor E. Nationally representative surveys show recent increases in the prevalence of overweight and obesity among women of reproductive age in Bangladesh, Nepal, and India. *J Nutr* 2009;139:2139–44.
 46. Tebekaw Y, Teller C, Colón-Ramos U. The burden of underweight and overweight among women in Addis Ababa, Ethiopia. *BMC Public Health* 2014;14:1126.
 47. Selvamani Y, Singh P. Socioeconomic patterns of underweight and its association with self-rated health, cognition and quality of life among older adults in India. *PLoS One* 2018;13:e0193979.
 48. Jonas W, Nissen E, Ransjö-Arvidson AB, *et al*. Short- and long-term decrease of blood pressure in women during breastfeeding. *Breastfeed Med*. 2008;3:103–9.
 49. Handlin L, Jonas W, Ransjö-Arvidson AB, *et al*. Influence of common birth interventions on maternal blood pressure patterns during breastfeeding 2 days after birth. *Breastfeed Med*. 2012;7:93–9.
 50. Stuebe AM, Schwarz EB, Grewen K, *et al*. Duration of lactation and incidence of maternal hypertension: a longitudinal cohort study. *Am J Epidemiol* 2011;174:1147–58.
 51. Sámano R, Martínez-Rojano H, Godínez Martínez E, *et al*. Effects of breastfeeding on weight loss and recovery of pregestational weight in adolescent and adult mothers. *Food Nutr Bull* 2013;34:123–30.
 52. Hatsu IE, McDougald DM, Anderson AK. Effect of infant feeding on maternal body composition. *Int Breastfeed J* 2008;3:18.
 53. López-Olmedo N, Hernández-Cordero S, Neufeld LM, *et al*. The associations of maternal weight change with breastfeeding, diet and physical activity during the postpartum period. *Matern Child Health J* 2016;20:270–80.
 54. Schalla SC, Witcomb GL, Haycraft E. Body shape and weight loss as motivators for breastfeeding initiation and continuation. *Int J Environ Res Public Health* 2017;14.