BMJ Open Awareness of risk factors and warning symptoms and attitude towards gastric cancer screening among the general public in China: a cross-sectional study

Qi Liu,¹ Xi Zeng,^{2,3} Wen Wang,¹ Ruo-lin Huang,¹ Yan-jin Huang,¹ Shan Liu,¹ Ying-hui Huang,¹ Ying-xin Wang,¹ Qing-hong Fang,¹ Guoping He,¹ Ying Zeng¹

To cite: Liu Q, Zeng X, Wang W, *et al.* Awareness of risk factors and warning symptoms and attitude towards gastric cancer screening among the general public in China: a cross-sectional study. *BMJ Open* 2019;9:e029638. doi:10.1136/bmjopen-2019-029638

QL and XZ contributed equally.

Received 03 February 2019
Revised 27 June 2019
Accepted 05 July 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Nursing, University of South China, Hengyang, China

²Hunan Province Key Laboratory of Tumor Cellular & Molecular Pathology, Cancer Research Institute, University of South China, Hengyang, China

³Department of Pathology, Hunan Province Cooperative Innovation Center for Molecular Target New Drug Study, University of South China, Hengyang, China

Correspondence to

Dr Ying Zeng;
zengying2003@126.com

ABSTRACT

Objectives This study aimed to assess the knowledge of risk factors and warning symptoms and attitude towards gastric cancer screening among the general population in China.

Setting Hunan province, China

Participants Individuals aged older than 18 years were recruited using a cluster sampling method.

Design A cross-sectional study, and a pretested structured questionnaire was used to assess participants' awareness of gastric cancer.

Primary and secondary outcome measures Knowledge level of risk factors and warning symptoms of gastric cancer, gastric cancer screening attitude, sociodemographic factors associated with gastric cancer knowledge and screening behaviour.

Results This study comprised 1200 participants with a mean age of 40.31 (SD 16.73) years, of whom 622 (51.8%) were women. The mean score for gastric cancer knowledge was 8.85/22 (SD 6.48). There were 47.0% of the participants who had a low knowledge level about the risk factors and warning symptoms of gastric cancer. In total, 83.8% believed screening is helpful for early detection of gastric cancer, and 15.2% had undergone gastric cancer screening. The most common reason for not undergoing screening was having 'no symptoms' (63.0%), followed by 'fear of undergoing gastroscopy' (38.1%). Independent factors related to lower knowledge levels included male sex, living in rural areas, lower educational level, working as a farmer and without a family history of gastric cancer ($p < 0.05$). Factors independently associated with screening behaviour included white-collar employment, higher income and having upper gastrointestinal tract diseases ($p < 0.05$).

Conclusions In China, people have poor knowledge about risk factors and warning symptoms of gastric cancer, but a majority have a positive attitude towards the benefits of gastric cancer screening. Being asymptomatic and having a fear of gastroscopy were the main self-reported reasons for not undergoing screening. These results highlight the urgent need for educational campaigns to improve gastric cancer awareness.

INTRODUCTION

Although the incidence of gastric cancer has been declining, it remains the fifth most

Strengths and limitations of this study

- ▶ We used quantitative methods to assess the level of gastric cancer knowledge among the general population.
- ▶ Our findings may be used as a reference for other countries with high incidence rates of gastric cancer and no screening programmes.
- ▶ The participants' screening history was self-reported, and thus, there may be biases in the accuracy of recalling information.
- ▶ Other factors possibly related to knowledge and screening, such as cancer risk awareness, belief or culture, were not explored in this study.

common cancer and the third leading cause of cancer-related deaths worldwide.¹ Particularly, the incidence and mortality rates of gastric cancer in China account for a staggering 42.6% and 45.0%, respectively, of the global rates, and the number of new cases continues to increase due to population growth and ageing.^{2,3}

Early gastric cancer has significantly better prognosis than advanced gastric cancer.^{4,5} However, the 5-year survival rate of gastric cancer is significantly lower in China than in Japan and Korea,⁶ suggesting delays in diagnosis.⁷ Factors contributing to this delay could be poor awareness of risk factors or symptoms, and a negative attitude towards gastric cancer screening. *Helicobacter pylori* (*H. pylori*) infection has been identified as a primary causative factor of gastric cancer, with almost 90% of non-cardia gastric cancer cases attributable to *H. pylori* infection.⁸ A previous study found that awareness of the association between *H. pylori* infection and gastric cancer is more likely to make a person undergo screening and receive treatment for *H. pylori* infection.⁹ This may primarily be because knowledge of the risk factors is a vital

aspect in developing cancer risk perceptions and further influencing the participation in cancer screening.^{10 11} Moreover, knowledge about risk factors is an important prerequisite to healthy behaviours.¹²

In addition, knowledge about warning symptoms is critical for early diagnosis of cancer. The presence of an abdominal lump, abdominal fullness and pain are warning symptoms of gastric cancer.¹³ A recent study showed that knowledge about warning symptoms can lead to earlier presentation to medical care, which could result in earlier diagnosis and better outcomes.¹⁴ Studies investigating public knowledge of risk factors or warning symptoms of gastric cancer in Korea and Iran reported a general lack of knowledge.^{15 16}

Gastric cancer screening via gastroscopy has been shown to reduce disease-related mortality as it detects precancerous polyps, and thus enables early diagnosis.^{17 18} An individual's attitude towards screening has been shown to be significantly related to screening behaviour.¹⁹ Despite the serious burden of gastric cancer in China, there are no nationwide screening guidelines or programmes.⁷ Due to its large population and the relative lack of medical and health resources, opportunistic screening with gastroscopy in asymptomatic people is the primary practice in China.^{20 21} However, opportunistic screening operates on a voluntary basis and depends only on requests from the individual or their physician.²² In clinical practice, at-risk individuals (eg, those with *H. pylori* infection and first-degree relatives with gastric cancer) are recommended for gastric cancer screening, for which they shoulder the entire cost.^{23–25} Compared with organised screening, opportunistic screening involves fewer formal decisions about whether to screen, whom to screen and at what intervals screening should be done.²²

In China, there is little information about public awareness of risk factors or warning symptoms and screening for gastric cancer. Therefore, it is important to assess gastric cancer awareness in the Chinese population as a basis for future practice. The primary aim of this study was to evaluate the knowledge about risk factors and warning symptoms, and the attitude towards gastric cancer screening among the general population in China. The secondary aim was to investigate sociodemographic factors associated with gastric cancer knowledge and screening behaviour.

METHODS

Setting and sample

This cross-sectional, descriptive study was conducted between March and July 2018. The minimum calculated sample size was 896, which was determined using the formula $N = [\mu\alpha^2 \times \pi \times (1 - \pi)] / \delta^2$,²⁶ in which the prevalence rate was 30% (π) based on the knowledge level of the pilot group, the significance level was 0.05 (α) and the allowable error was 0.03 (δ). Considering a non-response rate of 40%, a target sample of 1250 was finally determined.

Of the 32 medical centres in Hunan province, 11 were included in this study using a cluster random sampling method. People who visited the medical centres were invited to this study. With the support of medical centres, each trained interviewer was accompanied by a medical staff (physician or nurse) to introduce the study purpose. The inclusion criteria were (a) age ≥ 18 years, (b) ability to communicate verbally and (c) no history of cancer. The exclusion criteria were (a) age < 18 years, (b) language barrier and (c) history of cancer. Individuals who agreed to participate in this study completed a standardised face-to-face questionnaire about gastric cancer knowledge and screening attitude. Administering the questionnaire took approximately 10–20 min. To maximise the response rate, each participant received a gift worth ¥25 (about US\$3.5).

Study instrument

The questionnaire consisted of items about risk factors, warning symptoms and screening for gastric cancer. The risk factors and warning symptoms listed were those identified by the American Cancer Society and China Anti-Cancer Association. The questionnaire comprised four sections: (1) Sociodemographic characteristics, including gender, age, residence, marital status, educational level, occupation, monthly income, family history of gastric cancer, family history of cancer, personal health status and history of gastrointestinal disease. (2) Identification of risk factors of gastric cancer: participants were asked to judge whether the following 16 items were risk factors of gastric cancer: male sex, older age, *H. pylori* infection, stomach ulcer, previous stomach surgery, atrophic gastritis, family history of gastric cancer, salty diet, consumption of pickled foods, consumption of smoked foods, irregular diet, frequently eating of leftovers, frequent midnight snacking, smoking, drinking and stress. Responses included 'Yes', 'No' or 'Not sure/Do not know', with 'Yes' indicating that they identified the items to be risk factors. (3) Identification of warning symptoms of gastric cancer: participants were asked to judge whether the following six items were warning symptoms of gastric cancer: gastrointestinal bleeding, recurrent nausea and vomiting, weight loss, upper abdominal fullness, abdominal lump and abdominal pain. The response categories were similar to those in section (2). (4) Perceptions on the screening for gastric cancer: (i) 'Do you think gastric cancer can be prevented?' Options included 'Yes' and 'No'; (ii) 'Do you think early gastric cancer can be cured?' (Yes or No); (iii) 'Do you think screening can help to detect early gastric cancer?' (Yes or No); (iv) 'Have you ever undergone gastric cancer screening?' (Yes or No); (v) 'Why do you not undergo a screening for gastric cancer?' Options included 'Do not know the benefits of screening', 'Fear of undergoing gastroscopy', 'Worried about screening results', 'No symptoms', 'No time' and 'Financial limitations' (multiple answers were allowed) and (vi) 'Which screening method for gastric cancer would you prefer?' Options included 'Gastroscopy', 'Blood test' and 'None of them'.

For the second and third sections, each correct answer was scored one point, with the total scores ranging from 0 to 22 points. The respondents' knowledge was then categorised according to the total scores: high knowledge, 15–22 points; moderate knowledge, 8–14 points; and low knowledge, 0–7 points.²⁷

The reliability of the questionnaire was evaluated by pretesting it in 100 adults. The retest coefficients of the second, third and fourth sections were 0.852, 0.874 and 0.791, respectively, while the Cronbach's α coefficient were 0.879, 0.890 and 0.783, respectively.

Data analysis

All data were analysed using SPSS V.23.0. Sociodemographic characteristics and responses to each question were described using frequencies and percentages. Univariate logistic regression was used to determine the relationship between sociodemographic characteristics and knowledge and participation in gastric cancer screening. Variables with a p value ≤ 0.15 in the univariate analysis were entered into the multivariate logistic regression analysis to investigate the independent factors influencing knowledge and screening behaviour. For each gastric cancer knowledge model, response options for the dependent variable were categorised as 'low/moderate knowledge' and 'high knowledge'. Only the results of the multivariate analysis were presented using OR and 95% CI. A two-tailed p value of <0.05 was considered statistically significant.

Patient and public involvement

None of the patients was involved in the design or development of the research question and outcome measures. They were also not involved in the recruitment and conduct of the study. The participants were informed of the study results via text message after completion of the study.

RESULTS

Sociodemographic characteristics of the participants

Of the 1250 individuals given the questionnaire, 1230 responded, of whom 30 completed less than 50% of the questionnaire. Thus, the final response rate was 96% ($n=1200$). The mean age of the participants was 40.31 (SD 16.73) years. Of the 1200 participants, 622 (51.8%) were women, 766 (63.8%) lived in rural areas, 408 (34.0%) had high educational level (college or above) and 169 (14.1%) had high income ($\geq$ ¥5000). In total, 198 (16.5%) participants had a family history of cancer, 58 (4.8%) had a family history of gastric cancer and 248 (20.7%) had diseases of the upper gastrointestinal tract (table 1).

Knowledge about risk factors and warning symptoms of gastric cancer

Table 2 presents the participants' knowledge about gastric cancer. The mean knowledge score was 8.85 (SD 6.48). Of the 1200 participants, 564 (47.0%), 347

Table 1 Sociodemographic characteristics of participants ($n=1200$)

Characteristic	Number (%)
Gender	
Male	578 (48.2)
Female	622 (51.8)
Age (years)	
18–29	425 (35.4)
30–39	208 (17.3)
40–49	192 (16.0)
50–59	209 (17.4)
>60	166 (13.8)
Mean $\pm$ SD	40.31 $\pm$ 16.73
Marital status	
Single	467 (38.9)
Married	733 (61.1)
Residence	
City	434 (36.2)
Rural	766 (63.8)
Educational Level	
Uneducated or elementary school	294 (24.5)
High school	498 (41.5)
College or above	408 (34.0)
Occupation	
Farmer	399 (33.3)
Worker	283 (23.6)
White collar	243 (20.3)
Student	275 (22.9)
Income	
<¥2000 (<US\$300)	575 (47.9)
¥2000–5000 (US\$300–755)	456 (38.0)
>¥5000 (>US\$755)	169 (14.1)
Family history of gastric cancer	
Yes	58 (4.8)
No	1142 (95.2)
Family history of cancer	
Yes	198 (16.5)
No	1002 (83.5)
Health status	
Unhealthy	130 (10.8)
Suboptimal	672 (56.0)
Healthy	398 (33.2)
Diseases of the upper gastrointestinal tract	
Yes	248 (20.7)
No	952 (79.3)

(28.9%) and 289 (24.1%) had low, moderate and high knowledge of gastric cancer, respectively. The most highly recognised risk factor was irregular diet (59.8%),

Table 2 Participants' knowledge about risk factors and warning symptoms of gastric cancer (n=1200)

Category	Number	%
Risk factors of gastric cancer		
Irregular diet	717	59.8
Consumption of pickled foods	690	57.5
Drinking	681	56.8
Consumption of smoked foods	671	55.8
Salty diet	611	50.9
Stomach ulcer	587	48.9
Stress	538	44.8
Smoking	534	44.5
Frequent eating of leftovers	534	44.5
Atrophic gastritis	431	35.9
Older age	419	34.9
<i>H. pylori</i> infection	418	34.8
Family history of gastric cancer	415	34.6
Male sex	304	25.3
Previous stomach surgery	285	23.8
Frequent midnight snacking	225	18.8
Warning symptoms of gastric cancer		
Gastrointestinal bleeding	598	49.8
Abdominal lump	541	45.1
Abdominal pain	524	43.7
Upper abdominal fullness	501	41.8
Weight loss	452	37.7
Recurrent nausea and vomiting	413	34.4
Total knowledge level (22 points)		
Low (0–7)	564	47.0
Moderate (8–14)	347	28.9
High (15–22)	289	24.1
Mean score of gastric cancer risk factors and warning symptoms (22 points)	8.85 (SD 6.48)	

H. pylori, *Helicobacter pylori*.

followed by consumption of pickled foods (57.5%), drinking (56.8%), consumption of smoked foods (55.8%) and salty diet (50.9%). Less well-recognised risk factors included male sex (25.3%), previous stomach surgery (23.8%) and frequent midnight snacking (18.8%). The most highly recognised warning symptom was gastrointestinal bleeding (49.8%), followed by abdominal lump (45.1%) and abdominal pain (43.7%), while less well-recognised warning symptoms included weight loss (37.7%) and recurrent nausea and vomiting (34.4%).

Table 3 presents the outcomes of the multivariate analysis for factors associated with gastric cancer knowledge. In the univariate analysis, gender, age, residence, marital

status, educational level, occupation and family history of gastric cancer were significantly associated with gastric cancer knowledge ($p < 0.05$). These factors plus variables with p value < 0.15 in the univariate analysis were entered into the multivariate logistic regression model. The independent variables associated with knowledge included gender, age, residence, educational level, occupation and family history of gastric cancer ($p < 0.05$; table 3).

Participants who were found to be less knowledgeable about risk factors and warning symptoms of gastric cancer included male sex (OR 0.59; 95% CI 0.46 to 0.75), living in rural areas (OR 0.76; 95% CI 0.59 to 0.99), with lower educational level (uneducated and elementary school: OR 0.25; 95% CI 0.16 to 0.41), working as a farmer (OR 0.38; 95% CI 0.22 to 0.66) and those with a family history of gastric cancer (OR 0.20; 95% CI 0.12 to 0.34). Participants aged 18–29 years had better knowledge about gastric cancer than the other age groups (OR 3.50; 95% CI 2.10 to 5.81).

Attitudes towards gastric cancer screening

Table 4 presents the participants' attitudes towards gastric cancer screening. In total, 84.7% of participants thought gastric cancer could be prevented, 83.8% thought gastric cancer could be detected early and only 15.2% thought early gastric cancer could not be cured. Among the 1200 participants, 182 (15.2%) had undergone gastric cancer screening, including 97 participants aged older than 40 years, which is the recommended age range for gastric cancer screening. For the 1018 participants who did not undergo gastric cancer screening, the most common reason was 'no symptoms', followed by 'fear of undergoing gastroscopy' and being 'worried about screening results'. The participants were asked about the most acceptable screening method for gastric cancer and 60.2% preferred blood testing as their first choice, while only 29.8% chose endoscopy as their first option.

Table 5 presents a multivariate analysis of the factors associated with gastric cancer screening behaviour. In the univariate analysis, factors significantly associated with gastric cancer screening behaviour were age, marital status, occupation, income and diseases of the upper gastrointestinal tract ($p < 0.05$). These factors plus variables with p value < 0.15 in the univariate analysis were entered into the multivariate logistic regression model. Only three of these factors were independently associated with gastric cancer screening behaviour: occupation, income and diseases of the upper gastrointestinal tract ($p < 0.05$). White collar employees were more likely to have undergone gastric cancer screening compared with those in other occupations (OR 4.88; 95% CI 2.04 to 11.72). Participants with the lowest income were less likely to have undergone gastric cancer screening than those with higher incomes (OR 0.22; 95% CI 0.12 to 0.38). Further, participants without upper gastrointestinal diseases were less likely to have undergone gastric cancer screening than those with upper gastrointestinal diseases (OR 0.23; 95% CI 0.16 to 0.34).

Table 3 Multivariate analysis of factors associated with gastric cancer knowledge (n=1200)

Variable	Low/moderate, n (%)	High, n (%)	OR	95% CI
Gender				
Male	492 (85.1)	86 (14.9)	0.59	0.46 to 0.75
Female	419 (67.3)	203 (32.6)	1 (ref)	
Age (years)				
18–29	252 (59.3)	173 (40.7)	3.50	2.10 to 5.81
30–39	144 (69.2)	64 (30.8)	3.34	2.06 to 5.40
40–49	163 (84.9)	29 (15.1)	2.26	1.40 to 3.63
50–59	198 (94.8)	11 (5.2)	1.18	0.73 to 1.93
≥60	154 (92.8)	12 (7.2)	1 (ref)	
Residence				
Rural	611 (79.8)	155 (20.2)	0.76	0.59 to 0.99
City	300 (69.1)	134 (30.9)	1 (ref)	
Educational level				
Uneducated or elementary school	270 (91.8)	24 (8.2)	0.25	0.16 to 0.41
High school	426 (85.5)	72 (14.5)	0.43	0.30 to 0.61
College or above	215 (52.7)	193 (47.3)	1 (ref)	
Occupation				
Farmer	367 (92.0)	32 (8.0)	0.38	0.22 to 0.66
Worker	241 (85.2)	42 (14.8)	0.49	0.30 to 0.82
White collar	166 (68.4)	77 (31.6)	0.61	0.40 to 0.94
Student	137 (49.8)	138 (50.2)	1 (ref)	
Family history of gastric cancer				
No	885 (77.5)	257 (22.5)	0.20	0.12 to 0.34
Yes	26 (44.8)	32 (55.2)	1 (ref)	

Bold figures indicate the statistically significant findings ($p < 0.05$).
CI, confidence interval; OR, odds ratio; ref, reference.

DISCUSSION

Measuring and understanding public awareness about risk factors, warning symptoms and screening attitude could help in developing appropriate policies for gastric cancer prevention. In general, this study found that a large number of the population had poor knowledge about risk factors and warning symptoms of gastric cancer. However, the majority of participants had a positive attitude towards screening and its benefits. The most common reasons for not seeking screening included no symptoms and fear of undergoing gastroscopy.

Gastric cancer knowledge

In this study, the knowledge level was relatively lower than those reported in the Korean and Iranian populations,^{15 16} signifying poor awareness among the Chinese population regarding gastric cancer. Moreover, while participants had relatively good knowledge of lifestyle risk factors, they had a poorer knowledge of disease risk factors. For instance, more than 50% of participants identified irregular diet, drinking and consumption of pickled foods as

risk factors of gastric cancer, while less than 36% of participants identified atrophic gastritis, *H. pylori* infection and previous stomach surgery as risk factors. This result indicates that people are unfamiliar with these medical terms, which may imply a gap in educational interventions.

More than 50% of Chinese experience *H. pylori* infection.²⁸ In general, 24.2% of China's cancer burden is attributed to infection, of which 42% is attributed to *H. pylori* infection.²⁹ However, only 34.8% of participants in this study identified *H. pylori* infection as a risk factor of gastric cancer, and this was significantly lower compared with the rate among Koreans (58.3%).¹⁶ One of the reasons for this difference could be that Korea has national guidelines for gastric cancer screening. Further, healthcare professionals in Korea are more aware of the significance of gastric cancer knowledge, and thus provide better educational intervention.^{30 31} Another reason could be the high media exposure of *H. pylori* infection: due to the high prevalence of *H. pylori* infection, several advertisements about *H. pylori* infection are broadcast

Table 4 Gastric cancer screening perceptions among participants (n=1200)

Question	Number	%
Gastric cancer can be prevented		
Yes	1016	84.7
No	184	15.3
Early gastric cancer can be cured		
Yes	1018	84.8
No	182	15.2
Gastric cancer can be detected early by screening		
Yes	1006	83.8
No	194	16.2
Have you ever participated in gastric cancer screening?		
Yes	182	15.2
No	1018	84.8
Why do you not undergo gastric cancer screening? (multiple selections possible) (n=1018)*		
No symptoms	641	63.0
Fear of undergoing gastroscopy	388	38.1
Worried about screening results	196	19.3
Do not know the benefits of screening	143	14.0
No time	109	10.7
Financial limitations	92	9.0
Which screening test for GC would you prefer? (n=1200)		
Blood test	722	60.2
Gastroscopy	358	29.8
None of them	120	10.0

*Participants who did not undergo gastric cancer screening. GC, gastric cancer.

in Korea.³¹ In our study, more than 50% of participants did not identify any warning symptoms of gastric cancer, and this rate was higher than that among Iranians.¹⁵ This result shows that poor awareness of warning symptoms is a factor contributing to the high rate of advanced-stage diagnosis of gastric cancer in Chinese patients, and indicates that these warning symptoms could be a good target for awareness-raising initiatives.⁷

Knowledge of gastric cancer differed according to sociodemographic characteristics. Similar to a previous study,³² we found that women had better gastric cancer knowledge than men. This finding may be because women tend to be family caregivers, and thus typically have more interactions with the healthcare system, and get more opportunities to learn about cancer.^{33 34} Further, we found that middle-aged and older participants were

less knowledgeable about gastric cancer than younger individuals, which can be worrying because gastric cancer often occurs in people aged older than 40 years.³⁵ This suggests that middle-aged and older adults should be the key population for gastric cancer education. Participants from rural areas, who were uneducated or had primary education only, and who worked as farmers, showed lower knowledge about gastric cancer. Studies regarding lung or colorectal cancer awareness have reported similar findings, in which the lower socioeconomic groups were less knowledgeable.^{36 37} Evidence also suggests that the lower socioeconomic groups were more likely to have later-stage cancer at diagnosis.^{38 39} McCutchan *et al* reported that fearful and fatalistic beliefs about cancer and emotional barriers were also the reasons for delay in seeking medical help in cancer patients with lower socioeconomic status.⁴⁰ Thus, when educating this group about gastric cancer, attention should also be paid on their negative beliefs about cancer.

Gastric cancer screening attitude

Participants held positive attitudes towards screening and early diagnosis of gastric cancer, implying that they were aware of the value of early diagnosis and that the screening helps gastric cancer prevention. As expected, a small percentage of participants (15.2%) had undergone gastric cancer screening at least once. In China, gastroscopy is conducted via opportunistic screening and therefore, it relies on individuals voluntarily requesting it and burdening the cost.^{20 23} The response 'No symptoms' was the most common self-reported reason for not undergoing gastric cancer screening, which is concordant with findings from a previous study,¹⁶ implying that participants mistakenly believed screening was only required when symptoms appeared. Therefore, it is necessary to educate the public regarding the indications of cancer screening.

Moreover, 'fearing of undergoing gastroscopy' was also a major barrier for participation in screening, and the expressed willingness to have a blood test was markedly higher than undergoing gastroscopy. This finding reveals that the availability of reliable blood tests for gastric cancer could be a key driver of increased screening rates. Although gastroscopy remains the gold standard for detecting gastric cancer, it has the disadvantages of being painful and costly.²¹ Serologic testing is a more convenient and cost-effective method; however, although it has been used to identify the high-risk gastric cancer population, its effectiveness needs to be further clarified.⁴¹ The development of precision medicine may lead to novel alternative methods for early detection of gastric cancer.⁴²

'Financial limitations' was not found to be a reason for not undergoing gastric cancer screening; however, multivariate analysis showed that the high-income group or those with white-collar jobs were more likely to undergo screening than the low-income group. This may be partly because gastroscopy for gastric cancer screening is not covered by Chinese health insurance. It is also possible

Table 5 Multivariate analysis of factors associated with gastric cancer screening behaviour (n=1200)

Variable	Screened, n (%)	Not screened, n (%)	OR	95% CI
Occupation				
Farmer	59 (14.8)	340 (85.2)	3.66	1.63 to 8.19
Worker	53 (18.7)	230 (81.3)	3.35	1.42 to 7.91
White collar	62 (25.5)	181 (74.5)	4.88	2.04 to 11.72
Student	8 (2.9)	267 (97.1)	1 (ref)	
Income				
<¥2000 (<US\$300)	35 (6.1)	540 (93.9)	0.22	0.12 to 0.38
¥2000–5000 (US\$300–755)	94 (20.6)	363 (79.4)	0.58	0.38 to 0.89
>¥5000 (>US\$755)	53 (31.4)	116 (68.6)	1 (ref)	
Diseases of the upper gastrointestinal tract				
No	104 (10.9)	848 (89.1)	0.23	0.16 to 0.34
Yes	78 (31.5)	170 (68.5)	1 (ref)	

Bold figures indicate the statistically significant findings ($p < 0.05$).
CI, confidence interval; OR, odds ratio; ref, reference.

that high-income people have better health literacy, so they give more importance to cancer screening.⁴³

Taking one step forward

In Japan, a country with high gastric cancer incidence, cancer education and awareness is one of the specific initiatives of the Basic Plan to Promote Cancer Control Programs under the Cancer Control Act. This project was aimed to raise adolescent cancer awareness through cancer education courses among middle school students.⁴⁴ In China, the government has started to pay attention to public awareness about cancer. The Three Years Action Plan for Cancer Prevention and Control in China set a goal of attaining a 60% awareness rate of cancer prevention knowledge. In 2018, the China Digestive Union set 15–21 October as the gastric cancer prevention and awareness week, with the aim to popularise knowledge on gastric cancer prevention and improve awareness on screening and its benefits. However, data to evaluate the effectiveness of such interventions are lacking, and thus, this study could be used as a basis to measure the effectiveness of future public health campaigns.

Our study has shown that Chinese have low awareness regarding the risk factors and warning symptoms of gastric cancer. Further, they have some misconceptions about gastric cancer screening. There are several suggestions to improve public awareness of gastric cancer. First, the government could learn from initiatives in different countries, who used the media, such as TV, to broadcast knowledge on gastric cancer prevention.³¹ Moreover, the government should implement an organised gastric cancer screening programme and incorporate screening into insurance plans to improve people's screening behaviour.⁷ Second, health education platforms, such as through WeChat, should be set up at the hospital and community level, and this platform should be promoted

among the people. Third, health workers, especially community and oncology nurses, should be trained in raising public awareness about gastric cancer prevention. Less-known risk factors and warning symptoms found in this study need to be emphasised during education interventions. Further, more attention should be given to the lower-income population, who are at high risk for gastric cancer.⁴⁵ In this study, people with lower socioeconomic status not only had poorer knowledge, but also were less likely to be screened. Therefore, improving gastric cancer knowledge and financial subsidies for screening in this population is crucial to reduce the gastric cancer burden in China. In addition, men and the elderly should be targeted during education interventions.

Strengths and limitations

The main strength of this study is the high response rate, which could decrease the non-response bias.⁴⁶ This high response rate may be attributed to the medical centres' support, experienced interviewers, shorter survey length and a gift incentive. Further, face-to-face interview, which has the highest response rates compared with other survey methods,⁴⁶ was used in this study. However, this method inevitably cost a lot of manpower and material resources. Our study had several limitations. First, the participants' screening history was self-reported, and thus, the possibility of biases in the accuracy of recall information could not be eliminated. However, a previous study reported no significant difference in accuracy between self-reported and recorded screening history.⁴⁷ Second, sociocultural factors possibly related to knowledge and screening behaviour, such as beliefs and tradition, were not explored, because only a quantitative method was used. Moreover, the awareness of the gastric cancer risk, which is an important factor for cancer screening,⁴⁸ was not investigated. Qualitative or mixed-method studies

may be considered for a more comprehensive analysis of related factors. Third, the screening attitude was assessed under the Chinese gastric cancer screening system, which limits the generalisability of the results to other countries. However, the results of this study have valuable implications for improving gastric cancer screening systems in China and other developing countries.

CONCLUSIONS

This study indicates that people lack knowledge about the risk factors and warning symptoms of gastric cancer in China. However, the majority had a positive attitude towards the benefits of gastric cancer screening. Being asymptomatic and having fear of gastroscopy were the main self-reported reasons for not undergoing screening. Interventions should be developed to improve knowledge on gastric cancer and screening participation in China.

Acknowledgements The authors gratefully acknowledge the participants who participated in this study. We would like to thank Professor Xin Chen for providing statistical consultation. In particular, the advice of Professor Jyu-Lin Chen (University of California, San Francisco) on manuscript writing is sincerely appreciated.

Contributors Conceptualisation: YZ, QL and GH. Data curation: YZ, QL, WW and R-IH. Investigation: QL, R-IH, Y-JH, SL, Y-HH, Y-XW and Q-HF. Methodology: YZ and XZ. Software: WW. Writing—original draft: YZ and QL. Writing—review and editing: YZ and XZ.

Funding The research was funded by the National Natural Science Foundation of China (NSFC), Grant/award number: 81641112; the Hunan Provincial Natural Science Foundation of China, Grant/award number: 2019JJ50521 and 2019JJ40254; and the Key Project of Hunan Provincial Education Department, Grant/award number: 18A229; and supported by the University of South China Innovation Foundation for Postgraduate, Grant/award: 2018KYY300, and the University of South China Innovation & Entrepreneurship Foundation for Undergraduate, Grant/award: X2019208. This work was also supported by the Construction Program of the Key Discipline in Hunan Province, China (Basic Medicine Sciences in University of South China), Centre for Gastric Cancer Research of Hunan Province (University of South China) and Key Laboratory of Tumour Cellular & Molecular Pathology (University of South China), College of Hunan Province.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethic Review Committee of University of South China (Approval No.: 4304079009640).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
- Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359–E386.
- Chen W, Zheng R, Baade PD, et al. Cancer statistics in China, 2015. *CA Cancer J Clin* 2016;66:115–32.
- Coit DG, Andtbacka R, Anker CJ, et al. Melanoma, version 2.2013: featured updates to the NCCN guidelines. *J Natl Compr Canc Netw* 2013;11:395–407.
- Isobe Y, Nashimoto A, Akazawa K, et al. Gastric cancer treatment in Japan: 2008 annual report of the JGCA nationwide registry. *Gastric Cancer* 2011;14:301–16.
- Allemani C, Weir HK, Carreira H, et al. Global surveillance of cancer survival 1995–2009: analysis of individual data for 25 676 887 patients from 279 population-based registries in 67 countries (CONCORD-2). *The Lancet* 2015;385:977–1010.
- Zong L, Abe M, Seto Y, et al. The challenge of screening for early gastric cancer in China. *Lancet* 2016;388.
- Plummer M, Franceschi S, Vignat J, et al. Global burden of gastric cancer attributable to *Helicobacter pylori*. *Int J Cancer* 2015;136:487–90.
- Shin DW, Cho J, Kim SH, et al. Preferences for the "screen and treat" strategy of *Helicobacter pylori* to prevent gastric cancer in healthy Korean populations. *Helicobacter* 2013;18:262–9.
- Hay JL, Orom H, Kiviniemi MT, et al. "I don't know" my cancer risk: exploring deficits in cancer knowledge and information-seeking skills to explain an often-overlooked participant response. *Med Decis Making* 2015;35:436–45.
- McCaffery K, Wardle J, Waller J. Knowledge, attitudes, and behavioral intentions in relation to the early detection of colorectal cancer in the United Kingdom. *Prev Med* 2003;36:525–35.
- World Health Organization. Guide to cancer early diagnosis. Available: https://www.who.int/cancer/publications/cancer_early_diagnosis/en/ [Accessed 9 Oct 2018].
- Bai Y, Li Z-S, Zou D-W, et al. Alarm features and age for predicting upper gastrointestinal malignancy in Chinese patients with dyspepsia with high background prevalence of *Helicobacter pylori* infection and upper gastrointestinal malignancy: an endoscopic database review of 102,665 patients from 1996 to 2006. *Gut* 2010;59:722–8.
- Quaife SL, Forbes LJL, Ramirez AJ, et al. Recognition of cancer warning signs and anticipated delay in help-seeking in a population sample of adults in the UK. *Br J Cancer* 2014;110:12–18.
- Mansour-Ghanaei F, Joukar F, Soati F, et al. Knowledge about gastric carcinoma in North of Iran, a high prevalent region for gastric carcinoma: a population-based telephone survey. *Asian Pac J Cancer Prev* 2012;13:3361–6.
- Oh D-Y, Choi KS, Shin H-R, et al. Public awareness of gastric cancer risk factors and disease screening in a high risk region: a population-based study. *Cancer Res Treat* 2009;41:59–66.
- Kim H, Hwang Y, Sung H, et al. Effectiveness of gastric cancer screening on gastric cancer incidence and mortality in a community-based prospective cohort. *Cancer Res Treat* 2018;50:582–9.
- Gotoda T, Ishikawa H, Ohnishi H, et al. Randomized controlled trial comparing gastric cancer screening by gastrointestinal X-ray with serology for *Helicobacter pylori* and pepsinogens followed by gastrointestinal endoscopy. *Gastric Cancer* 2015;18:605–11.
- Moutel G, Duchange N, Lièvre A, et al. Low participation in organized colorectal cancer screening in France: underlying ethical issues. *Eur J Cancer Prev* 2019;28:27–32.
- Shen L, Zhou C, Liu L, et al. Application of oral contrast trans-abdominal ultrasonography for initial screening of gastric cancer in rural areas of China. *Dig Liver Dis* 2017;49:918–23.
- Nie Y, Wu K, Yu J, et al. A global burden of gastric cancer: the major impact of China. *Expert Rev Gastroenterol Hepatol* 2017;11:651–61.
- Kim BJ, Heo C, Kim BK, et al. Effectiveness of gastric cancer screening programs in South Korea: organized vs opportunistic models. *World J Gastroenterol* 2013;19:736–41.
- Di L, Wu H, Zhu R, et al. Multi-Disciplinary team for early gastric cancer diagnosis improves the detection rate of early gastric cancer. *BMC Gastroenterol* 2017;17:147.
- Liu WZ. Screening and early detection of gastric cancer. *Chin J Gastroenterol* 2018;23:129–32.
- Leung WK, Wu M-shiang, Kakugawa Y, et al. Screening for gastric cancer in Asia: current evidence and practice. *Lancet Oncol* 2008;9:279–87.
- Hajian-Tilaki K. Sample size estimation in epidemiologic studies. *Caspian J Intern Med* 2011;2:289–98.
- Taha H, Al Jaghbeer M, Al-Sabbagh MQ, et al. Knowledge and practices of colorectal cancer early detection examinations in Jordan: a cross sectional study. *Asian Pac J Cancer Prev* 2019;20:831–8.
- Nagy P, Johansson S, Molloy-Bland M. Systematic review of time trends in the prevalence of *Helicobacter pylori* infection in China and the USA. *Gut Pathog* 2016;8:8.
- Plummer M, de Martel C, Vignat J, et al. Global burden of cancers attributable to infections in 2012: a synthetic analysis. *Lancet Glob Health* 2016;4:e609–16.

30. Lee JH, Kim JG, Jung HK, *et al.* [Synopsis on clinical practice guideline of gastric cancer in Korea: an evidence-based approach]. *Korean J Gastroenterol* 2014;63:66–81.
31. Sin M-K, Kim I-H. Facilitators of and barriers to gastric cancer screening among Korean Americans. *Cancer Nurs* 2017;40:E59–E65.
32. Crane M, Scott N, O'Hara BJ, *et al.* Knowledge of the signs and symptoms and risk factors of lung cancer in Australia: mixed methods study. *BMC Public Health* 2016;16:508.
33. Arber S, Ginn J. Gender differences in informal caring. *Health Soc Care Community* 1995;3:19–31.
34. Qin L, Xu H. A cross-sectional study of the effect of health literacy on diabetes prevention and control among elderly individuals with prediabetes in rural China. *BMJ Open* 2016;6:e011077.
35. Charvat H, Sasazuki S, Inoue M, *et al.* Prediction of the 10-year probability of gastric cancer occurrence in the Japanese population: the JPHC study cohort II. *Int J Cancer* 2016;138:320–31.
36. Simon AE, Juszczuk D, Smyth N, *et al.* Knowledge of lung cancer symptoms and risk factors in the U.K.: development of a measure and results from a population-based survey. *Thorax* 2012;67:426–32.
37. TT S, Goh JY, Tan J, *et al.* Level of colorectal cancer awareness: a cross sectional exploratory study among multi-ethnic rural population in Malaysia. *BMC Cancer* 2013;13:376.
38. Rutherford MJ, Ironmonger L, Ormiston-Smith N, *et al.* Estimating the potential survival gains by eliminating socioeconomic and sex inequalities in stage at diagnosis of melanoma. *Br J Cancer* 2015;112 Suppl 1:S116–S123.
39. Macleod U, Mitchell ED, Burgess C, *et al.* Risk factors for delayed presentation and referral of symptomatic cancer: evidence for common cancers. *Br J Cancer* 2009;101 Suppl 2:S92–S101.
40. McCutchan GM, Wood F, Edwards A, *et al.* Influences of cancer symptom knowledge, beliefs and barriers on cancer symptom presentation in relation to socioeconomic deprivation: a systematic review. *BMC Cancer* 2015;15:1000.
41. Hamashima C. Current issues and future perspectives of gastric cancer screening. *World J Gastroenterol* 2014;20:13767–74.
42. Yeh JM, Hur C, Ward Z, *et al.* Gastric adenocarcinoma screening and prevention in the era of new biomarker and endoscopic technologies: a cost-effectiveness analysis. *Gut* 2016;65:563–74.
43. Oldach BR, Katz ML. Health literacy and cancer screening: a systematic review. *Patient Educ Couns* 2014;94:149–57.
44. Wakao F. [The present state of cancer education in Japan and its future]. *Gan To Kagaku Ryoho* 2015;42:920–3.
45. Uthman OA, Jadidi E, Moradi T. Socioeconomic position and incidence of gastric cancer: a systematic review and meta-analysis. *J Epidemiol Community Health* 2013;67:854–60.
46. De Leeuw ED, Hox J, Dillman D. *International Handbook of survey methodology*. Routledge, 2012.
47. Lo SH, Waller J, Vrinten C, *et al.* Self-Reported and objectively recorded colorectal cancer screening participation in England. *J Med Screen* 2016;23:17–23.
48. Salimzadeh H, Bishehsari F, Delavari A, *et al.* Cancer risk awareness and screening uptake in individuals at higher risk for colon cancer: a cross-sectional study. *BMJ Open* 2016;6:e013833.