BMJ Open Characteristics of stakeholder involvement in systematic and rapid reviews: a methodological review in the area of health services research

Jonas Feldmann, Milo Alan Puhon, Margot Mütsch

To cite: Feldmann J, Puhon MA, Mütsch M. Characteristics of stakeholder involvement in systematic and rapid reviews: a methodological review in the area of health services research. *BMJ Open* 2019;9:e024587. doi:10.1136/bmjopen-2018-024587

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-024587>).

Received 11 June 2018

Revised 10 July 2019

Accepted 17 July 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Epidemiology, Biostatistics and Prevention Institute, University of Zurich, Zurich, Switzerland

Correspondence to

Dr Margot Mütsch;
margot.muetsch@uzh.ch

ABSTRACT

Objective Engaging stakeholders in reviews is considered to generate more relevant evidence and to facilitate dissemination and use. As little is known about stakeholder involvement, we assessed the characteristics of their engagement in systematic and rapid reviews and the methodological quality of included studies. Stakeholders were people with a particular interest in the research topic.

Design Methodological review.

Search strategy Four databases (Medline, Embase, Cochrane database of systematic reviews, databases of the University of York, Center for Reviews and Dissemination (CRD)) were searched based on an a priori protocol. Four types of reviews (Cochrane and non-Cochrane systematic reviews, rapid and CRD rapid reviews) were retrieved between January 2011 and October 2015, pooled by potential review type and duplicates excluded. Articles were randomly ordered and screened for inclusion and exclusion criteria until 30 reviews per group were reached. Their methodological quality was assessed using AMSTAR and stakeholder characteristics were collected.

Results In total, 57 822 deduplicated citations were detected with potential non-Cochrane systematic reviews being the biggest group (56 986 records). We found stakeholder involvement in 13% (4/30) of Cochrane, 20% (6/30) of non-Cochrane, 43% (13/30) of rapid and 93% (28/30) of CRD reviews. Overall, 33% (17/51) of the responding contact authors mentioned positive effects of stakeholder involvement. A conflict of interest statement remained unmentioned in 40% (12/30) of non-Cochrane and in 27% (8/30) of rapid reviews, but not in Cochrane or CRD reviews. At most, half of non-Cochrane and rapid reviews mentioned an a priori study protocol in contrast to all Cochrane reviews.

Conclusion Stakeholder engagement was not general practice, except for CRD reviews, although it was more common in rapid reviews. Reporting factors, such as including an a priori study protocol and a conflict of interest statement should be considered in conjunction with involving stakeholders.

INTRODUCTION

Evidence synthesis remains a rapidly growing field. There are different methodological approaches and formats depending on the

Strengths and limitations of this study

- A systematic approach to assess stakeholder engagement in systematic and rapid reviews was used to fill the gap and to provide an overview of the current situation.
- The different review types were able to inform on stakeholder involvement and pointed to potential improvements.
- With the scope of this study being restricted to health services research we intended to focus on a best-case approach where the context is considered most important and therefore, stakeholder involvement might be most regularly, but the overall findings might not be applicable beyond this domain.
- The majority of studies were from high-income countries limiting the generalisability beyond this setting.
- The perception and the reporting of stakeholder involvement might be influenced by the lack of a uniform definition of stakeholders and depend on the criteria used.

research question and the intended use of the review, such as scoping, systematic or rapid reviews as well as realist syntheses, policy briefs and health technology assessments (HTAs). We will focus on full systematic reviews as a well-established review type in health-care and on rapid reviews as an emerging one. A traditional full systematic review is a review that ‘attempts to collate all empirical evidence and that fits prespecified eligibility criteria in order to answer a specific, usually narrow research question or intervention’.¹ Within systematic reviews, Cochrane full systematic reviews were established as ‘gold standard’ in knowledge synthesis. A couple of studies compared Cochrane and non-Cochrane systematic reviews and found that the reporting of Cochrane systematic reviews was the most complete one,² that they were less prone to bias due to greater transparency in reporting as well as due to the quality criteria

used, such as the risk of bias assessment.³ Furthermore, Hopewell *et al* described that the inclusion of grey literature routinely performed in Cochrane systematic reviews limited publication bias and provided more conservative treatment effects compared with non-Cochrane systematic reviews without including grey literature.⁴

Rapid reviews are characterised by a less complex research question and aim to synthesise evidence within a shorter time period, with time frames ranging from 1 week to 9 months.^{5,6} They might, therefore, be prone to be of lower validity as a consequence of accelerating and streamlining the review process.⁵ However, in response to an increasing demand from stakeholders, rapid reviews are being performed more frequently than before.⁶ Only few formal definitions of different rapid reviews exist and few studies have examined their methodology.⁷ To cover the variety of rapid reviews we included rapid reviews listed in medical databases as well as other rapid reviews from the databases of the University of York, Center for Reviews and Dissemination (www.crd.york.ac.uk/CRDWeb), such as the Dare reviews (Database of Abstracts of Reviews of Effects) or HTAs (CRD rapid reviews). Rapid reviews target specific audiences including government policy-makers, healthcare institutions, health professionals and patient associations and research questions are often tailored to these stakeholders.^{8,9}

Stakeholder as a very broad term may include anyone affected by an issue and/or anyone who can provide input on the topic.^{10,11} Hence, not only decision makers, health professionals and their organisations are targeted, but also citizen or patients, other researchers and the media. Involving stakeholders in health services research might be possible at all stages of systematic or rapid reviews and this is often referred to as co-production. Thus, stakeholder engagement aims participatory research and might direct the research question, define the scope and context of the review as well as contribute to the literature search, the evidence synthesis and interpretation and might facilitate dissemination and use. Therefore, it enhances the relevance of findings for policy and practice and contributes to the sustainability of health systems.^{12–14} We especially focused on stakeholder involvement among health services research. This embraced 'the multi-disciplinary field of scientific studies regarding social, financing and personal factors, organisational structures and processes, health technologies, and how these factors affect access to, the quality and cost of healthcare, and ultimately, our health and well-being'.¹⁵ Here, we considered the context of conducted studies and dissemination issues as most important, and we expected health services research to be the research field where stakeholders were involved most regularly.

Study aim

To date, little is known about the extent of stakeholder engagement in systematic and rapid reviews and there have been few efforts to directly report the specific effects regarding their involvement. We consider this information

to be relevant for integrated knowledge translation, the dissemination and acceptance of systematic and rapid reviews in policy and practice. We focused on systematic and rapid reviews as they represent different types of evidence synthesis and are well-established in research and practice. In addition, reviews needed to belong to the area of health services research as an established field for systematic and rapid reviews as well as a best-case sample with a considerable extent of stakeholder involvement. We aimed to assess the extent and characteristics of stakeholder engagement in published systematic and rapid reviews and to specifically determine reporting characteristics.

METHODS

Study design

Based on the prespecified protocol this methodological review was performed to assess characteristics and reporting of stakeholder engagement in random samples of systematic and rapid reviews in the field of health services research (online supplementary file S1). A total of four types of reviews, two groups of each, systematic and rapid reviews, were assessed as they were considered to exhibit potentially differences in stakeholder involvement. This included Cochrane systematic reviews, non-Cochrane systematic reviews, rapid reviews and rapid reviews of the databases of the University of York, CRD rapid reviews. For the reporting of this study the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist was used (online supplementary file S2).

Search strategy and screening

For systematic reviews, we searched the Cochrane database of systematic reviews for articles including the term 'systematic review' in their title or abstract (Cochrane systematic reviews) and EBSCO Medline and Embase with excluding the term 'Cochrane' in title or abstract (non-Cochrane systematic reviews). Rapid reviews were searched in EBSCO Medline and Embase and in the databases of the University of York, CRD rapid reviews (www.crd.york.ac.uk/CRDWeb). Sample search strategies are depicted in online supplementary files S3a and S3b. We included studies published between January 2011 and December 2015 without language restriction. Our search was completed on 22 October 2015 and therefore, the search results for 2015 did not cover the whole year. All results were pooled by review type and duplicates were excluded based on authors, journal and publication year. For each of the four lists of potential study types, each reference was given a unique random number using the `sample()` function available in R.¹⁶ Each list was then sorted by the random number, and the articles were screened in order for inclusion and exclusion criteria by two reviewers until a total of 30 studies were reached per group (JF and MM tested their consistency in assessing the inclusion and exclusion criteria for about 10 different

studies and MM checked the assignment for about half of the study sample). Full text screening was performed to ensure the decision. One review author (JF) performed data extraction, supervised and spot-checked by a second researcher (MM). Disagreements were resolved by consensus.

Sample size and selection of review groups

As rapid reviews were often targeted to specific audiences,^{8,9} they were expected to involve stakeholders to a greater extent. We estimated a proportion of stakeholder involvement of 0.70 for rapid reviews and of 0.25 for systematic reviews, which resulted in a minimal expected difference of 0.45. For a two group-comparison, rapid and systematic reviews, with $n=25$ per group there will be an estimated power of 0.88 and an alpha of 0.023. To account for a potential subgroup analysis, we included a sample of $n=30$ reviews per group, with four review groups (Cochrane and non-Cochrane systematic reviews, rapid and CRD rapid reviews). This resulted in an overall sample of 120 studies.

Inclusion and exclusion criteria

We included reviews targeting health services research as defined by Lohr¹⁵: "Health services research is the multidisciplinary field of scientific investigation that studies how social factors, financing systems, organizational structures and processes, health technologies, and personal behaviors affect access to health care, the quality and cost of health care, and ultimately, our health and well-being. Its research domains are individuals, families, organizations, institutions, communities, and populations". Therefore, we included HTAs (eg, rapid reviews as part of HTAs) targeting effectiveness as well as meta-analyses and systematic reviews for effectiveness and utility, reviews of pharmaceutical trials under everyday conditions, reviews of basic research in care-related fields, reviews of quality research or of methodological developments in the field of health services research, reviews about the development and application of new technologies (ehealth) and the implementation of knowledge into clinical practice. We excluded reviews that focused exclusively on economic or cost analyses, performed a narrative review, an overview of reviews or a protocol of a systematic review in order not to mix so different methodological approaches. The following study types were also excluded: reviews of efficacy studies without assessing the quality of life, reviews including clinical efficacy trials phase I–III, epidemiological reviews to assess determinants or risk factors, non-human studies, establishment of databases or registries, reviews with unclear design or description of the intervention.

Data extraction

Stakeholders were defined as people with a particular interest in the research topic (but were not members of the primary research team).^{10,17} We evaluated any kind of stakeholder involvement and recorded different groups

of stakeholders: institutional healthcare providers, representatives of hospitals or community services, patients/consumers, participants of government agencies and healthcare policymakers at Federal, State and local levels, associations of health professionals and researchers (if not members of the primary study team). If available, we identified their fields and stages of involvement as well as their contribution to the study. This included all review steps, such as formulating the research question(s), determining study characteristics, contributing to the writing of the protocol, participating within the review process with searching, screening, data extraction, synthesising, interpreting the study results and/or establishing recommendations. In addition, we extracted the following data: institution where the review was performed, contact details of the corresponding author, the type of intervention, study setting, characteristics of the population (sample size, age range, sex), funding, declared conflict of interest, year of publication. The methodological quality of included articles was assessed using the AMSTAR tool, a measurement tool for the assessment of the methodological quality of systematic reviews.¹⁸ Where available, our rating was compared with an already existing one from the 'health evidence' or 'health systems evidence' platforms.^{19,20} Both quality ratings were then categorised as strong (≥ 8), moderate (4–7) or weak (< 4). In the absence of a quality assessment tool for rapid reviews, we used AMSTAR also for rapid reviews, being aware that this might result in lower rating scores due to abbreviated procedures. In addition, the use of the PRISMA checklist was recorded when mentioned in the articles.²¹ AMSTAR ratings are presented as online supplementary file S4, study characteristics as online supplementary file S5.

To ascertain stakeholder engagement as well as to assess the extent or stages of their involvement, preformulated questions were sent by email to the contact authors of the included articles. We asked them, if there had been any stakeholder involvement (yes or no) and if so, to specify the number of stakeholders, the stages of involvement and their self-assessed estimation of the effect of stakeholder engagement on the review outcome. In case of missing responses, one reminder was sent.

Patient and Public involvement

None involved.

RESULTS

Our search identified 57 822 citations remaining after exclusion of duplicates. Although, not all of these records will fulfil the inclusion criteria, such as for example, being designed as a systematic review, we assigned them as potential review group. For the screening step, the counts of the articles that needed to be screened to reach the final set of studies are depicted in figure 1.^{22–141} The excluded studies are presented in online supplementary table S1.

Figure 1 Flow diagram of study selection. CRD, databases of the University of York, Centre for Reviews and Dissemination; RR, rapid review; SR, systematic review.

Table 1 presents the characteristics of the included studies. Cochrane systematic reviews generally focused on the prevention and treatment of specific conditions, whereas 30% of non-Cochrane systematic reviews were categorised with a community-based topic, with a wide range of participants and possible outcomes and a broader range of study settings. Of note, 23% of each, rapid reviews and CRD rapid reviews, were categorised as health system intervention, because they focused on methodological questions or health quality research.

We detected noticeable differences when comparing the median AMSTAR scores: Cochrane systematic reviews and CRD rapid reviews showed higher median scores than non-Cochrane systematic reviews and rapid reviews. Where available, we also collected existing AMSTAR ratings from the 'health evidence' or 'health systems evidence' platforms. For 68% (13/19 study ratings) we found a congruent classification as strong, moderate or weak, respectively (online supplementary file S4). In non-Cochrane reviews information about methodological specifications of included studies were often lacking. As an example, all 30 Cochrane systematic reviews (100%) mentioned a pre-existing review protocol, whereas a high amount of non-Cochrane systematic reviews (59%) and of rapid reviews (43%) did not clarify whether there had been a protocol or not. Similarly, all 30 Cochrane systematic reviews and all but one CRD rapid reviews (97%) included a conflict of interest statement, whereas 40% of non-Cochrane systematic reviews and 27% of rapid reviews lacked such a paragraph. All reviews were funded by national or international governmental or institutional sources. Only one rapid review reported the acceptance of additional funding from a pharmaceutical company.

Characteristics of the study population by type of included review are presented in online supplementary table S2.

Any stakeholder involvement was mentioned in 13% of Cochrane systematic reviews, 20% of non-Cochrane systematic reviews, 43% of rapid reviews and 93% of CRD rapid reviews. Except for CRD rapid reviews, where the involvement of stakeholders was routinely reported in the methods or appendix section, about half of the stakeholder involvement was reported in the articles and the other half was confirmed via email by contact authors and remained unmentioned in the reviews. With 67% the email response rate was highest for rapid reviews. When comparing the amount of stakeholder involvement in different review topics, there were notable differences: the proportion of stakeholder involvement in reviews focusing on prevention or treatment of specific conditions was lower than in those targeting health system interventions. Interestingly, the proportion of stakeholder involvement has been quite constant between the years 2012 (5/15, 33%) and 2014 (12/34, 34%) even though the amount of rapid reviews has increased substantially (table 2). There was, however, a remarkable increase in the proportion of stakeholder involvement for the year 2015 (22/32, 69%), but this was mostly due to the fact that a majority of CRD rapid reviews, which presented the highest proportion of stakeholder involvement, was indexed in 2015.

CRD rapid reviews turned out to have by far the highest proportion of reported stakeholder involvement. The types of stakeholders engaged were listed in the appendix of 26/30 articles, but it was not specified what they had specifically contributed to the review, for example, if their contribution had affected the final results and conclusions of the reviews (table 3). One author mentioned in the review that the reason for involving stakeholders was to understand the clinical perspective.

We detected stakeholder involvement in 43% (13/30) of rapid reviews. In contrast to CRD rapid reviews, there was usually a small number of stakeholders engaged. They were involved at different phases, such as determining study characteristics, formulating the research question, within the review process in general, and less commonly in protocol writing, result synthesis and the interpretation of findings. One article involved stakeholder throughout all stages, in one case the phase of involvement was not specified. In total, 85% of the authors confirmed, that the stakeholders had significantly contributed to the review. Of these, 62% mentioned that involving stakeholders led to an increased impact of the review and enabled to focus on the needs of target groups, which made the review more relevant to for example, patients or policymakers. Stakeholders had commissioned one third of the rapid reviews or had asked for evidence. Only two of 13 authors (15%) did not mention any substantial effect. In two cases one of the funding sources was involved as stakeholder.

We identified stakeholder involvement in 20% (6/30) of non-Cochrane systematic reviews. In four cases the stakeholder involvement was confirmed via email, two articles mentioned the involvement of stakeholders. Only two types of stakeholders were reported: policy-makers and researchers. If mentioned, all phases of

Table 1 Characteristics of included reviews by review type

	Cochrane SR (%)	Non-Cochrane SR (%)	RR (%)	CRD RR (%)	Total (%)
Random sample (% of total per group)	30 (6)	30 (0.1)	30 (14)	30 (32)	120 (0.2)
Geographic location of the corresponding author (% of random sample)					
Africa	2 (7)	0	0	0	2 (2)
Asia	5 (17)	3 (10)	1 (3)	0	9 (8)
Australia	3 (10)	5 (17)	2 (7)	0	10 (8)
Europe	11 (37)	13 (43)	13 (43)	2 (7)	39 (33)
North America	7 (23)	8 (27)	14 (47)	28 (93)	57 (48)
South America	2 (7)	1 (3)	0	0	3 (3)
Thematic focus					
Prevention	11 (37)	5 (17)	4 (13)	7 (23)	27 (23)
Treatment	18 (60)	14 (47)	15 (50)	16 (53)	63 (53)
Health system	1 (3)	2 (7)	7 (23)	7 (23)	17 (14)
Community-based*	0	9 (30)	4 (13)	0	13 (11)
Funding					
Governmental, institutional or WHO	30 (100)	30 (100)	30 (100)	30 (100)	120 (100)
Additional funding by company	0	0	1 (3)	0	1 (1)
Conflict of interest (COI)					
Declared none	21 (70)	15 (50)	19 (63)	29 (97)	84 (70)
COI declared	9 (30)	3 (10)	3 (10)	0	15 (13)
Unmentioned	0	12 (40)	8 (27)	1 (3)	21 (18)
Study protocol					
Yes, mentioned in article	30 (100)	5 (17)	4 (13)	1 (3)	40 (33)
Yes, mentioned in correspondence	0	8 (27)	9 (30)	0	17 (14)
No, confirmed by correspondence	0	2 (7)	4 (13)	0	6 (5)
Unmentioned	0	15 (50)	13 (43)	29 (97)	57 (48)
AMSTAR rating					
Median (range)	11 (7, 11)	7 (3, 11)	4 (2, 10)	8 (5, 11)	8 (2, 11)

AMSTAR: measurement tool for the 'assessment of the methodological quality of systematic reviews', not applicable questions were not counted. Higher AMSTAR scores indicate higher quality.

*With exclusion of treatment or preventive interventions.

CRD, databases of the University of York, Center for Reviews and Dissemination; RR, rapid review; SR, systematic review.

involvement occurred about equally. A total 67% (4/6) study authors reported significant benefit from the stakeholders' contributions. One author mentioned that stakeholder supported the researchers to understand different perspectives of the problem, and 33% (2/6) reported no effect. In one case, the stakeholder was involved as funding source.

Cochrane systematic reviews turned out to be the group with the smallest amount of stakeholder involvement. All stakeholder engagement was confirmed and specified by email. The types of stakeholders engaged were patients, caregivers, professional organisations and researchers. They were mostly involved for providing feedback during the review process, in one case they contributed in formulating the review question. One review involved

stakeholders at all stages. A total of 50% (2/4) of the authors reported substantial benefit, the other 50% (2/4) reported no significant effect on study results or conclusions. Two authors had involved stakeholders to make the review more relevant for its target audience, one author claimed that stakeholders helped him to refine the research question. One author involved stakeholders to get direct consumer feedback before publishing the review. No stakeholders were involved in funding.

DISCUSSION

The involvement of stakeholders varies by review type. Our findings suggest that rapid reviews tend to involve stakeholders more than twice as frequently than systematic

Table 2 General characteristics of stakeholder involvement (SI) by review type

	Cochrane SR (%) (n=30)	Non-Cochrane SR (%) (n=30)	RR (%) (n=30)	CRD RR (%) (n=30)	Total (%) (N=120)
SI mentioned in article	2 (7)	2 (7)	6 (20)	28 (93)	38 (32)
SI only mentioned in correspondence	2 (7)	4 (13)	7 (23)	0	13 (11)
Total SI	4 (13)	6 (20)	13 (43)	28 (93)	51 (43)
SI per year					
2011	0/5 (0)	0/1 (0)	0/3 (0)	0/0 (0)	0/9 (0)
2012	3/7 (43)	1/5 (20)	1/3 (33)	0/0 (0)	5/15 (33)
2013	0/9 (0)	0/6 (0)	3/5 (60)	9/10 (90)	12/30 (40)
2014	1/7 (14)	3/9 (30)	6/15 (40)	2/3 (67)	12/34 (35)
2015	0/2 (0)	2/9 (22)	3/4 (75)	17/17 (100)	22/32 (69)
Number of stakeholders involved per review (% of SI)					
1	2 (50)	2 (33)	2 (15)	0	6 (12)
2–4	1 (25)	1 (17)	4 (31)	2 (7)	8 (16)
>4	1 (25)	3 (50)	4 (31)	26 (93)	34 (67)
Unspecified	0	0	3 (23)	0	3 (6)
Types of stakeholders (multiple roles possible) (% of SI)					
Patients/consumers	2 (50)	0	2 (15)	5 (18)	9 (18)
Professional organisations	1 (25)	0	0	5 (18)	6 (12)
Caregivers	2 (50)	0	3 (23)	23 (82)	28 (55)
Researchers	1 (25)	3 (50)	2 (15)	0	6 (12)
Policymakers	0	3 (50)	4 (31)	3 (11)	10 (20)
Unspecified	0	0	2 (15)	4 (14)	6 (12)
Funding source (% of SI)					
Governmental or institutional funding	4 (100)	6 (100)	13 (100)	28 (100)	51 (100)
Funding source involved as stakeholder	0	1 (17)	2 (15)	0	3 (6)
Thematic focus (% per review type per topic)					
Prevention	0/11 (0)	0/5 (0)	1/4 (25)	6/7 (86)	7/27 (26)
Treatment	4/18 (22)	3/14 (21)	6/15 (40)	16/16 (100)	29/63 (46)
Health system	0/1 (0)	1/2 (50)	5/7 (71)	6/7 (86)	12/17 (71)
Community-based	0/0 (0)	2/9 (22)	1/4 (25)	0/0 (0)	3/13 (23)

CRD, databases of the University of York, Center for Reviews and Dissemination; RR, rapid review; SR, systematic review.

reviews. On average, they also involved a greater number of stakeholders per review. In addition, we detected considerable differences in the phases in which stakeholders were involved. Rapid reviews involved them at very early stages of the review process, such as determining the intentional study characteristics or formulating the research question. Furthermore, it seemed to be much more common for policymakers, who were the most frequent group of stakeholders involved in rapid reviews, or other stakeholder groups, to substantially contribute to rapid reviews than to systematic reviews.

Of note, the majority of rapid and non-Cochrane systematic review authors reported that the involved stakeholders had positive and considerable effects on the study results, making the review more relevant for the targeted audience. However, the minority of rapid and

non-Cochrane systematic review authors reporting no effect of stakeholder involvement on the review process, may illustrate that the stakeholders' contribution to non-Cochrane systematic reviews was seen as an additional and welcomed feature to the review, but not as a substantial part. This shows the importance that within and between review types an information and experiences exchange between researchers could benefit the stakeholder engagement.

A strength of this study is to provide an overview with its rather low amount of stakeholder engagement in the assessed review types despite known benefits regarding the use of evidence in policy and practice. With respect to the different procedures used by review types, the reported experiences of stakeholder engagement, for example, in rapid reviews, could further benefit and

Table 3 Engagement characteristics of stakeholder involvement (SI) by review type

	Cochrane SR (%) (n=30)	Non-Cochrane SR (%) (n=30)	RR (%) (n=30)	CRD RR (%) (n=30)	Total (%) (N=120)
Stages of SI (multiple stages possible) (% of SI)					
Research question	1 (25)	1 (17)	4 (31)	0	6 (12)
Study characteristics	0	0	6 (46)	0	6 (12)
Protocol writing	0	1 (17)	2 (15)	0	3 (6)
Review process	3 (75)	1 (17)	5 (38)	0	9 (18)
Interpretation of findings	0	1 (17)	1 (8)	0	2 (4)
Result synthesis	0	0	2 (15)	0	2 (4)
Multiple or all stages	1 (25)	2 (33)	1 (8)	28 (100)	32 (63)
Unspecified	0	2 (33)	1 (8)	0	3 (6)
Reasons mentioned for SI (% of SI)					
Stakeholder commissioned/requested review	0	1 (17)	4 (31)	0	5 (10)
To get consumer feedback	1 (25)	0	0	0	1 (2)
To refine research question	1 (25)	0	0	0	1 (2)
To make the review more relevant for target audience	2 (50)	0	8 (62)	0	10 (20)
To understand clinical perspective	0	1 (17)	0	1 (4)	2 (4)
Unspecified	0	4 (67)	1 (7)	27 (96)	32 (63)
Effect of SI mentioned by authors (% of SI)					
Substantial or positive effect	2 (50)	4 (67)	11 (85)	0	17 (33)
Negative effect	0	0	0	0	0
No effect mentioned	2 (50)	2 (33)	2 (15)	28 (100)	34 (67)
Explicit encouragement for involving stakeholders in further studies (% of reviews)	0	1 (3)	1 (3)	1 (3)	3 (2)

CRD, databases of the University of York, Center for Reviews and Dissemination; RR, rapid review; SR, systematic review.

facilitate stakeholder engagement in other review types, for example in systematic reviews.

One limitation of this study is the low response rate to our emails by contact authors. Given the fact that nearly half of our emails to study authors remained unanswered or could not be sent, the rate of unmentioned stakeholder involvement might still be higher than our numbers suggest. Of note, there was no considerable difference between systematic reviews and rapid reviews in the percentage of unmentioned stakeholder involvement.

Although our search is not very recent this article highlights the current situation and there is a call for action. Of course, an updated assessment might be needed in the years following.

Although when using a broad definition of the term 'stakeholder' including everyone with a particular interest in the research topic (but who are not members of the primary research team), this term was not consistently used by contact authors. In their emails, some authors listed peer reviewers as stakeholders or one author erroneously mentioned members of the research team, who performed literature research and data extraction as stakeholders. In addition, interdisciplinary knowledge exchange is an important part of evidence-based research,

but there is a difference whether experts from other fields were included as researchers in the study team or whether they were considered as stakeholders. We did not count any experts or peer reviewers as stakeholders. Furthermore, discussion is needed, whether the funding body of a review might contribute as a stakeholder and how a potentially associated conflict of interest could be avoided.

The reported involvement of stakeholders corresponded with Cottrell *et al* who mentioned that stakeholders might contribute to different study types and evidence phases.¹⁴² Based on, Keown *et al*¹⁴³ concluded, that stakeholder involvement led to an increased relevance and depth of review findings, more clarity in defining research questions, broader dissemination of their results and increased awareness of target groups. Although they mentioned that this engagement process required flexibility and might be resource-intensive and time-intensive, they nevertheless concluded that involving stakeholders facilitated implementation and should be indispensable for future research.

Although the overall proportion of stakeholder involvement has not yet increased prominently in the past few years, there have still been considerable efforts in creating

standardised procedures for involving stakeholders in evidence synthesis. Keown *et al* identified five opportunities in the systematic review process, where potential stakeholders could be engaged on a regular base.¹⁴³ The CRD rapid reviews included were mainly performed by Health Quality Ontario (Canada) and routinely held expert panels, including physicians, caregivers and sometimes, consumer representatives and professional organisations. The fact that they included stakeholders on a regular basis might confirm, what Keown *et al*¹⁴³ had already suggested: the engagement of audience members interested in or affected by the investigated topic definitively resulted in more benefits than limitations. We, therefore, suggest that future researcher involve stakeholders more broadly in the process of evidence synthesis, to increase the relevance and acceptance of the knowledge transfer. One example of an organisation ensuring that stakeholders are involved in research is the James Lind Alliance (<http://www.jla.nihr.ac.uk>).

In future, the reporting of stakeholder involvement should be improved and its effects better evaluated and communicated. Stakeholder engagement could also be included in reporting checklists of all review types.

Acknowledgements We are indebted to Julia Kreis and Aline Flatz for reviewing the protocol and our thanks go to Sarah Haile for generating the random lists. Parts of this work have been presented at the Global Evidence Summit September 2017 in Cape Town, SA.

Contributors All authors conceived and designed the study. JF collected and interpreted the data and essentially contributed to the writing of the manuscript. MAP and MM supervised the overall research and critically revised the manuscript. All authors approved the final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Khangura S, Konnyu K, Cushman R, *et al*. Evidence summaries: the evolution of a rapid review approach. *Syst Rev* 2012;1:10.
2. Page MJ, Shamseer L, Altman DG, *et al*. Epidemiology and reporting characteristics of systematic reviews of biomedical research: a cross-sectional study. *PLoS Med* 2016;13:e1002028.
3. Jadad AR, Cook DJ, Jones A, *et al*. Methodology and reports of systematic reviews and meta-analyses: a comparison of Cochrane reviews with articles published in paper-based journals. *JAMA* 1998;280:278–80.
4. Hopewell S, Clarke MJ, Lefebvre C, *et al*. Handsearching versus electronic searching to identify reports of randomized trials. *Cochrane Database Syst Rev* 2007;24.
5. Ganann R, Ciliska D, Thomas H. Expediting systematic reviews: methods and implications of rapid reviews. *Implementation Sci* 2010;5.
6. Watt A, Cameron A, Sturm L, *et al*. Rapid versus full systematic reviews: validity in clinical practice? *ANZ J Surg* 2008;78:1037–40.
7. Tricco AC, Antony J, Zarin W, *et al*. A scoping review of rapid review methods. *BMC Med* 2015;13:224.
8. Watt A, Cameron A, Sturm L, *et al*. Rapid reviews versus full systematic reviews: an inventory of current methods and practice in health technology assessment. *Int J Technol Assess Health Care* 2008;24:133–9.
9. Hailey DM. Health technology assessment in Canada: diversity and evolution. *Med J Aust* 2007;187:286–8.
10. Mallery C, Ganachari D, Fernandez J, *et al*. *Innovative methods in Stakeholder engagement: an environmental scan*. Rockville, MD: Agency for Healthcare Research and Quality, 2012.
11. Concannon TW, Fuster M, Saunders T, *et al*. A systematic review of stakeholder engagement in comparative effectiveness and patient-centered outcomes research. *J Gen Intern Med* 2014;29:1692–701.
12. Boote J, Wong R, Booth A. 'Talking the talk or walking the walk?' A bibliometric review of the literature on public involvement in health research published between 1995 and 2009. *Health Expect* 2015;18:44–57.
13. Kreis J, Puhan MA, Schünemann HJ, *et al*. Consumer involvement in systematic reviews of comparative effectiveness research. *Health Expect* 2013;16:323–37.
14. Morley RF, Norman G, Golder S, *et al*. A systematic scoping review of the evidence for consumer involvement in organisations undertaking systematic reviews: focus on Cochrane. *Res Involv Engagem* 2016;2.
15. Lohr KN, Steinwachs DM. Health services research: an evolving definition of the field. *Health Serv Res* 2002;37:7–9.
16. R Core Team. R: a language and environment for statistical computing. Available: <https://www.R-project.org/2018>
17. Pollock A, Campbell P, Struthers C, *et al*. Stakeholder involvement in systematic reviews: a protocol for a systematic review of methods, outcomes and effects. *Res Involv Engagem* 2017;3.
18. Shea BJ, Grimshaw JM, Wells GA, *et al*. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7:10.
19. McMaster University Hamilton (Canada). Health evidence 2016. Available: <http://www.healthevidence.org> [Accessed 06.04.2016].
20. Mc Master Health Forum (McMaster University Hamilton (Canada). Health systems evidence 2016. Available: <https://www.healthsystemsevidence.org> [Accessed 06.04.2016].
21. Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006–12.
22. Akl EA, Oxman AD, Herrin J, *et al*. Framing of health information messages. *Cochrane Database of Systematic Reviews* 2011;22.
23. Bjelakovic G, Gluud LL, Nikolova D, *et al*. Vitamin D supplementation for prevention of cancer in adults. *Cochrane Database Syst Rev* 2014;366.
24. Carney T, Myers BJ, Louw J, *et al*. Brief school-based interventions and behavioural outcomes for substance-using adolescents. *Cochrane Database Syst Rev* 2014;(2).
25. Chen W, Zhang Y, Li X, *et al*. Chinese herbal medicine for diabetic peripheral neuropathy. *Cochrane Database Syst Rev* 2013;10.
26. Cheuk Daniel KL, Chiang Alan KS, Lee Tsz L, *et al*. Vaccines for prophylaxis of viral infections in patients with hematological malignancies. *Cochrane Database of Syst Rev* 2011;3.
27. Fede G, Germani G, Gluud C, *et al*. Propylthiouracil for alcoholic liver disease. *Cochrane Database Syst Rev* 2011;79.
28. Fonner Virginia A, Denison J, Kennedy Caitlin E, *et al*. Voluntary counseling and testing (VCT) for changing HIV-related risk behavior in developing countries. *Cochrane Database of Syst Rev* 2012;9.
29. Garrison SR, Allan GM, Sekhon RK, *et al*. Magnesium for skeletal muscle cramps. *Cochrane Database Syst Rev* 2012;173.
30. Jefferson T, Rivetti A, Di Pietrantonj C, *et al*. Vaccines for preventing influenza in healthy children. *Cochrane Database Syst Rev* 2012;8.
31. Jin H, Leng Q, Li C, *et al*. Dietary flavonoid for preventing colorectal neoplasms. *Cochrane Database Syst Rev* 2012;17.
32. Jorge EC, Jorge EN, El Dib RP, *et al*. Early light reduction for preventing retinopathy of prematurity in very low birth weight infants. *Cochrane Database Syst Rev* 2013;91.
33. Junker AE, Als-Nielsen B, Gluud C, *et al*. Dopamine agents for hepatic encephalopathy. *Cochrane Database Syst Rev* 2014;30.
34. Karmali KN, Davies P, Taylor F, *et al*. Promoting patient uptake and adherence in cardiac rehabilitation. *Cochrane Database Syst Rev* 2014;15.
35. Kwong Joey SW. Yoga for secondary prevention of coronary heart disease. *Cochrane Database Syst Rev* 2015;7.
36. Lager KE, Mistri AK, Khunti K, *et al*. Interventions for improving modifiable risk factor control in the secondary prevention of stroke. *Cochrane Database Syst Rev* 2014;52.

37. Marti-Carvajal AJ, Knight-Madden JM, Martinez-Zapata MJ, *et al.* Interventions for treating leg ulcers in people with sickle cell disease. *Cochrane Database Syst Rev* 2014;81.
38. Moi JH, Sriranganathan MK, Edwards CJ, *et al.* Lifestyle interventions for acute gout. *Cochrane Database Syst Rev* 2013;11.
39. Nava F, Ghilotti F, Maggi L, *et al.* Biologics, colchicine, corticosteroids, immunosuppressants and interferon-alpha for neuro-Behçet's syndrome. *Cochrane Database Syst Rev* 2014;12.
40. Njei B, Kongnyuy EJ, Kumar S, *et al.* Optimal timing for antiretroviral therapy initiation in patients with HIV infection and concurrent cryptococcal meningitis. *Cochrane Database Syst Rev* 2013;50(11).
41. Charles O I, Nagpal S, Musekiwa A, *et al.* Home- or community-based programmes for treating malaria. *Cochrane Database Syst Rev* 2013;5.
42. Olsen O, Clausen JA, Cochrane Pregnancy and Childbirth Group. Planned Hospital birth versus planned home birth. *Cochrane Database Syst Rev* 2012;312.
43. Overdevest Gijsbert M, Jacobs W, Vleggeert-Lankamp C, *et al.* Effectiveness of posterior decompression techniques compared with conventional laminectomy for lumbar stenosis. *Cochrane Database Syst Rev* 2015;3.
44. Peckham EJ, Nelson EA, Greenhalgh J, *et al.* Homeopathy for treatment of irritable bowel syndrome. *Cochrane Database Syst Rev* 2013;26.
45. Rahimi-Movaghar A, Amin-Esmaili M, Hefazi M, *et al.* Pharmacological therapies for maintenance treatments of opium dependence. *Cochrane Database Syst Rev* 2013;66.
46. Shields L, Zhou H, Pratt J, *et al.* Family-centred care for hospitalised children aged 0-12 years. *Cochrane Database Syst Rev* 2012;45.
47. Smart NA, Marshall BJ, Daley M, *et al.* Low-Fat diets for acquired hypercholesterolaemia. *Cochrane Database Syst Rev* 2011;20.
48. Taylor F, Huffman Mark D, Macedo Ana F, *et al.* Statins for the primary prevention of cardiovascular disease. *Cochrane Database Syst Rev* 2013;1.
49. Taylor Matthew J, Rudkin L, Bullemor-Day P, *et al.* Strategies for managing sexual dysfunction induced by antidepressant medication. *Cochrane Database Syst Rev* 2013;5.
50. Tejani AM, Wasdell M, Spiwak R, *et al.* Carnitine for fatigue in multiple sclerosis. *Cochrane Database Syst Rev* 2012;218.
51. Thomas RE, Lorenzetti D, Spragins W, *et al.* Mentoring adolescents to prevent drug and alcohol use. *Cochrane Database Syst Rev* 2011;1.
52. Agarwal R, Goldenberg M, Perry R, *et al.* The quality of life of adults with attention deficit hyperactivity disorder: a systematic review. *Innov Clin Neurosci* 2012;9:10-21.
53. Ahmed AA, Alsharief E, Alsharief A. Intensive versus conventional glycemic control: what is best for patients with type 2 diabetes? *Diabetes Metab Syndr* 2013;7:48-51.
54. Albrecht B, Staiger PK, Hall K, *et al.* Benzodiazepine use and aggressive behaviour: a systematic review. *Aust N Z J Psychiatry* 2014;48:1096-114.
55. Bak A, Browne J, Loveday HP, *et al.* Systematic review of nutritional interventions to prevent healthcare-associated infections in undernourished elderly. *Proceedings of the Nutrition Society* 2013;72.
56. Blasco MA, Redleaf MI. Cochlear implantation in unilateral sudden deafness improves tinnitus and speech comprehension: meta-analysis and systematic review. *Otol Neurotol* 2014;35:1426-32.
57. Bourke L, Homer KE, Thaha MA, *et al.* Interventions to improve exercise behaviour in sedentary people living with and beyond cancer: a systematic review. *Br J Cancer* 2014;110:831-41.
58. Chien T-J, Liu C-Y, Chang Y-F, *et al.* Acupuncture for treating aromatase Inhibitor-Related arthralgia in breast cancer: a systematic review and meta-analysis. *The Journal of Alternative and Complementary Medicine* 2015;21:251-60.
59. Collins CE, Burrows TL, Bray J, *et al.* Effectiveness of parent-centred interventions for the prevention and treatment of childhood overweight and obesity in community settings: a systematic review. *JBIM Database System Rev Implement Rep* 2013;11:180-257.
60. Foersch M, Jacobs C, Wriedt S, *et al.* Effectiveness of maxillary protraction using facemask with or without maxillary expansion: a systematic review and meta-analysis. *Clin Oral Investig* 2015;19:1181-92.
61. Gibbs M, Winsper C, Marwaha S, *et al.* Cannabis use and mania symptoms: a systematic review and meta-analysis. *European Psychiatry* 2015;30.
62. Guerin M, Grimmer K, Kumar S. Community services' involvement in the discharge of older adults from hospital into the community. *Int J Integr Care* 2013;13:e032.
63. MB H, Xu H, Bai PD, *et al.* Obesity has multifaceted impact on biochemical recurrence of prostate cancer: a dose-response meta-analysis of 36,927 patients. *Medical Oncology* 2014;31.
64. Johantgen M, Fountain L, Zangaro G, *et al.* Comparison of labor and delivery care provided by certified nurse-midwives and physicians: a systematic review, 1990 to 2008. *Women's Health Issues* 2012;22:e73-81.
65. Lankhaar JAC, de Vries WR, Jansen JACG, *et al.* Impact of overt and subclinical hypothyroidism on exercise tolerance: a systematic review. *Res Q Exerc Sport* 2014;85:365-89.
66. Llewellyn A, Simmonds M, Owen CG, *et al.* Childhood obesity as a predictor of morbidity in adulthood: a systematic review and meta-analysis. *Obes Rev* 2016;17:56-67.
67. Maglione MA, Das L, Raean L, *et al.* Safety of vaccines used for routine immunization of US children: a systematic review. *Pediatrics* 2014;134:325-37.
68. Matsuda A, Yamaoka K, Tango T, *et al.* Effectiveness of psychoeducational support on quality of life in early-stage breast cancer patients: a systematic review and meta-analysis of randomized controlled trials. *Quality of Life Research* 2014;23:21-30.
69. Panagioti M, Gooding PA, Triantafyllou K, *et al.* Suicidality and posttraumatic stress disorder (PTSD) in adolescents: a systematic review and meta-analysis. *Soc Psychiatry Psychiatr Epidemiol* 2015;50:525-37.
70. Rodgers M, Asaria M, Walker S, *et al.* The clinical effectiveness and cost-effectiveness of low-intensity psychological interventions for the secondary prevention of relapse after depression: a systematic review. *Health Technol Assess* 2012;16:1-129.
71. Rolka EJ, Silverman MJ. A systematic review of music and dyslexia. *The Arts in Psychotherapy* 2015;46:24-32.
72. Rozet F, Bastide C, Beuzebec P, *et al.* Prise en charge des Tumeurs de la prostate à faible Risque évolutif. *Progrès en Urologie* 2015;25:1-10.
73. Rush CJ, Campbell RT, Jhund PS, *et al.* Falling cardiovascular mortality in heart failure with reduced ejection fraction and implications for clinical trials. *JACC Heart Fail* 2015;3:603-14.
74. Scott IC, Tan R, Stahl D, *et al.* The protective effect of alcohol on developing rheumatoid arthritis: a systematic review and meta-analysis. *Rheumatology* 2013;52:856-67.
75. Siffredi V, Anderson V, Leventer RJ, *et al.* Neuropsychological profile of agenesis of the corpus callosum: a systematic review. *Dev Neuropsychol* 2013;38:36-57.
76. Sipe TA, Finnie RKC, Knopf JA, *et al.* Effects of mental health benefits legislation: a community guide systematic review. *Am J Prev Med* 2015;48:755-66.
77. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ* 2012;346:e7492.
78. Tricco AC, Antony J, Khan PA, *et al.* Safety and effectiveness of dipeptidyl peptidase-4 inhibitors versus intermediate-acting insulin or placebo for patients with type 2 diabetes failing two oral antihyperglycaemic agents: a systematic review and network meta-analysis. *BMJ Open* 2014;4:e005752.
79. Vasconcellos F, Seabra A, Katzarzyk PT, *et al.* Physical activity in overweight and obese adolescents: systematic review of the effects on physical fitness components and cardiovascular risk factors. *Sports Medicine* 2014;44:1139-52.
80. Wang Z. Age-Dependent decline of association between obesity and mortality: a systematic review and meta-analysis. *Obes Res Clin Pract* 2015;9:1-11.10.1016/j.orcp.2014.01.006
81. Wu S, Cohen D, Shi Y, *et al.* Economic analysis of physical activity interventions. *Am J Prev Med* 2011;40:149-58.
82. Canadian Agency for Drugs and Technologies in Health. *Sedation and anesthesia options for diagnostic procedures: a review of clinical effectiveness and guidelines.* Ottawa (ON: Canadian Agency for Drugs and Technologies in Health, 2015.
83. Attree P, French B, Milton B, *et al.* The experience of community engagement for individuals: a rapid review of evidence. *Health Soc Care Community* 2011;19:250-60.
84. Barry MM, D'Eath M, Sixsmith J. Interventions for improving population health literacy: insights from a rapid review of the evidence. *J Health Commun* 2013;18:1507-22.
85. Beall RF, Baskerville N, Golfam M, *et al.* Modes of delivery in preventive intervention studies: a rapid review. *Eur J Clin Invest* 2014;44:688-96.
86. Beck CR, Sokal R, Arunachalam N, *et al.* Neuraminidase inhibitors for influenza: a review and public health perspective in the aftermath of the 2009 pandemic. *Influenza Other Respi Viruses* 2013;7:14-24.

87. Boyce T, Holmes A. Addressing health inequalities in the delivery of the human papillomavirus vaccination programme: examining the role of the school nurse. *PLoS One* 2012;7:e43416.
88. Brunton G, Paraskeva N, Caird J, *et al.* Psychosocial predictors, assessment, and outcomes of cosmetic procedures: a systematic rapid evidence assessment. *Aesthetic Plast Surg* 2014;38:1030–40.
89. Butow P, Price MA, Shaw JM, *et al.* Clinical pathway for the screening, assessment and management of anxiety and depression in adult cancer patients: Australian guidelines. *Psychooncology* 2015;24:987–1001.
90. Costello RB, Lentino CV, Boyd CC, *et al.* The effectiveness of melatonin for promoting healthy sleep: a rapid evidence assessment of the literature. *Nutr J* 2014;13.
91. Crawford C, Lee C, Bingham J, *et al.* Sensory art therapies for the self-management of chronic pain symptoms. *Pain Medicine* 2014;15:S66–S75.
92. Crawford C, Lee C, May T, *et al.* Physically oriented therapies for the self-management of chronic pain symptoms. *Pain Medicine* 2014;15:S54–S65.10.1111/pme.12410
93. Dorrestijn PM, Ipenburg NA, Murphy KJ, *et al.* Rapid systematic review of normal audiometry results as a predictor for benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg* 2014;150:919–24.
94. Geddes R, Frank J, Haw S. A rapid review of key strategies to improve the cognitive and social development of children in Scotland. *Health Policy* 2011;101:20–8.
95. Hithersay R, Strydom A, Moulster G, *et al.* Carer-led health interventions to monitor, promote and improve the health of adults with intellectual disabilities in the community: a systematic review. *Res Dev Disabil* 2014;35:887–907.
96. Humphreys DK, Ogilvie D. Synthesising evidence for equity impacts of population-based physical activity interventions: a pilot study. *International Journal of Behavioral Nutrition and Physical Activity* 2013;10.
97. Jiwa M, McManus A, Dadich A. Continuity of cancer care: where do primary care practitioners fit in? *Cancer Forum* 2013;37:31–4.
98. Kreindler SA, Cui Y, Metge CJ, *et al.* Patient characteristics associated with longer emergency department stay: a rapid review. *Emerg Med J* 2016;33.
99. Kwok E, Konnyu KJ, Skidmore B, *et al.* Effectiveness and safety of emergency department short-stay units: a rapid review. *Canadian Journal of Emergency Medicine* 2012;14:S5–S6.
100. Lal S, Adair CE. E-mental health: a rapid review of the literature. *Psychiatric Services* 2014;65:24–32.
101. Lee C, Crawford C, Schoemaker E, *et al.* Movement therapies for the self-management of chronic pain symptoms. *Pain Medicine* 2014;15:S40–S53.
102. Lee C, Crawford C, Swann S, *et al.* Multimodal, integrative therapies for the self-management of chronic pain symptoms. *Pain Medicine* 2014;15:S76–S85.
103. Mann R, Gilbody S. Validity of two case finding questions to detect postnatal depression: a review of diagnostic test accuracy. *J Affect Disord* 2011;133:388–97.
104. McMurrin M. Individual-Level interventions for alcohol-related violence: a rapid evidence assessment. *Criminal Behaviour and Mental Health* 2012;22:14–28.
105. Menear M, Briand C. Implementing a continuum of evidence-based psychosocial interventions for people with severe mental illness: part 1—Review of major initiatives and implementation strategies. *The Canadian Journal of Psychiatry* 2014;59:178–86.
106. Moe-Byrne T, Chambers D, Harden M, *et al.* Behaviour change interventions to promote prescribing of generic drugs: a rapid evidence synthesis and systematic review. *BMJ Open* 2014;4:e004623.
107. Muñoz K, Caballero A, White K. Effectiveness of questionnaires for screening hearing of school-age children: a comprehensive literature review. *Int J Audiol* 2014;53:910–4.
108. Ndumbe-Eyoh S, Moffatt H. Intersectional action for health equity: a rapid systematic review. *BMC Public Health* 2013;13:1056.
109. Saeed S, Golfam M, Beall RF, *et al.* Effectiveness of individual-focused interventions to prevent chronic disease. *Eur J Clin Invest* 2014;44:882–90.
110. Trivedy CR, Cooke MW, Visits UR. Unscheduled return visits (URV) in adults to the emergency department (ED): a rapid evidence assessment policy review. *Emerg Med J* 2015;32:324–9.
111. Yu J, Wang Y, Li Y, *et al.* The dA Vinci surgical system: a rapid review of the clinical and economic evidence. *Value Health* 2014;17.
112. Health Technology Assessment Database. Aerobic exercise training in patients with heart failure: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000505/frame.html>
113. Health Technology Assessment Database. Criteria for referral to home care: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000501/frame.html>
114. Health Technology Assessment Database. Physical activity counselling: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000508/frame.html>
115. Health Technology Assessment Database. Care coordination for postacute stroke, chronic obstructive pulmonary disease, and heart failure clients: an economic rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000492/frame.html>
116. Health Technology Assessment Database. Sodium restriction in heart failure: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000504/frame.html>
117. Health Technology Assessment Database. Cognitive-Behavioural therapy for anxiety and depression in patients with chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000486/frame.html>
118. Health Technology Assessment Database. Communication of discharge instructions: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000500/frame.html>
119. Health Technology Assessment Database. Prophylactic antibiotics for individuals with chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000489/frame.html>
120. Health Technology Assessment Database. Respiratory therapy services in home care for individuals with chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000491/frame.html>
121. Health Technology Assessment Database. Exercise programs after pulmonary rehabilitation for patients with chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000488/frame.html>
122. Baidoo-sonso S. Effect of supportive interventions on informal caregivers of people at the end of life: a rapid review (structured Abstract), 2014. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000095/frame.html>
123. Brenner S. Optimal timing to begin an active rehabilitation program after a hip fracture: a rapid review (structured Abstract), 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001040/frame.html>
124. Brenner S. Effectiveness and safety of thrombolytics for the treatment of ischemic stroke: a rapid review (structured Abstract), 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001030/frame.html>
125. Bungay H, Vella-Burrows T. The effects of participating in creative activities on the health and well-being of children and young people: a rapid review of the literature. *Perspect Public Health* 2013;133:44–52.
126. Ghazipura M. Optimal timing for antibiotic administration in patients with community-acquired pneumonia: a rapid review (structured Abstract), 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001051/frame.html>
127. Ghazipura M. Community versus inpatient rehabilitation in hip fracture patients: a rapid review (Structured abstract). Health Technology Assessment Database [Internet], 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001045/frame.html>
128. Ghazipura M. Cardiovascular and aerobic exercise in postacute stroke patients: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000494/frame.html>
129. Ghazipura M. Home-Based versus centre-based rehabilitation for community-dwelling postacute stroke patients: an economic rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000493/frame.html>
130. Kabali C. Self-Management and educational interventions in the postacute stroke population: a rapid review (structured Abstract), 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000498/frame.html>
131. Lambrinos A. Preoperative consultation: a rapid review (structured Abstract). health technology assessment database, 2014. Available:

- <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001064/frame.html>
132. Lambrinos A. Medication reconciliation at discharge: a rapid review (structured Abstract). health technology assessment database, 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000507/frame.html>
 133. Lambrinos A. Home-Based exercise programs in heart failure: a rapid review (structured Abstract). health technology assessment database, 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000506/frame.html>
 134. Ling D. Role of screening for respiratory syncytial virus or influenza and empirical antiviral treatment for patients with community-acquired pneumonia: a rapid review (structured Abstract). health technology assessment database, 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001055/frame.html>
 135. McCurdy BR. Inhospital physiotherapy for acute exacerbations of chronic obstructive pulmonary disease (AECOPD): a rapid review (structured Abstract). health technology assessment database, 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001027/frame.html>
 136. McCurdy BR. Action plans for individuals with chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract). health technology assessment database, 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001026/frame.html>
 137. Miani C, Ball S, Pitchforth E, *et al.* Organisational interventions to reduce length of stay in hospital: a rapid evidence assessment (structured Abstract). health technology assessment database, 2014. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001458/frame.html>
 138. Nikitovic M. Intensity of rehabilitation after hip fracture: a rapid review (structured Abstract). health technology assessment database, 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001044/frame.html>
 139. Schaink A. The effectiveness of statins for primary prevention: a rapid review (structured Abstract). health technology assessment database, 2013. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32014001037/frame.html>
 140. Wang M. Airway clearance techniques for patients with stable chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract). health technology assessment database, 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000487/frame.html>
 141. Wang M. Pulmonary rehabilitation in the home versus other settings for individuals with chronic obstructive pulmonary disease (COPD): a rapid review (structured Abstract). health technology assessment database, 2015. Available: <http://onlinelibrary.wiley.com/o/cochrane/clhta/articles/HTA-32015000495/frame.html>
 142. Cottrell E, Whitlock E, Kato E, *et al.* *Defining the benefits of Stakeholder engagement in systematic reviews*. Rockville (MD: AHRQ Methods for Effective Health Care, 2014.
 143. Keown K, Van Eerd D, Irvin E. Stakeholder engagement opportunities in systematic reviews: knowledge transfer for policy and practice. *J Contin Educ Health Prof* 2008;28:67–72.