BMJ Open Association between educational level and total and cause-specific mortality: a pooled analysis of over 694 000 individuals in the Asia Cohort Consortium

Keming Yang,¹ Ying Zhang,² Eiko Saito,³ Md Shafiur Rahman,⁴ Prakash Chandra Gupta,⁵ Norie Sawada,⁶ Akiko Tamakoshi,⁷ Yu-Tang Gao,⁸ Woon-Puay Koh,⁹ Xiao-Ou Shu,¹⁰ Ichiro Tsuji,¹¹ Atsuko Sadakane,¹² Chisato Nagata,¹³ San-Lin You,¹⁴ Jian-Min Yuan,¹⁵ Myung-Hee Shin,¹⁶ Yu Chen,¹⁷ Wen-Harn Pan,¹⁸ Mangesh S Pednekar,⁵ Shoichiro Tsugane,⁶ Hui Cai,¹⁰ Yong-Bing Xiang,⁸ Kotaro Ozasa,¹² Yasutake Tomata,¹¹ Seiki Kanemura,¹¹ Yumi Sugawara,¹¹ Keiko Wada,¹³ Renwei Wang,¹⁵ Yoon-Ok Ahn,¹⁹ Keun-Young Yoo,²⁰ Habibul Ahsan,²¹ Kee Seng Chia,²² Paolo Boffetta,²³ Daehee Kang,¹⁹ John D Potter,²⁴ Manami Inoue,⁶ Wei Zheng,¹⁰ Hongmei Nan¹

To cite: Yang K, Zhang Y, Saito E, *et al*. Association between educational level and total and cause-specific mortality: a pooled analysis of over 694 000 individuals in the Asia Cohort Consortium. *BMJ Open* 2019;**9**:e026225. doi:10.1136/bmjopen-2018-026225

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-026225>).

Received 22 August 2018
Revised 11 March 2019
Accepted 28 June 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Hongmei Nan; hnan@iu.edu

ABSTRACT

Objective To study the association of educational level and risk of death from all causes, cardiovascular disease (CVD) and cancer among Asian populations.

Design A pooled analysis of 15 population-based cohort studies.

Setting and participants 694 434 Asian individuals from 15 prospective cohorts within the Asia Cohort Consortium.

Interventions None.

Main outcome measures HRs and 95% CIs for all-cause mortality, as well as for CVD-specific mortality and cancer-specific mortality.

Results A total of 694 434 participants (mean age at baseline=53.2 years) were included in the analysis. During a mean follow-up period of 12.5 years, 103 023 deaths were observed, among which 33 939 were due to cancer and 34 645 were due to CVD. Higher educational levels were significantly associated with lower risk of death from all causes compared with a low educational level ($\leq$ primary education); HRs and 95% CIs for secondary education, trade/technical education and $\geq$ university education were 0.88 (0.85 to 0.92), 0.81 (0.73 to 0.90) and 0.71 (0.63 to 0.80), respectively ($p_{\text{trend}}=0.002$). Similarly, HRs (95% CIs) were 0.93 (0.89 to 0.97), 0.86 (0.78 to 0.94) and 0.81 (0.73 to 0.89) for cancer death, and 0.88 (0.83 to 0.93), 0.77 (0.66 to 0.91) and 0.67 (0.58 to 0.77) for CVD death with increasing levels of education (both $p_{\text{trend}} < 0.01$). The pattern of the association among East Asians and South Asians was similar compared with $\leq$ primary education; HR (95% CI) for all-cause mortality associated with $\geq$ university education was 0.72 (0.63 to 0.81) among 539 724 East Asians (Chinese, Japanese and Korean) and 0.61 (0.54 to 0.69) among 154 710 South Asians (Indians and Bangladeshis).

Conclusion Higher educational level was associated with substantially lower risk of death among Asian populations.

Strengths and limitations of this study

- This is the first and largest (n=694 434) pooled analysis of Asians to evaluate the impact of educational level on mortality in Asia.
- Our study is a pooling project of 15 prospective cohort studies from multiple countries with long-term follow-up in the Asia Cohort Consortium (ACC).
- This analysis is based on high-quality individual data collected from participating cohorts and harmonised using a standardised process at ACC coordinating centre.
- Our study revealed an inverse association of educational levels and risk of death in Asian populations; in Asia, further studies examining the determinants of educational inequalities in mortality are warranted.

INTRODUCTION

Research has demonstrated that socioeconomic status (SES) has a significant impact on individual health status in terms of mortality, morbidity and disability.^{1 2} Educational attainment is closely related to income, occupation, access to medical care and lifestyle habits. As a major determinant of SES, education has been inversely associated with mortality, an overall measure of health.^{3 4}

Several cohort analyses have examined the relationship between education and mortality in the Western countries. For example, an analysis of two large American Cancer Society cohorts followed from 1959 to 1996 showed that low educational level was associated with higher all-cause death rates.⁵ The follow-up of the US National Longitudinal Mortality Study

(2002–2011) found that both all-cause and cause-specific mortalities were higher among the least-educated than among the most-educated groups, and educational disparities in mortality were more apparent in those aged 50–64 years than those aged 66–79 years.⁶ In the Europe, the inverse association between education with total and cause-specific mortality was also supported by a study of eight western European populations (1990–1997).⁷ Leinsalu *et al*⁸ examined the educational inequalities in mortality in four Eastern European countries (1990–2000); they found mortality rates decreased in similar patterns in all educational groups in Poland and Hungary, whereas in Estonia and Lithuania, mortality rates decreased among the highly educated group but increased among those with low education. Generally, over the past two decades, mortality has declined substantially in lower socioeconomic groups in most European countries; however, relative mortality inequalities widened because the declines were less marked in lower socioeconomic populations.⁹

In Asia, however, relatively few studies have assessed the education/mortality relationship among populations in this area. Further, such investigations were previously conducted only among one single country or area, such as South Korea,¹⁰ the city of Wuhan in China,¹¹ Japan^{12 13} and India.¹⁴ Based on the information from the UNESCO Institute for Statistics (<http://data.uis.unesco.org>), compared with European countries and the USA, mean years of schooling and higher education rates are relatively lower whereas disease and death burdens are relatively higher in Asian countries.^{15–17} Therefore, in this current study, we seek to examine the association of education and mortality in combined Asian populations by using pooled data from 15 prospective cohort studies in the Asia Cohort Consortium (ACC). The large sample size of this pooled analysis provided a strong statistical power to quantify the impact of educational levels on all-cause and major cause-specific mortalities in Asia.

METHODS

Study population

Our study is a pooling project of prospective cohort studies in the ACC, a collaborative effort committed to studying the aetiology of diseases in Asian populations.

Figure 1 Flowchart of Asian individuals eligible for inclusion in the study.

The ACC includes more than 20 cohorts representing Japan, China, South Korea, India, Taiwan, Bangladesh and Singapore. Cohorts were identified through a systematic search of the literature in early 2008, followed by a survey sent to each cohort to assess data availability. Details of the ACC and each participating cohorts have been presented elsewhere.^{18 19} A total of 15 cohorts had collected information about educational attainment and therefore have been included in this pooled analysis. Individual data from participating cohorts was collected and harmonised for the statistical analysis. This study was approved by the ethics committees of each individual cohort study and by the institutional review board of the ACC coordinating centre (Fred Hutchinson Cancer Research Center, Seattle, USA; National Cancer Center, Tokyo, Japan).

Patient and public involvement

Patients or public were not involved in this study.

Data harmonisation

Relevant data from each of the participating cohorts were transferred and harmonised at the ACC coordinating centre. Harmonisation involved several rounds of discussions to ensure that variables were correctly interpreted and extracted. Data were checked for illogical or missing values, and queries were sent back for clarification. The distributions of individual variables were explored to identify false or implausible values. All personal identifiers were removed, but study-specific ID numbers were retained to facilitate referral of all queries to the individual cohort.

Education level and outcome measurements

We excluded from our analysis participants without complete information on educational level (n=31 036), vital status (n=1347), baseline age (n=3947) or follow-up time (n=90). After these exclusions, 694 434 subjects from the 15 participating cohorts were included, among whom 539 724 were East Asians (Chinese (including cohorts from mainland China, Taiwan, Singapore), Japanese and Koreans) and 154 710 were South Asians (Indians and Bangladeshis). Among all included subjects, 344 702 were men and 349 732 were women (figure 1).

Information about completed education was collected by each cohort through questionnaire and harmonised into the following groups: ≤primary education (ie, no formal education or completed only primary education), secondary education (ie, high school education), technical/trade education (ie, associate degree) and ≥university education (ie, undergraduate or graduate education). Data on all-cause and cause-specific mortality were ascertained through linkage to death certificate data or active follow-up. The diagnosis of cause-specific mortality was made according to the International Classification of Diseases, 9th or 10th Revision (ICD-9 or ICD-10): all cancer (ICD-9 codes: 140–208; ICD-10 codes: C00–C97),

cardiovascular diseases (CVDs) (ICD-9 codes: 390–459; ICD-10 codes: I00–I99).

Statistical analysis

Cox proportional hazards regression models were utilised to estimate HRs and 95% CIs for the association between educational level and the risk of death, with '≤primary education' as the reference group for the estimation. The ages of the subjects when they entered and exited the cohort were used to define the time scale in the Cox models. The entry time was defined as age at the baseline interview, and the exit time was defined as age at death or last follow-up, whichever occurred earlier.

We built up two models: the crude model, adjusted only for baseline age and gender; and the multivariable-adjusted model, which was further adjusted for other potential confounders, including residential location (urban, rural), marital status (single, married, widowed/separated/divorced), body mass index (BMI) (underweight, normal, overweight, obese), smoking status (never, ever), alcohol drinking status (never, ever) and physical activity (<1 hour/week, 1–2 hours/week, 3–4 hours/week, ≥5 hours/week). Cox models were performed for each cohort, and random-effects meta-analyses were conducted to summarise results across cohorts. In all models, the impact of educational level on the risk of death was assumed to be cohort-specific.²⁰ We assumed that the log-HRs associated with education level had a fixed-effect component that was common to all cohorts within each country and a random effect that was cohort-specific. The random effects for log-HRs were assumed to be normally distributed, with mean zero; namely, we assumed that β_{ij} , the estimated log-HR for the j th educational level in an i th cohort, follows the distribution $\beta_{ij} \sim N(\beta_j, \sigma^2_{ij} + \tau^2_j)$, where σ^2_{ij} is the within-study variance of β_{ij} and τ^2_j is the between-cohort variance of β_{ij} , as estimated from the Cox regression model.^{21 22}

We also conducted stratified analyses by the covariates described above and assessed statistical significance of interaction using the Wald test for cross-product terms of covariates and education in the Cox models adjusted for other confounding factors. Cox model estimation for each cohort was performed using the PHREG procedure in SAS, V.9.4 (SAS Institute). The meta-analysis estimation was performed using STATA, V.14.0 (StataCorp LP). All tests were two-sided, and $p < 0.05$ was considered statistically significant.

RESULTS

After exclusions, a total of 694 434 participants were included in the 15 participating cohorts; half (49.6%) were women. Nearly half (50.2%) of study participants received no formal education or completed only primary education. Approximately 9% of study participants had received university or higher degrees at baseline. Our analysis included 103 023 deaths during a mean follow-up period of 12.5 years, among which 33 939 (33.0%) were from cancer, 34 645 (33.6%) were from CVD and the remaining 34 439

(33.4%) were from other causes. Selected characteristics of cohorts included in the present study are listed in [table 1](#).

We found inverse associations of education levels with all-cause mortality among our total population as well as populations from different countries or regions ([table 2](#)). When adjusted for age, sex, residential location, marital status, BMI, smoking, alcohol drinking status and physical activity, HRs for mortality were slightly attenuated compared with crude models (age- and sex-adjusted only), but remained statistically significant. No substantial difference was observed between the two models in terms of the pattern of the association. Among all combined cohorts, higher education level was significantly associated with lower risk of death from all causes; compared with ≤primary education, multivariable-adjusted HRs and 95% CIs for secondary education, trade/technical education and ≥university education were 0.88 (0.85 to 0.92), 0.81 (0.73 to 0.90) and 0.71 (0.63 to 0.80), respectively ($p_{\text{trend}} = 0.002$). The pattern of association among East Asians and South Asians appeared similar; regarding all-cause mortality compared with ≤primary education, multivariable-adjusted HRs and 95% CIs for ≥university education was 0.72 (0.63 to 0.81) among 539 724 East Asians (Chinese, Japanese and Korean) and 0.61 (0.54 to 0.69) among 154 710 South Asians (Indians and Bangladeshis).

A similar pattern of association was observed for cause-specific mortality due to CVD and cancer, while the strength of the inverse association was weaker for deaths due to cancer than those due to CVD and all causes ([table 3](#)). Compared with low educational level (ie, ≤primary education), higher educational level was significantly associated with lower cancer-specific mortality (multivariable-adjusted HR (95% CI) for secondary education=0.93 (0.89 to 0.97), for trade/technical education=0.86 (0.78 to 0.94) and for ≥university degree=0.81 (0.73 to 0.89), $p_{\text{trend}} = 0.006$), as well as lower CVD-specific mortality (multivariable-adjusted HR (95% CI) for secondary education=0.88 (0.83 to 0.93), for trade/technical education=0.77 (0.66 to 0.91) and for ≥university degree=0.67 (0.58 to 0.77), $p_{\text{trend}} = 0.002$). As shown in [table 3](#), the significant gradient of the inverse association between education and cancer/CVD-specific mortality also appeared in the East Asian population-only analysis ($p < 0.05$). In South Asians, compared with the reference group (ie, ≤primary education), individuals with ≥university education had significantly lower risk of death from CVD ($p < 0.05$). The association between education and cancer-specific mortality was not statistically significant among South Asians probably due to the small numbers of deaths across different educational groups among South Asians.

We further examined the association between educational levels and risk of death due to all causes stratified by covariates (online supplementary table 1). We observed statistically significant effect modifications by age, gender, residential location, marital status, physical activity, smoking and alcohol consumption status on the association between education and all-cause mortality (p for interactions < 0.01). However, for all covariates except

Table 1 Selected characteristics of participating cohorts

Cohort	No of subjects	Dates of enrolment	Mean follow-up years	Women (%)	Mean age (SD) at entry	Median (range) of age at entry	Education level				All causes		Cause of death (%)		
							≤ Primary (N (%))	Secondary (N (%))	Trade/technical (N (%))	≥University (N (%))	# death (mortality%)	CVD (N (%))	Cancer (N (%))	Other (N (%))	
Mainland China															
SMHS	60611	2002–2006	9.4	0	55.3 (9.7)	53 (40–75)	4084 (6.7)	20334 (33.6)	21858 (36.1)	14335 (23.7)	5364 (8.9)	1772 (33.0)	2328 (43.4)	1264 (23.6)	
SWHS	74925	1996–2000	14.9	100	52.6 (9.1)	50 (40–71)	16181 (21.6)	27682 (37.0)	20890 (27.9)	10172 (13.6)	7651 (10.2)	2490 (32.5)	3219 (42.1)	1942 (25.4)	
SCS	18100	1986–1989	16.3	0	55.3 (5.7)	56 (35–76)	5156 (28.5)	8472 (46.8)	1067 (5.9)	3405 (18.8)	4983 (27.5)	1686 (33.8)	1974 (39.6)	1323 (26.6)	
Taiwan															
CBCSP	23810	1991–1992	15.2	49.7	47.3 (10.0)	48(29–66)	14998 (63.0)	3279 (13.8)	3560 (15.0)	1973 (8.3)	2767 (11.6)	558 (20.2)	1013 (36.6)	1196 (43.2)	
CVDFACTS	5142	1990–1993	14.9	55.9	47.5 (15.6)	48 (18–92)	2366 (46.0)	2031 (39.5)	–	745 (14.5)	825 (16.0)	220 (26.7)	218 (26.4)	387 (46.9)	
Singapore															
SCHS	63247	1993–1999	11.5	55.8	56.5 (8.0)	56 (43–83)	45379 (71.8)	14621 (23.1)	2223 (3.5)	1024 (1.6)	10682 (16.9)	3708 (34.7)	3887 (36.4)	3087 (28.9)	
Japan															
JACC	69951	1988–1990	12.5	58.6	57.4 (10.0)	58 (40–79)	41978 (60.0)	18724 (26.8)	6918 (9.9)	2331 (3.3)	9717 (13.9)	3011 (31.0)	3610 (37.2)	3096 (31.9)	
JPHC	40692	1990–1992	21	52.1	49.5 (5.9)	50 (40–59)	21248 (52.2)	14761 (36.3)	2752 (6.8)	1931 (4.7)	6608 (16.2)	1738 (26.3)	2744 (41.5)	2126 (32.2)	
Miyagi	43525	1990	16.2	52.1	52.0 (7.6)	52 (40–64)	26101 (60.0)	11651 (26.8)	4147 (9.5)	1626 (3.7)	5086 (11.7)	1376 (27.1)	2359 (46.4)	1351 (26.6)	
Ohsaki	46990	1995	10.8	51.6	60.1 (10.3)	61 (40–80)	28387 (60.4)	15060 (32.1)	2601 (5.5)	942 (2.0)	7517 (16.0)	2422 (32.2)	2654 (35.3)	2441 (32.5)	
Takayama	31106	1992	13.6	54.2	55.9 (12.8)	55 (35–101)	19730 (63.4)	9013 (29.0)	–	2363 (7.6)	6076 (19.5)	2026 (33.3)	1771 (29.2)	2279 (37.5)	
LSS	47928	1963–1993	22	59.9	52.1 (13.6)	51 (19–99)	21889 (45.7)	20605 (43.0)	–	5434 (11.3)	24818 (51.8)	9109 (36.7)	6734 (27.1)	8975 (36.2)	
Republic of Korea															
SeoulIM	13697	1992–1993	15.6	0	49.2 (5.2)	49 (25–82)	133 (1.0)	6227 (45.5)	–	7337 (53.6)	894 (6.5)	154 (17.2)	493 (55.2)	247 (27.6)	
India															
Mumbai	142973	1991–1997	5.2	40.9	50.5 (11.1)	49 (35–98)	92639 (64.8)	43908 (30.7)	–	6426 (4.5)	9147 (6.4)	4008 (43.8)	797 (8.7)	4342 (47.5)	
Bangladesh															
HEALS	11737	2000–2002	12.3	57.1	37.1 (10.1)	36 (17–75)	8705 (74.2)	2915 (24.8)	117 (1.0)	–	888 (7.6)	367 (41.3)	138 (15.5)	383 (43.1)	
Total	694434	1963–2006	12.5	49.6	53.2 (10.7)	52 (17–101)	348974 (50.2)	219283 (31.6)	66133 (9.5)	60044 (8.7)	103023 (14.8)	34645 (33.6)	33939 (33.0)	34439 (33.4)	

Participants without complete information on educational level, vital status, baseline age or follow-up time were excluded; among all participants, 12 people aged 17 years and 2 people aged 101 years.

Participants who died from unknown causes were excluded from analysis.

CBCSP, Community-based Cancer Screening Project; CVDFACTS, Cardiovascular Disease risk FACTor Two-township Study; HEALS, Health Effects for Arsenic Longitudinal Study Bangladesh; JACC, Japan Collaborative Cohort Study; JPHC, Japan Public Health Center-based prospective Study; LSS, Life Span Study Cohort; Miyagi, Miyagi Cohort; Mumbai, Mumbai Cohort Study; Ohsaki, Ohsaki National Health Insurance Cohort Study; SCHS, Singapore Chinese Health Study; SCS, Shanghai Cohort Study; SMHS, Shanghai Men's Health Study; SWHS, Shanghai Women's Health Study; SeoulIM, Seoul Male Cancer Cohort; Takayama, Takayama Study.

Table 2 Association of educational level with risk of death from all causes in selected study populations in Asia

Population	Number		Model 1	Model 2
	Participants	Deaths	HR (95% CI)	HR (95% CI)
All cohorts combined				
≤Primary education	348 974	64 948	1	1
Secondary education	219 283	26 652	0.86 (0.82 to 0.90)	0.88 (0.85 to 0.92)
Trade/technical education	66 133	5195	0.76 (0.68 to 0.86)	0.81 (0.73 to 0.90)
≥University education	60 044	6228	0.66 (0.57 to 0.77)	0.71 (0.63 to 0.80)
P for trend			0.002	0.002
Mainland China				
≤Primary education	25 421	7227	1	1
Secondary education	56 488	5882	0.79 (0.76 to 0.83)	0.82 (0.79 to 0.85)
Trade/technical education	43 815	2702	0.64 (0.60 to 0.69)	0.69 (0.66 to 0.73)
≥University education	27 912	2187	0.50 (0.48 to 0.54)	0.56 (0.53 to 0.59)
P for trend			<0.001	<0.001
Taiwan				
≤Primary education	17 364	2777	1	1
Secondary education	5310	478	0.83 (0.75 to 0.92)	0.85 (0.70 to 1.02)
Trade/technical education	3560	202	0.75 (0.64 to 0.87)	0.82 (0.70 to 0.95)
≥University education	2718	135	0.50 (0.42 to 0.60)	0.57 (0.48 to 0.68)
P for trend			0.03	0.08
Singapore				
≤Primary education	45 379	8810	1	1
Secondary education	14 621	1596	0.76 (0.72 to 0.81)	0.81 (0.77 to 0.86)
Trade/technical education	2223	205	0.67 (0.58 to 0.76)	0.75 (0.65 to 0.86)
≥University education	1024	71	0.52 (0.41 to 0.66)	0.63 (0.50 to 0.80)
P for trend			0.002	0.03
Japan				
≤Primary education	159 333	38 765	1	1
Secondary education	89 814	15 833	0.92 (0.87 to 0.98)	0.94 (0.88 to 0.99)
Trade/technical education	16 418	2075	0.90 (0.81 to 1.00)	0.92 (0.82 to 1.02)
≥University education	14 627	3149	0.89 (0.80 to 0.99)	0.89 (0.80 to 0.99)
P for trend			0.47	0.36
Republic of Korea				
≤Primary education	133	17	1	1
Secondary education	6227	480	0.75 (0.46 to 1.22)	0.77 (0.47 to 1.26)
Trade/technical education	–	–	–	–
≥University education	7337	397	0.48 (0.30 to 0.78)	0.52 (0.32 to 0.86)
P for trend			<0.001	<0.001
India				
≤Primary education	92 639	6663	1	1
Secondary education	43 908	2195	0.83 (0.79 to 0.87)	0.89 (0.85 to 0.94)
Trade/technical education	–	–	–	–
≥University education	6426	289	0.55 (0.48 to 0.61)	0.61 (0.54 to 0.69)
P for trend			<0.001	<0.001
Bangladesh				
≤Primary education	8705	689	1	1

Continued

Table 2 Continued

Population	Number		Model 1	Model 2
	Participants	Deaths	HR (95% CI)	HR (95% CI)
Secondary education	2915	188	0.83 (0.70 to 0.98)	0.92 (0.77 to 1.08)
Trade/technical education	117	11	0.92 (0.51 to 1.68)	1.15 (0.63 to 2.10)
≥University education	–	–	–	–
P for trend			0.73	0.47
East Asians				
≤Primary education	247 630	57 596	1	1
Secondary education	172 460	24 269	0.86 (0.82 to 0.91)	0.88 (0.84 to 0.93)
Trade/technical education	66 016	5184	0.76 (0.67 to 0.86)	0.80 (0.72 to 0.89)
≥University education	53 618	5939	0.70 (0.57 to 0.79)	0.72 (0.63 to 0.81)
P for trend			0.008	0.006
South Asians				
≤Primary education	101 344	7352	1	1
Secondary education	46 823	2383	0.83 (0.79 to 0.87)	0.89 (0.85 to 0.94)
Trade/technical education	117	11	0.92 (0.51 to 1.68)	1.15 (0.63 to 2.10)
≥University education	6426	289	0.55 (0.48 to 0.61)	0.61 (0.54 to 0.69)
P for trend			0.02	0.03

Model 1: adjust for baseline age and sex.

Model 2: adjust for baseline age, sex, urban/rural residence, marital status, body mass index, smoking status, alcohol consumption status and physical activity.

East Asians include participants from mainland China, Taiwan, Singapore, Republic of Korea and Japan; South Asians include participants from India and Bangladesh.

residential location and marital status, we did not find material differences among stratified groups in terms of the risk estimates of education on total mortality, although *p* value for interaction was statistically significant.

DISCUSSION

In this large (694 434) pooled analysis of Asian populations, we found that higher educational level was associated with a substantially reduced risk of death from all causes, CVD and cancer. This inverse association presented a dose–response pattern. The mortality disparity did not appear to be explained solely by lifestyle factors such as the level of physical activity, smoking and alcohol consumption. Our study provides convincing evidence for the link between education and risk of death in Asian populations.

Our findings in Asian populations are generally similar to the results in studies carried out in Western countries. Kitagawa and Hauser, using 1960 US death records and census data, first examined the SES/mortality relationship. They found not only that both men and women with higher education have lower mortality rates and live longer than those with lower education but also that there is a significant gradient across the seven ordered categories of completed schooling.²³ An additional 4 years of education lowered 5-year mortality by 1.8%, according to the US National Bureau of Economic Research.²⁴ A low level of education was also found to be associated with

increased all-cause mortality compared with high levels of education in cohorts with type II diabetes^{25–27} and acute coronary syndrome.²⁸ Recently, Mackenbach *et al* studied the trends in health inequalities in 27 European countries (1980–2014); they found that, in Western Europe, all-cause mortality has declined steadily in both low-educated and highly educated men and women; the trend in mortality was generally stable in both education groups.²⁹ In Western Europe, absolute inequalities have usually decreased due to the larger absolute declines among the low-educated groups, but relative inequalities have generally increased because relative mortality declines were larger in the highly educated.²⁹ However, since the 1990s, relative and absolute inequalities in mortality have both increased in Eastern Europe.^{8,9} The recent work by Mackenbach *et al* found that mortality has also begun to decline among the low-educated population in Eastern European countries; absolute inequalities in mortality have started to decrease as well.²⁹

The inverse association between education and mortality was also found among older people in low-income and middle-income countries by a population-based cohort study of 12 373 people aged ≥65 years from Latin America, China and India.¹⁵ In 2014, Vathesatogkit *et al* performed and published the first meta-analysis of studies from Asia on the association between SES and mortality.³⁰ Consistent with our findings, they found overall that those with the lowest level of education experienced a 1.4-fold higher risk

Table 3 Association of educational level with risk of death from cardiovascular diseases and cancers in selected study populations in Asia

Population	CVD			Cancer		
	Number of deaths	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Number of deaths	Model 1 HR (95% CI)	Model 2 HR (95% CI)
All cohorts combined						
≤Primary education	22 723	1	1	19 465	1	1
Secondary education	8559	0.86 (0.81 to 0.91)	0.88 (0.83 to 0.93)	9626	0.90 (0.86 to 0.95)	0.93 (0.89 to 0.97)
Trade/technical education	1478	0.73 (0.61 to 0.88)	0.77 (0.66 to 0.91)	2322	0.81 (0.72 to 0.91)	0.86 (0.78 to 0.94)
≥University education	1885	0.63 (0.53 to 0.75)	0.67 (0.58 to 0.77)	2526	0.75 (0.64 to 0.86)	0.81 (0.73 to 0.89)
P for trend		0.003	0.002		0.01	0.006
Mainland China						
≤Primary education	2700	1	1	2548	1	1
Secondary education	1858	0.80 (0.75 to 0.85)	0.83 (0.78 to 0.89)	2612	0.84 (0.74 to 0.95)	0.88 (0.79 to 0.97)
Trade/technical education	729	0.56 (0.48 to 0.66)	0.61 (0.51 to 0.73)	1334	0.72 (0.57 to 0.92)	0.79 (0.65 to 0.96)
≥University education	661	0.47 (0.42 to 0.51)	0.52 (0.45 to 0.60)	1027	0.58 (0.45 to 0.75)	0.66 (0.57 to 0.78)
P for trend		<0.001	0.001		0.03	0.03
Taiwan						
≤Primary education	619	1	1	931	1	1
Secondary education	97	0.76 (0.55 to 1.06)	0.78 (0.53 to 1.15)	169	0.88 (0.73 to 1.06)	0.91 (0.77 to 1.09)
Trade/technical education	33	0.71 (0.50 to 1.03)	0.77 (0.53 to 1.11)	81	0.78 (0.62 to 1.00)	0.84 (0.65 to 1.07)
≥University education	29	0.52 (0.36 to 0.77)	0.61 (0.42 to 0.90)	50	0.54 (0.40 to 0.73)	0.63 (0.47 to 0.84)
P for trend		0.20	0.39		0.08	0.13
Singapore						
≤Primary education	3079	1	1	3150	1	1
Secondary education	536	0.77 (0.70 to 0.84)	0.81 (0.74 to 0.90)	627	0.78 (0.71 to 0.85)	0.84 (0.77 to 0.92)
Trade/technical education	80	0.79 (0.63 to 0.99)	0.87 (0.69 to 1.08)	75	0.62 (0.49 to 0.78)	0.71 (0.57 to 0.90)
≥University education	13	0.30 (0.17 to 0.51)	0.35 (0.20 to 0.60)	35	0.64 (0.45 to 0.89)	0.80 (0.57 to 1.12)
P for trend		<0.001	0.002		0.24	0.77
Japan						
≤Primary education	13 187	1	1	12 159	1	1
Secondary education	4910	0.89 (0.81 to 0.97)	0.91 (0.83 to 0.99)	5736	0.95 (0.92 to 0.99)	0.96 (0.93 to 1.00)
Trade/technical education	632	0.90 (0.78 to 1.04)	0.92 (0.80 to 1.06)	830	0.93 (0.84 to 1.02)	0.94 (0.85 to 1.04)
≥University education	953	0.84 (0.72 to 0.96)	0.83 (0.73 to 0.95)	1147	0.89 (0.84 to 0.95)	0.89 (0.84 to 0.95)
P for trend		0.49	0.38		0.10	0.09
Republic of Korea						
≤Primary education	4	1	1	7	1	1
Secondary education	74	0.51 (0.19 to 1.40)	0.58 (0.21 to 1.60)	252	0.97 (0.46 to 2.06)	0.99 (0.46 to 2.11)
Trade/technical education	–	–	–	–	–	–
≥University education	76	0.40 (0.15 to 1.10)	0.51 (0.18 to 1.41)	234	0.69 (0.33 to 1.47)	0.74 (0.35 to 1.59)
P for trend		0.14	0.43		<0.001	0.002
India						
≤Primary education	2868	1	1	564	1	1
Secondary education	987	0.89 (0.83 to 0.97)	0.90 (0.84 to 0.98)	200	0.88 (0.74 to 1.04)	0.99 (0.84 to 1.18)

Continued

Table 3 Continued

Population	CVD			Cancer		
	Number of deaths	Model 1 HR (95% CI)	Model 2 HR (95% CI)	Number of deaths	Model 1 HR (95% CI)	Model 2 HR (95% CI)
Trade/technical education	–	–	–	–	–	–
≥University education	153	0.67 (0.57 to 0.79)	0.68 (0.58 to 0.80)	33	0.77 (0.54 to 1.10)	0.92 (0.65 to 1.32)
P for trend		<0.001	<0.001		0.48	0.70
Bangladesh						
≤Primary education	266	1	1	106	1	1
Secondary education	97	1.11 (0.88 to 1.41)	1.08 (0.84 to 1.38)	30	0.84 (0.56 to 1.26)	0.97 (0.63 to 1.47)
Trade/technical education	4	0.88 (0.33 to 2.38)	0.88 (0.32 to 2.38)	2	1.04 (0.26 to 4.24)	1.38 (0.33 to 5.69)
≥University education	–	–	–	–	–	–
P for trend		0.65	0.68		0.76	0.63
East Asians						
≤Primary education	19589	1	1	18795	1	1
Secondary education	7475	0.84 (0.79 to 0.90)	0.87 (0.82 to 0.92)	9396	0.90 (0.86 to 0.95)	0.93 (0.89 to 0.97)
Trade/technical education	1474	0.73 (0.61 to 0.87)	0.77 (0.66 to 0.91)	2320	0.81 (0.72 to 0.91)	0.86 (0.78 to 0.94)
≥University education	1732	0.62 (0.51 to 0.75)	0.67 (0.57 to 0.78)	2493	0.74 (0.64 to 0.87)	0.80 (0.72 to 0.89)
P for trend		0.01	0.01		0.02	0.01
South Asians						
≤Primary education	3134	1	1	670	1	1
Secondary education	1084	0.97 (0.79 to 1.19)	0.95 (0.81 to 1.11)	230	0.87 (0.75 to 1.02)	0.99 (0.84 to 1.16)
Trade/technical education	4	0.88 (0.33 to 2.38)	0.88 (0.32 to 2.38)	2	1.04 (0.26 to 4.24)	1.38 (0.33 to 5.69)
≥University education	153	0.67 (0.57 to 0.79)	0.68 (0.58 to 0.80)	33	0.77 (0.54 to 1.10)	0.92 (0.65 to 1.32)
P for trend		0.17	0.14		0.61	0.80

Model 1: adjust for baseline age and sex.

Model 2: adjust for baseline age, sex, urban/rural residence, marital status, body mass index, smoking status, alcohol consumption status and physical activity.

East Asians include participants from mainland China, Taiwan, Singapore, Republic of Korea and Japan; South Asians include participants from India and Bangladesh.

CVD, cardiovascular disease.

of mortality than the highest level of education (all cause: RR=1.41, 95% CI=1.29 to 1.52; CVD: RR=1.66, 95% CI=1.23 to 2.25; cancer: RR=1.16, 95% CI=1.07 to 1.27). However, we note that—unlike our pooled analysis—due to the nature of a meta-analysis based only on reported estimates from previous publications, the detailed educational levels in the comparison groups were not specified in the study by Vathesatogkit *et al.*³⁰

Educational attainment may not have a strong causal relationship with adult mortality. One nationwide quasi-experiment was conducted to examine the causal effect of education on mortality among 1.2 million Swedes;³¹ the exposure was a 1-year increase in compulsory schooling as an educational reform implemented in Sweden from 1949 to 1962. No significant difference in all-cause mortality between the experimental and control groups was found during the entire follow-up (HR=0.98, 95% CI=0.95 to 1.01) or among those aged ≤40 years (HR=1.03, 95% CI=0.98 to

1.07). Risk of death from all causes in the experimental group aged >40 years was lower than that in the control group with marginally significant association (HR=0.96, 95% CI=0.93 to 0.99).³¹ Interestingly, utilising a longitudinal dataset of Danish twins, Behrman *et al* found only weak evidence of an association between educational level and adult mortality among identical twin pairs who shared both the same genetic background and similar childhood social environments; but stronger evidence of the education/mortality relationship was found among fraternal twin pairs and unrelated individuals.³²

Understanding the mechanisms by which educational attainment influences death risk is highly important for public policy making. There are several possible explanations. On one hand, education may affect health outcomes via people's SES (eg, occupation, income, wealth), social resources (eg, access to health information and healthcare services), health-related behaviours

(eg, smoking, alcohol consumption, dietary habits, physical activity) and cognitive skills (eg, communication with physicians and nurses). On the other hand, a person's educational attainment can also act as a surrogate measurement for early-life factors such as parental SES, physical and mental health, childhood environment and social context^{4 33}; in other words, education could lie on the pathway between those early-life factors and health outcomes (eg, mortality). Education seems to have an important latent effect on mortality into late life, suggesting that the adverse effect of low educational attainment and socioeconomic sequelae may accumulate across the life course.³⁴

A study of 17 European countries (1970–2010) indicated that behavioural factors, including smoking and excessive alcohol consumption, were important contributors to the between-country variations in the magnitude of socioeconomic inequalities in mortality.³⁵ By comparing estimates (ie, HRs) obtained from the crude model (age- and sex-adjusted only) and the model further adjusted for each modifiable covariate individually, we found that smoking explained the most of the education–mortality association in our pooled data. Tobacco smoking is associated with a substantially increased risk of death among Asian adults, accounting for approximately 2 million deaths in adults aged over 45 years throughout Asia in 2004.¹⁸ Research has shown that smoking rates were higher among lower-educated people in many countries and the educational differences in smoking were more apparent in younger population than older generations.^{36–38} In our current study, the inverse associations between education and risk of death are significant in both never-smokers and ever-smokers; the magnitude of the association is slightly larger among those who never smoked (online supplementary table 1). In a study of 14 European countries, Gregoraci *et al* found that smoking-attributable mortality was inversely associated with socioeconomic levels defined by education and occupation in 2000–2004.³⁹ Though the contribution of smoking to socioeconomic inequalities in mortality has been reduced since 1990–2004, smoking remains one of the most important intervention targets for reducing health inequalities in Europe.³⁹ In Asian populations, further studies examining the determinants of socioeconomic inequalities in mortality are warranted.

Our study has several strengths. First, this pooling project is the first and largest study evaluating the education/mortality relationship using combined Asian populations. Second, in contrast with a meta-analysis based on data from previous publications, our pooled analysis provides more accurate and reliable estimates by directly utilising individual data with standardised exposure measurement. Besides, the detailed covariate information collected in these cohorts enables a careful control for potential confounding and evaluation of effect modifications. We also acknowledge some limitations. First, because educational attainment data are self-reported at baseline in each cohort, we cannot rule out the possibility of misclassification. Additionally, there may be

heterogeneity in education level across cohorts. However, we used a standardised harmonisation process on a range of environmental variables, including education level across 15 participating cohorts. Results presented in tables 2 and 3 show the consistency of the association between educational level and risk of death on a per-study level, and the pooled risk estimate is similar to that found in prior studies.^{5 27} Thus, bias due to self-report and heterogeneity in educational level is likely to be minimal. Second, we also cannot completely rule out the possibility that some participants might have had further education during follow-up after baseline enrolment (mean age at baseline=53.2 years). However, in Asian countries, pursuit of further education among older adults is not as popular as in Western countries. Thus, such missing information seems less likely to affect our findings.

CONCLUSION

Despite the limitations mentioned above, our study provides the best estimate of the impact of education on all-cause as well as major cause-specific mortality in Asian populations to date. Our findings may contribute to better public policy decisions, especially regarding increasing educational opportunities for Asian populations as a powerful intervention to reduce both morbidity and mortality.

Author affiliations

¹Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, Indianapolis, Indiana, USA

²Department of Biostatistics, College of Public Health, University of Nebraska Medical Center, Omaha, Nebraska, USA

³Division of Cancer Statistics Integration, Center for Cancer Control and Information Services, National Cancer Center, Tokyo, Japan

⁴Department of Global Health Policy, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

⁵Healis - Sekhsaria Institute for Public Health, Navi Mumbai, India

⁶Epidemiology and Prevention Group, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan

⁷Faculty of Medicine, Hokkaido University, Sapporo, Japan

⁸State Key Laboratory of Oncogene and Related Genes & Department of Epidemiology, Shanghai Cancer Institute, Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

⁹Health Services and Systems Research, Duke-NUS Medical School, Singapore

¹⁰Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, Tennessee, USA

¹¹Tohoku University Graduate School of Medicine, Sendai, Japan

¹²Department of Epidemiology, Radiation Effects Research Foundation, Hiroshima, Japan

¹³Department of Epidemiology and Preventive Medicine, Gifu University Graduate School of Medicine, Gifu, Japan

¹⁴School of Medicine, Big Data Research Centre, Fu-Jen Catholic University, Taipei, Taiwan

¹⁵Division of Cancer Control and Population Sciences, UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

¹⁶Department of Social and Preventive Medicine, Sungkyunkwan University School of Medicine, Seoul, The Republic of Korea

¹⁷Department of Population Health, Department of Environmental Medicine, New York University School of Medicine, New York City, New York, USA

¹⁸Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan

¹⁹Department of Preventive Medicine, Seoul National University College of Medicine, Seoul, The Republic of Korea

²⁰Armed Forces Capital Hospital, Seoul National University College of Medicine, Seoul, The Republic of Korea

²¹Department of Public Health Sciences, The University of Chicago, Chicago, Illinois, USA

²²Epidemiology Program, Saw Swee Hock School of Public Health, National University of Singapore, Singapore

²³Icahn School of Medicine at Mount Sinai, New York, New York, USA

²⁴Centre for Public Health Research, Massey University, Wellington, New Zealand

Acknowledgements The authors would like to thank all research team members and participants of each cohort study for their contribution to this research.

Contributors HN was responsible for the study concept and design. PCG, NS, AT, Y-TG, W-PK, X-OS, IT, AS, CN, S-LY, J-MY, M-HS, YC, W-HP, MSP, ST, HC, Y-BX, KO, YT, SK, YS, KW, RW, Y-OA, K-YY, HA, KSC, PB, DK, JDP, MI and WZ were involved in data collection and contributed to study materials. ES and MSR were involved in data management. KY analyzed the data and drafted the manuscript. All authors (KY, YZ, ES, MSR, PCG, NS, AT, Y-TG, W-PK, X-OS, IT, AS, CN, S-LY, J-MY, M-HS, YC, W-HP, MSP, ST, HC, Y-BX, KO, YT, SK, YS, KW, RW, Y-OA, K-YY, HA, KSC, PB, DK, JDP, MI, WZ, HN) contributed to the interpretation of the results and revision of the manuscript critically for important intellectual content. KY and HN are guarantors. All authors, external and internal, had full access to all of the data (including statistical reports and tables) in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding There is no funding for this specific analysis. The cohorts participating in the pooled analysis were supported by the following grants: Japan Public Health Center-based prospective Study (JPHC Study), National Cancer Center Research and Development Fund (23-A-31(toku) and 26-A-2) (since 2011) and a Grant-in-Aid for Cancer Research from the Ministry of Health, Labour and Welfare of Japan (from 1989 to 2010); Japan Collaborative Cohort Study (JACC), National Cancer Center Research and Development Fund, A Grant-in-Aid for Cancer Research; Grant for Health Services and Grant for Comprehensive Research on Cardiovascular and Life-Style Related Diseases from the Ministry of Health, Labour and Welfare, Japan; Grant for the Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan; Shanghai Women's Health Study (SWHS), the US National Cancer Institute (grant numbers R37 CA070867 and UM1 CA182910); Singapore Chinese Health Study (SCHS), National Institutes of Health (R01CA144034, UM1CA182876); Shanghai Men's Health Study (SMHS), the US National Cancer Institute (R01 CA082729 and UM1 CA173640); Shanghai Cohort Study (SCS), National Institutes of Health (R01CA144034, UM1CA182876); Ohsaki National Health Insurance Cohort Study, Grants-in-Aid for Cancer Research and for the Third Term Comprehensive Ten-Year Strategy for Cancer Control (H21-3jigan-ippan-003), Ministry of Health, Labour and Welfare; Miyagi Cohort, Grants-in-Aid for Cancer Research and for the Third Term Comprehensive Ten-Year Strategy for Cancer Control (H21-3jigan-ippan-003), Ministry of Health, Labour and Welfare; Life Span Study Cohort, The Japanese Ministry of Health, Labour and Welfare and the US Department of Energy; Takayama Study, National Cancer Center Research and Development Fund; CardioVascular Disease risk FACTor Two-township Study (CVDFACTS), Department of Health, Taiwan (DOH80-27, DOH81-021, DOH8202-1027, DOH83-TD-015 and DOH84-TD-006); Community-Based Cancer Screening Program (CBCSP), National Science Council and Department of Health, Taiwan; Seoul Male Cancer Cohort, National R&D Program for Cancer Control, Ministry of Health and Welfare, Republic of Korea (0520160-1); Health Effects of Arsenic Longitudinal Study (HEALS), National Institutes of Health (P42ES010349, R01CA102484, R01CA107431); Mumbai Cohort Study, International Agency for Research on Cancer, Lyon, France; Clinical Trials Service Unit, Oxford, UK; World Health Organisation, Geneva, Switzerland. The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval This study was approved by the ethics committees of each individual cohort study and by the institutional review board of the ACC coordinating center (Fred Hutchinson Cancer Research Center, Seattle, USA; National Cancer Center, Tokyo, Japan).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement The datasets analysed during the current study (deidentified participant data) are available from the corresponding author upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Adler NE, Newman K. Socioeconomic disparities in health: pathways and policies. *Health Aff* 2002;21:60-76.
- Mackenbach JP, Stirbu I, Roskam A-JR, et al. Socioeconomic inequalities in health in 22 European countries. *N Engl J Med Overseas Ed* 2008;358:2468-81.
- Samir KC, Lentzner H. The effect of education on adult mortality and disability: a global perspective Vienna Yearbook of Population Research; 2010: 201-35.
- Hummer RA, Hernandez EM. The effect of educational attainment on adult mortality in the United States. *Popul Bull* 2013;68:1-16.
- Steenland K, Henley J, Thun M. All-Cause and cause-specific death rates by educational status for two million people in two American cancer Society cohorts, 1959-1996. *Am J Epidemiol* 2002;156:11-21.
- Ma J, Altekruse S, Cosgrove C, et al. Educational disparities in mortality between adults aged 50-64 and 66-79 years, U.S. *Am J Prev Med* 2017;52:728-34.
- Huisman M, Kunst AE, Bopp M, et al. Educational inequalities in cause-specific mortality in middle-aged and older men and women in eight Western European populations. *The Lancet* 2005;365:493-500.
- Leinsalu M, Stirbu I, Vågerö D, et al. Educational inequalities in mortality in four eastern European countries: divergence in trends during the post-communist transition from 1990 to 2000. *Int J Epidemiol* 2009;38:512-25.
- Mackenbach JP, Kulháňová I, Artnik B, et al. Changes in mortality inequalities over two decades: register based study of European countries. *BMJ* 2016;353.
- Khang Y-H, Lynch JW, Kaplan GA. Health inequalities in Korea: age- and sex-specific educational differences in the 10 leading causes of death. *Int J Epidemiol* 2004;33:299-308.
- Liang J, McCarthy JF, Jain A, et al. Socioeconomic gradient in old age mortality in Wuhan, China. *J Gerontol B Psychol Sci Soc Sci* 2000;55:S222-S233.
- Fujino Y, Tamakoshi A, Iso H, et al. A nationwide cohort study of educational background and major causes of death among the elderly population in Japan. *Prev Med* 2005;40:444-51.
- Ito S, Takachi R, Inoue M, et al. Education in relation to incidence of and mortality from cancer and cardiovascular disease in Japan. *Eur J Public Health* 2008;18:466-72.
- Pednekar MS, Gupta R, Gupta PC. Illiteracy, low educational status, and cardiovascular mortality in India. *BMC Public Health* 2011;11:567.
- Ferri CP, Acosta D, Guerra M, et al. Socioeconomic factors and all cause and cause-specific mortality among older people in Latin America, India, and China: a population-based cohort study. *PLoS Med* 2012;9:e1001179.
- UNESCO. The UNESCO Institute for statistics. Available: <http://data.uis.unesco.org> [Accessed 15 Feb 2019].
- Ryan CL, Bauman K. Educational attainment in the United States: 2015. U.S. census bureau. Available: <https://www.census.gov/content/dam/Census/library/publications/2016/demo/p20-578.pdf> [Accessed 22 Feb 2019].
- Zheng W, McLerran DF, Rolland BA, et al. Burden of total and cause-specific mortality related to tobacco smoking among adults aged ≥ 45 years in Asia: a pooled analysis of 21 cohorts. *PLoS Med* 2014;11:e1001631.
- Zheng W, McLerran DF, Rolland B, et al. Association between body-mass index and risk of death in more than 1 million Asians. *N Engl J Med* 2011;364:719-29.
- Smith CT, Williamson PR, Marson AG. An overview of methods and empirical comparison of aggregate data and individual patient data results for investigating heterogeneity in meta-analysis of time-to-event outcomes. *J Eval Clin Pract* 2005;11:468-78.
- Brockwell SE, Gordon IR. A comparison of statistical methods for meta-analysis. *Stat Med* 2001;20:825-40.
- DerSimonian R, Laird N. Meta-Analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
- Kitagawa EM, Hauser PM. *Differential mortality in the United States: a study in socioeconomic epidemiology*. Cambridge, Massachusetts: Harvard University Press, 1973.

24. National Bureau of Economic Research. The effects of education on health. Available: <http://www.nber.org/digest/mar07/w12352.html> [Accessed 8 Aug 2017].
25. Blomster JI, Zoungas S, Woodward M, *et al.* The impact of level of education on vascular events and mortality in patients with type 2 diabetes mellitus: results from the advance study. *Diabetes Res Clin Pract* 2017;127:212–7.
26. Saydah SH, Imperatore G, Beckles GL. Socioeconomic status and mortality: contribution of health care access and psychological distress among U.S. adults with diagnosed diabetes. *Diabetes Care* 2013;36:49–55.
27. Rawshani A, Svensson A-M, Zethelius B, *et al.* Association between socioeconomic status and mortality, cardiovascular disease, and cancer in patients with type 2 diabetes. *JAMA Intern Med* 2016;176:1146–54.
28. Notara V, Panagiotakos DB, Kogias Y, *et al.* The impact of educational status on 10-year (2004–2014) cardiovascular disease prognosis and all-cause mortality among acute coronary syndrome patients in the Greek acute coronary syndrome (GREECS) longitudinal study. *J Prev Med Public Health* 2016;49:220–9.
29. Mackenbach JP, Valverde JR, Artnik B, *et al.* Trends in health inequalities in 27 European countries. *Proc Natl Acad Sci U S A* 2018;115:6440–5.
30. Vathesatogkit P, Batty GD, Woodward M. Socioeconomic disadvantage and disease-specific mortality in Asia: systematic review with meta-analysis of population-based cohort studies. *J Epidemiol Community Health* 2014;68:375–83.
31. Lager ACJ, Torssander J. Causal effect of education on mortality in a quasi-experiment on 1.2 million Swedes. *Proc Natl Acad Sci U S A* 2012;109:8461–6.
32. Behrman JR, Kohler H-P, Jensen VM, *et al.* Does more schooling reduce hospitalization and delay mortality? new evidence based on Danish twins. *Demography* 2011;48:1347–75.
33. Hayward MD, Hummer RA, Sasson I. Trends and group differences in the association between educational attainment and U.S. adult mortality: implications for understanding education's causal influence. *Soc Sci Med* 2015;127:8–18.
34. Pollitt RA, Rose KM, Kaufman JS. Evaluating the evidence for models of life course socioeconomic factors and cardiovascular outcomes: a systematic review. *BMC Public Health* 2005;5:7.
35. Mackenbach JP, Bopp M, Deboosere P, *et al.* Determinants of the magnitude of socioeconomic inequalities in mortality: a study of 17 European countries. *Health Place* 2017;47:44–53.
36. Cavelaars AE, Kunst AE, Geurts JJ, *et al.* Educational differences in smoking: international comparison. *BMJ* 2000;320:1102–7.
37. Tabuchi T, Kondo N. Educational inequalities in smoking among Japanese adults aged 25–94 years: nationally representative sex- and age-specific statistics. *J Epidemiol* 2017;27:186–92.
38. Pampel F, Legleye S, Goffette C, *et al.* Cohort changes in educational disparities in smoking: France, Germany and the United States. *Soc Sci Med* 2015;127:41–50.
39. Gregoraci G, van Lenthe FJ, Artnik B, *et al.* Contribution of smoking to socioeconomic inequalities in mortality: a study of 14 European countries, 1990–2004. *Tob Control* 2017;26:260–8.