BMJ Open The impact of general practitioners' gender on process indicators in Hungarian primary healthcare: a nation-wide cross-sectional study

Nóra Kovács,¹ Orsolya Varga,¹ Attila Nagy,¹ Anita Pálinkás,¹ Valéria Sipos,¹ László Körösi,² Róza Ádány,^{1,3,4} János Sándor¹

To cite: Kovács N, Varga O, Nagy A, *et al*. The impact of general practitioners' gender on process indicators in Hungarian primary healthcare: a nation-wide cross-sectional study. *BMJ Open* 2019;**9**:e027296. doi:10.1136/bmjopen-2018-027296

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-027296>).

Received 15 October 2018
Revised 03 August 2019
Accepted 16 August 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Preventive Medicine, Faculty of Public Health, University of Debrecen, Debrecen, Hungary

²National Institute of Health Insurance Fund Management, Budapest, Hungary

³WHO Collaborating Centre on Vulnerability and Health, Debrecen, Hungary

⁴MTA-DE Public Health Research Group, University of Debrecen, Debrecen, Hungary

Correspondence to

Dr Orsolya Varga;
varga.orsolya@sph.unideb.hu

ABSTRACT

Objectives The objectives of our study were (1) to investigate the association between gender of the general practitioner (GP) and the quality of primary care in Hungary with respect to process indicators for GP performance and (2) to assess the size of the gender impact.

Study design A nation-wide cross-sectional study was performed in 2016.

Setting and participants The study covered all general medical practices in Hungary (n=4575) responsible for the provision of primary healthcare (PHC) for adults. All GPs in their private practices are solo practitioners.

Main outcome measures Multilevel logistic regression models were used to analyse the association between GP gender and process indicators of PHC, and attributable proportion (AP) was calculated.

Results 48% of the GPs (n=2213) were women in the study. The crude rates of care provided by female GPs were significantly higher for seven out of eight evaluated indicators than those provided by male GPs. Adjusted for practice, physician and patient factors, GP gender was associated with the haemoglobin A1c (HbA1c) measurement: OR=1.18, 95% CI (1.14 to 1.23); serum creatinine measurement: OR=1.14, 95% CI (1.12 to 1.17); lipid measurement: OR=1.14, 95% CI (1.11 to 1.16); eye examination: OR=1.06, 95% CI (1.03 to 1.08); mammography screening: OR=1.05, 95% CI (1.03 to 1.08); management of patients with chronic obstructive pulmonary disease: OR=1.05, 95% CI (1.01 to 1.09) and the composite indicator: OR=1.08, 95% CI (1.07 to 1.1), which summarises the number of care events and size of target populations of each indicator. The AP at the specific indicators varied from 0.97% (95% CI 0.49% to 1.44%) of influenza immunisation to 8.04% (95% CI 7.4% to 8.67%) of eye examinations.

Conclusion Female GP gender was an independent predictor of receiving higher quality of care. The actual size of the gender effect on the quality of services seemed to be notable. Factors behind the gender effect should receive more attention in quality improvement particularly in countries where the primary care is organised around solo practices.

INTRODUCTION

The gender composition of physicians has changed in the last decades in all organisation

Strengths and limitations of this study

- The nation-wide study on process indicators of primary healthcare (PHC) covered all general medical practices in Hungary responsible for the provision to adults, avoided selection bias.
- Our investigation could estimate the population-level impact of the general practitioner (GP) gender effect.
- Although the analyses were controlled for several factors (GP age; patient age, gender, relative education; and practice size, types of settlement and regional location) influencing the quality of PHC, by involving other potential confounding factors (such as onset/duration of a chronic diseases, preferences and conditions of patients), reliability of our estimations for risks and impacts could be further improved.

for economic co-operation and development (OECD) countries. The proportion of female physicians has increased from 38% (in 2000) to 46% (in 2015), and this trend is expected to continue.¹ The Central and Eastern European countries showed the highest share of female physicians in 2015. Extremes of this share were observed in Latvia (74.4%) and in Estonia (74.6%), but the strongest growing trend was in the Netherlands, where the proportion of female physicians increased from 35.3% to 52.6% from 2000 to 2015. However, changes in gender proportions are not homogeneous in the different medical specialties; some, such as family medicine, are more affected.^{1–4}

According to the international literature, there is solid evidence that the physician's gender influences the effectiveness of medical care in primary healthcare (PHC).^{5–9} Female physicians in primary care tend to deliver more preventive services,^{10–18} they have better adherence to guidelines of chronic diseases^{8, 19} and they achieve

better intermediate treatment outcomes (blood pressure, haemoglobin A1c (HbA1c) and cholesterol level), compared with their male colleagues.^{20 21}

Several arguments could partially explain the existing differences between the quality of services provided by the two genders. Female physicians more often use patient-centred approach; moreover, they effectively facilitate patient participation in the medical exchange.^{8 22 23} According to studies in clinical practice, female physicians spend significantly more time with their patients and have longer consultations.^{6 24–26} Others have underlined the significance of the patient-centred communication style, which is more frequently applied by female physicians.^{8 27 28} Empathy is a further aspect of physicians' behaviour associated with better patient compliance and outcomes in chronic disease care^{23 29–31} and a higher level of empathy is a well-known characteristic of women.^{32 33} Additionally, the level of empathy seems to have an impact on career choices; individuals with higher empathy tend to prefer the primary care specialty.^{30 33 34} Similar specialty preference was found among both medical students and physicians.³⁰

The association between the gender of the provider (especially general practitioners (GPs)) and the effectiveness of the service is well demonstrated in many settings, but the exploration of how the gender of the physician actually affects the care practices and assessments of the size of the effect are insufficient.

Hungary provides an ideal field to examine the gender effect at the level of primary care in detail. The country has one of the highest shares of female physicians of general medicine and paediatrics, with a total share of 55.9% in 2015¹ and 56.2% in 2017.³⁵ In Hungary, the gender of the GP can be used as a potential factor affecting general medical practice (GMP)-level performance indicators because each individual GP works in their own primary care district with a territorial supply obligation (municipalities are responsible for the provision of primary care for the local population within their territory). Since 2009, the performance of providers has been monitored for quality indicators by the National Institute of Health Insurance Fund Management (NIHIFM), which have contracts with all GPs in the country. This monitoring exclusively applies process indicators to assess the quality of chronic disease management in primary care. The evaluation of GMPs and the provision of financial incentives are based on reaching desired target values for these indicators.³⁶

The objectives of our study were (1) to investigate the association between GP gender and the quality of primary care with respect to various process indicators for GMP performance and (2) to assess the size of the gender impact.

METHODS

Setting

The healthcare system in Hungary is based on compulsory health insurance with universal coverage. Primary care services are provided by GPs working in solo practices; therefore, one GMP is owned and operated by one GP.

In vacant GMPs, the services are provided by temporary GPs with restricted availability in time and place. The GPs are contracted with the NIHIFM, and they have territorial supply obligation, but patients can choose and change their primary care provider without any restriction.³⁷

Study design

A nation-wide cross-sectional study was performed, which covered all GMPs in Hungary responsible for the adult primary care in 2016 (n=4852). Demographic data of 7 207 186 clients (above 18 years) and 4575 GPs, the GMPs' organisational characteristics and data on performance indicators for GMPs were provided by NIHIFM. Vacant GMPs without a permanent GP during the study period (5.71%, n=277) were excluded, as this secondary analysis aimed to evaluate the influence of GPs' gender on PHC indicators.

Patient and public involvement

Patients and the public were not involved in this study.

Patient-level, physician-level and organisational characteristics of GMPs

Patient-specific and GMP-specific data were provided by NIHIFM. The number of adults registered in each GMP was determined by gender and age groups of 18–19, 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, 80–84, 85–89, and 90 years and above in the database. The age group of 65–69 (and 60–64 where 65 years and above patients were not available) was used as a reference. The age of the GP was a continuous variable. The male gender of GPs and patients were used as a reference in the analyses.

GMPs were categorised by the practice size (number of insured people registered in each GMP as follows: <800, 801–1200, 1201–1600, 1601–2000 and >2000; the categories are defined by NIHIFM), and types of settlement (rural and urban). The regional location of each GMP was described by the county (Baranya, Bács-Kiskun, Békés, Borsod-Abaúj-Zemplén, Csongrád, Fejér, Győr-Moson-Sopron, Hajdú-Bihar, Heves, Komárom-Esztergom, Nógrád, Pest, Somogy, Szabolcs-Szatmár-Bereg, Jász-Nagykun-Szolnok, Tolna, Vas, Veszprém and Zala counties) and the capital city (Budapest). The practice size of 1201–1600, the rural settlement and Budapest were used as reference categories.

The relative education was used to indicate the educational attainment of adults registered in each GMP. This indicator was calculated using gender, age group (7–19, 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, 80–84, 85–89 and 90 years and above), and settlement-specific data from 2011 Hungarian Census data provided by the Hungarian Central Statistical Office. The summarised length of education was calculated for each settlement. The expected number of school years was determined for adults in each settlement by the demographic characteristics of the settlement and the national reference values.

The ratio of observed and expected values, representing the settlement-specific relative education, was used to express the relative education of adults living in a certain settlement compared with the national reference level. A weighted settlement-specific relative education variable was calculated for GMPs providing care for patients from more than one settlement.³⁸

Performance indicators for GMPs

Each routine indicator of NIHIFM on immunisation, cancer screening and chronic disease management were used to assess the performance of GMPs: (1) proportion of patients with influenza immunisation; (2) proportion of patients who participated in screening mammography; (3) proportion of patients with hypertension with serum creatinine measurements; (4) proportion of patients with diabetes and/or hypertension with lipid profile measurements; (5) proportion of beta-blocker use among patients with ischaemic heart disease; (6) proportion of patients with diabetes with HbA1c measurements; (7) proportion of patients with diabetes who attended eye examination and (8) proportion of patients with chronic obstructive pulmonary disease (COPD) who participated in pulmonary functional testing. A composite indicator

(proportion of patients who received recommended care) was calculated by summarising the number of care events and the size of the target groups by each studied indicator. Indicators reflect the proportion of patients who received the care in each GMP. Details about the target groups and definitions of the assessed process indicators are shown in table 1.

Statistical analysis

Differences between baseline characteristics of GMPs were evaluated by the Pearson χ^2 test for categorical variables and two-sample t-test for continuous variables.

We used multilevel logistic regression models because of the binary outcome variables and in order to account for the clustering effect of patients within physicians. The models were adjusted for characteristics of patients (age and gender), GPs (age) and GMPs (practice size, types of settlement, geographical location and relative education) to assess the effect of the gender of the GP on their performance in case of each indicator. ORs with the corresponding 95% CIs, and robust standard errors were estimated. We presented intraclass correlation coefficient, which shows the proportion of total variance explained by the physician as grouping factor. The Hosmer-Lemeshow

Table 1 Definition and the target group of the studied PHC indicators

Indicator	Target group	Definition
Influenza immunisation	Patients over 65 years	Proportion of PHC patients, age 65 years and older, who received an influenza immunisation in the previous 12 months
Mammography screening	45–65 years-old female PHC patients	Proportion of female PHC patients, age 45–65 years, who received mammography in the previous 24 months
Serum creatinine measurement	Patients who redeemed antihypertensive drug at least four times in the previous 12 months (considered as patients with hypertension)	Proportion of PHC patients with hypertension screened for serum creatinine in the previous 12 months
Lipid measurement	Patients who redeemed antihypertensive drug at least four times in the previous 12 months (considered as patients with hypertension)	Proportion of PHC patients with hypertension and/or diabetes screened for lipid abnormalities in the previous 12 months
Beta-blocker application	Patients with AMI or CABG or PTCA	Proportion of patients with AMI, CABG or PTCA who used beta-blockers in the previous 12 months
HbA1c measurement	ATC A10 drug users	Proportion of PHC patients with diabetes mellitus screened for HbA1c in the previous 12 months
Eye examination	ATC A10 drug users	Proportion of PHC patients with diabetes mellitus who attended eye examination in the previous 12 months
Management of COPD	ATC R03 drug users and patients with COPD	Proportion of PHC patients with COPD who attended pulmonary function testing in the previous 12 months
Composite indicator	Total number of patients of each studied indicator	Proportion of PHC patients provided with recommended care; a general indicator of performance of GMP by summarising the number of care events and size of target populations of each studied indicator

AMI, acute myocardial infarction; CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; GMP, general medical practice; HbA1c, haemoglobin A1c; PHC, primary healthcare; PTCA, percutaneous transluminal coronary angioplasty.

test was used to determine the goodness of fit of the model (by comparing predicted and observed outcomes in deciles of predicted risk), which verified the adequacy of the model.

We calculated the missed care events, that is the number of care events not meeting the number of events which would be expected if the care was provided by female GP. The expected number of care events was calculated for GMPs with male GPs, using the demographic composition of adults belong to GMPs and considering female GPs as a reference, to describe the gender impact on service use in the whole country. Both observed and expected number of events were summarised for the whole country. The differences between the summarised observed and expected number of care events were calculated for GMPs with male GPs to quantify the differences in the number of care events compared with female GPs. Attributable proportion (AP) with 95% CI was also calculated by the summarised number of events to show the percentage of care events that can be attributed to the male GP gender.

Statistical significance was considered $p < 0.05$. Statistical analyses were performed using STATA IC V.13.0 software.

RESULTS

Descriptive statistics

The studied 4575 GPs (providing care for adults) consisted of 2213 (48.4%) women, and 2362 (51.6%) men ($p < 0.001$). Female GPs were younger (55.3 years vs 57.8 years; $p < 0.001$), more likely to work in urban regions (75.8% vs 62.0%; $p < 0.001$) and had smaller sized GMPs ($p < 0.001$). Patients of female physicians were more likely to be female (54.6% vs 52.31%; $p < 0.001$). The mean relative education was higher in GMPs with female GPs (1.026 vs 0.989; $p < 0.001$). The patient, physician and organisational characteristics of GMPs are summarised in [table 2](#).

Female GPs were younger. In the group of female GPs under 55 years and male GPs above 65 years were over-represented ([figure 1](#)).

According to the crude values of the PHC indicators (aggregated by the gender of the GP), statistically significant differences were found between male and female physicians. Patients of female GPs were more likely to have had serum creatinine (70.9% vs 66.5%; $p < 0.001$), lipid (64.5% vs 59.7%; $p < 0.001$) and HbA1c (80.6% vs 76.8%; $p < 0.001$) measurements; ophthalmological examinations (41.3% vs 38.2%; $p < 0.001$); and mammography screening (46.7% vs 44.7%; $p < 0.001$). They were also more likely to be managed properly in cases of COPD (79.1% vs 77.3%; $p < 0.001$). Although there was statistically significant differences between genders related to vaccination against influenza ($p = 0.004$), the size of difference was negligible (20.1% vs 19.9%). Patients with male GPs were more likely to receive beta-blockers (52.3% vs 53.5%; $p < 0.001$) ([table 3](#)).

Multilevel analyses

According to the multilevel logistic regression analysis, patients of female GPs were significantly more likely to receive care according to guidelines. The female gender of GPs was associated with hypertension and diabetes care-related indicators (HbA1c measurement, serum creatinine measurement, lipid measurement and eye examination), mammography screening, management of COPD patients and the composite indicator. No association was observed between influenza immunisation, beta-blocker application and GP gender. The proportion of total variability attributable to physician after adjusting for characteristics of physician, patient and practice varied between 2.4% and 17.9% across the indicators ([table 4](#)). Gender had one of the strongest effects among all studied variables on hypertension and diabetes care indicators (HbA1c, serum creatinine and lipid measurement). Results of the regression models are presented in Supplementary Table 1-2.

Population-level impact

Overall, 5.9% (95% CI 5.77% to 6.02%) of missed care events are attributable to male GP gender, which corresponds to 144 871 care events in the country per year. Restricting this summary to indicators with a significant association with GP gender (omitting influenza vaccination and beta-blocker application), the missed number of interventions was 144 373.

Compared with the GMPs with female GPs, the highest AP was observed for lipid measurement (AP=7.93%, 95% CI 7.71% to 8.16%), whereas the lowest AP was observed for influenza immunisation (AP=0.97%, 95% CI 0.49% to 1.44%). A notable percentage of missed care events was attributed to GPs' male gender in the case of serum creatinine measurement (AP=6.4%, 95% CI 6.18% to 6.61%), eye examinations (AP=8.04%, 95% CI 7.4% to 8.67%) and HbA1c measurements (AP=5.09%, 95% CI 4.65% to 5.54%). Mammography screening was not performed for 4.36% (95% CI 4.00% to 4.71%) of women in the target group, whereas COPD management was missed for 2.33% (95% CI 1.61% to 3.04%) of patients if the GP was male. Only beta-blocker application showed more care events that were attributed to male gender of GPs (AP=2.42%, 95% CI 1.53% to 3.31%) ([table 5](#)).

DISCUSSION

Main findings

Our results showed that GP gender was independently associated with quality of care with respect to process indicators for cancer screening and the management of chronic diseases (hypertension, diabetes and COPD). Patients of female GPs were more likely to have their HbA1c levels, serum creatinine, lipid measured, as well as their ophthalmological examination and pulmonary function testing implemented for monitoring purposes. Moreover, female patients of female GPs received significantly more mammography. Gender effect seems to have

Table 2 Patient-level, physician-level and organisational characteristics of GMPs

	All GMPs	GMPs with male GPs	GMPs with female GPs	P value
Number of GMPs	n=4575	n=2362 (51.6%)	n=2213 (48.4%)	
Age of GP, mean (SD)	56.6 (11.2)	57.8 (11.3)	55.3 (10.9)	<0.001
Number GMPs' patients				
Male	3 360 154 (46.6%)	1 794 560 (47.7%)	1 565 594 (45.5%)	<0.001
Female	3 847 032 (53.4%)	1 968 309 (52.3%)	1 878 723 (54.6%)	
Total	7 207 186	3 762 869	3 444 317	
Age (years) groups of patients				
18–19	121 802 (1.69%)	67 609 (1.8%)	54 193 (1.57%)	<0.001
20–24	483 058 (6.7%)	256 773 (6.82%)	226 285 (6.57%)	
25–29	539 957 (7.49%)	280 817 (7.46%)	259 140 (7.52%)	
30–34	553 972 (7.69%)	283 279 (7.53%)	270 693 (7.86%)	
35–39	690 907 (9.59%)	352 765 (9.37%)	338 142 (9.82%)	
40–44	752 110 (10.44%)	389 203 (10.34%)	362 907 (10.54%)	
45–49	647 431 (8.98%)	339 710 (9.03%)	307 721 (8.93%)	
50–54	534 270 (7.41%)	284 021 (7.55%)	250 249 (7.27%)	
55–59	561 965 (7.8%)	298 506 (7.93%)	263 459 (7.65%)	
60–64	665 236 (9.23%)	347 758 (9.24%)	317 478 (9.22%)	
65–69	542 777 (7.53%)	283 530 (7.53%)	259 247 (7.53%)	
70–74	416 851 (5.78%)	215 507 (5.73%)	201 344 (5.85%)	
75–79	316 345 (4.39%)	165 255 (4.39%)	151 090 (4.39%)	
80–84	213 912 (2.97%)	111 762 (2.97%)	102 150 (2.97%)	
85–89	117 420 (1.63%)	61 119 (1.62%)	56 301 (1.63%)	
>90	49 322 (0.68%)	25 378 (0.67%)	23 944 (0.7%)	
Relative education, mean (SD)	1.000 (0.1)	0.989 (0.2)	1.026 (0.1)	<0.001
Types of settlement				
Urban	3143 (68.7%)	1465 (62.0%)	1678 (75.8%)	<0.001
Rural	1432 (31.3%)	897 (37.9%)	535 (24.2%)	
GMP size (number of patients)				
<800	126 (2.8%)	66 (2.9%)	60 (2.7%)	<0.001
801–1200	646 (14.1%)	322 (13.6%)	324 (14.6%)	
1201–1600	1464 (32.0%)	690 (29.2%)	774 (34.9%)	
1601–2000	1415 (30.9%)	742 (31.4%)	673 (30.4%)	
>2000	924 (20.2%)	542 (22.9%)	382 (17.3%)	

GMP, general medical practice; GP, general practitioner.

a clinical relevance mostly on hypertension and diabetes care-related indicators, considering both the higher effect of the GP's gender (HbA1c measurement: OR=1.18, 95% CI 1.14 to 1.23, serum creatinine: OR=1.14, 95% CI 1.12 to 1.17 and lipid measurement: OR=1.14, 95% CI 1.11 to 1.16) and size of the affected population compared with other indicators (mammography screening, eye examination and management of COPD) where we also found statistically significant differences. A lack of gender influence was observed only for influenza immunisation and for beta-blocker usage.

The size of this gender effect proved to be notable. A high proportion ($AR_{total}=5.9\%$) of missed care events could be attributed to this gender effect. More than 144 000 care events per year were missed (out of 2 600 827 expected events) due to gender effect among patients of male GPs in Hungary. Primarily, missed lipid (59 723 missed laboratory investigations) and serum creatinine measurements (52 130 missed laboratory investigations) were responsible for this impact. Missed examinations among patients with diabetes mellitus (9801 missed HbA1c measurements and 7686 missed eye examinations)

Figure 1 The age and gender distribution of Hungarian GPs responsible for the adult primary healthcare in 2016. GPs, general practitioners.

and missed breast cancer screenings (13 285 screening tests) had a secondary, but significant, contribution.

Comparison with other studies

Others found similar gender differences in the management of diabetes^{8 20 21 39 40} and cancer^{8 41} screening. In addition, several studies have found an association between the management of hypertension and the gender of the physician. Female GPs tend to reach the blood pressure and lipid-level treatment goals more often^{42 43} and tend to focus better on cardiovascular risk factor control.^{15 39 42 43} Altogether, a positive association was observed between female gender and better quality of PHC in our investigation, in line with international experiences. Besides, our findings are consistent with other studies found that relatively low percentage of the variance is attributable to physicians on process measures after adjusting for characteristics of the physician, patient and practice.^{44 45}

Strength and limitations

A strength of this study was that it covered the entire Hungarian population, avoiding selection bias in the analysis of the association between GP gender and PHC indicators and resulting in the representativeness of our results for the whole Hungarian PHC population. As a consequence, the main strength of our investigation was that it could estimate the population-level impact of the GP gender effect, which has rarely been investigated as reflected by a scarcity of relevant publications.

A limitation of the study is that we were not able to monitor the changes over time due to the cross-sectional design.

	Number of patients received the care		Number of people in target group		Proportion of patients received the care		P value
	Male GPs	Female GPs	Male GPs	Female GPs	Male GPs	Female GPs	
Influenza immunisation	172 150	159 893	865 187	796 380	19.9%(19.8–19.9)	20.1%(19.9–20.2)	0.004
Mammography screening	304 907	299 137	681 141	640 293	44.7%(44.6–44.9)	46.7%(46.5–46.9)	<0.001
Serum creatinine measurement	814 819	785 950	1 225 158	1 108 168	66.5%(66.4–66.6)	70.9%(70.8–71.0)	<0.001
Lipid measurement	752 737	735 502	1 261 490	1 139 843	59.7%(59.6–59.8)	64.5%(64.4–64.6)	<0.001
Beta-blocker application	48 202	42 184	90 112	80 723	53.5%(53.2–53.8)	52.3%(51.9–52.6)	<0.001
HbA1c measurement	192 370	179 474	250 575	222 690	76.8%(76.6–76.9)	80.6%(80.4–80.8)	<0.001
Eye examination	95 617	91 995	250 575	222 690	38.2%(37.9–38.4)	41.3%(41.1–41.5)	<0.001
Management of COPD	75 154	69 835	97 213	88 345	77.3%(77.1–77.6)	79.1%(78.8–79.3)	<0.001
Composite indicator	2 455 956	2 363 970	4 721 451	4 299 132	52.0%(51.9–52.1)	54.9%(54.9–55.0)	<0.001

COPD, chronic obstructive pulmonary disease; GPs, general practitioners; HbA1c, haemoglobin A1c.

Table 3 The number of patients received the care, the number of people in the target groups and the proportion of patients received the care in 2016 for the whole country by gender of GPs with 95% CIs

Table 4 Influence of female gender of GPs (OR, 95% CIs, robust standard errors) on the primary healthcare indicators according to multilevel logistic regression analysis controlled for age of GP, age and gender of the patients, relative education of patients, practice size, types of settlement and regional location of GMP in 2016 and the ICC of tested models

Indicators	OR	95% CI	Robust SE	ICC (%)
Influenza immunisation (over 65 years)	1.03	0.98 to 1.09	0.028	17.9
Mammography screening (age 45–65 years)	1.05	1.03 to 1.08	0.014	4.8
Serum creatinine measurement	1.14	1.12 to 1.17	0.014	4.1
Lipid measurement	1.14	1.11 to 1.16	0.014	4.6
Beta-blocker application	0.98	0.96 to 1.01	0.013	2.4
HbA1c measurement	1.18	1.14 to 1.23	0.022	8.4
Eye examination	1.06	1.03 to 1.08	0.013	3.3
Management of COPD	1.05	1.01 to 1.09	0.019	4.9
Composite indicator	1.08	1.07 to 1.10	0.008	1.8

Reference group is male GPs.

COPD, chronic obstructive pulmonary disease; GMP, general medical practice; GPs, general practitioners; HbA1c, haemoglobin A1c; ICC, intraclass correlation coefficient.

The relative educational level, which was used to indicate the educational attainment for GMPs, cannot be considered as a year-specific and GMP-specific indicator. The information on education was not available for the investigated year, and the relative educational level of the adults living in the settlement (settlement-specific) was considered, regardless of the number of GPs in one settlement.

Additionally, the fact that the range of variables in the NIHIFM database which could be included in the analysis is limited, careful interpretation is required. Although the analyses were controlled for GPs' and patients' age and gender, educational attainment, GMP practice size, types of settlement and regional location, there were confounding factors that were not included in our models, limiting the reliability of the presented risk measures. Selection of the GP may be affected by patients preferences and expectations, as patients have free

choice of healthcare provider in Hungary. Since most of the GPs in Hungary are self-employed workers organising the provision on their own authority, the length of their working hours, the length of the consultation time and the availability of support staff were not measurable. The availability of support staff in solo practices or work in group practice could be an important factor in determining whether and how a patient will be treated.

Our analysis, which was based on data on solo GP practices, may differ from findings in group GP practices where professional cooperation with other providers may mitigate the effects of any non-clinical factors (eg, gender of GP).

Further research need

Our findings on impact of GP gender (mainly for HbA1c, serum creatinine and lipid measurement) suggest that further consideration of the effect is needed to identify

Table 5 Number of observed and expected care events in 2016, with absolute differences and attributable proportion estimations (with 95% CI) in Hungary among 3 762 869 patients provided by male GPs

Indicators	Number of observed cases	Number of expected cases	Absolute difference	Attributable proportion
Influenza immunisation	172 150	173 813	−1 663	0.97% (0.49%–1.44%)
Screening mammography	304 907	318 192	−13 285	4.36% (4%–4.71%)
Serum creatinine measurement	814 819	866 949	−52 130	6.4% (6.18%–6.61%)
Lipid measurement	752 737	812 459	−59 723	7.93% (7.71%–8.16%)
Beta-blocker application	48 202	47 037	1 165	2.42% (1.53%–3.31%)
HbA1c measurement	192 370	202 170	−9 801	5.09% (4.65%–5.54%)
Eye examination	95 617	103 302	−7 686	8.04% (7.4%–8.67%)
Management of COPD	75 154	76 901	−1 748	2.33% (1.61%–3.04%)
Total (composite indicator)	2 455 956	2 600 827	−144 871	5.9% (5.77%–6.02%)

COPD, chronic obstructive pulmonary disease; GPs, general practitioners; HbA1c, haemoglobin A1c.

the details and mechanisms behind the gender effect to improve the adequacy of targeted interventions. Patients with early stage diabetes mellitus are more cooperative with female GPs, but patients with advanced stages of the disease, realising the seriousness of the consequences, are more sensitive to male GP directions. The onset/duration of a chronic disease and accordingly the complex needs of patients may modify the gender effect.⁴⁶ Therefore, it would be worthwhile to investigate the potential influences on performance indicators that are differentiated according to the preferences^{47–49} and type of their health problem.⁵⁰

Furthermore, there is a possible extension of our presented investigation. Analysing outcome indicators on the prognosis of patients with chronic diseases in PHC is obviously required to more convincingly describe the public health impact of the GP gender effect.

Implications

Poor-quality healthcare services put a great burden on payers and society everywhere.⁵¹ The effective utilisation of experiences of high-income countries⁵² addressing gender differences in provision is crucial in low-income and middle-income countries such as Hungary. High-quality PHC is a shared goal of the patients, the GPs and the state.

According to our observations, a possible way to improve the quality of PHC is to raise awareness about the significance of the GPs' gender effect and its underlying causes. Keeping in mind that large sample size can detect even the smallest differences, the interpretation needs to consider that these small differences may have limited importance at population level; our results indicate the significance and urge the expansion of interventions aiming at improved communication skills of medical students and GPs, and at shifting the attitude of GPs. Incorporating practical knowledge on how the communication strategies and empathy in clinical practice influence the quality of services (which is often overlooked by medical students and practitioners) may facilitate the development of a patient-centred care culture.

It is well demonstrated that a behaviour change targeting improved patient outcomes is feasible by training physicians and medical students.²³ Since teaching patient-oriented communication skills to medical students increases their competency, as well as patients' satisfaction and health outcomes. Training in effective communication skills, embedded into the medical school curriculum, is widely recommended.^{53–56} Our results suggest that the scope of this training should be supplemented by gender-related attitudes and behavioural features.

These findings can be utilised in gender impact assessment regarding the feminisation of the primary care physician workforce.^{57 58}

CONCLUSIONS

Our results suggest the existence of a gender effect in Hungary on performance of GPs working in solo practices. Provision of guideline-recommended care was

observed more often in patients of female GPs. The actual impact of the gender effect on the quality of services was found to be notable, emphasising that the communication style and attitude associated with female gender should be considered as factors influencing the quality of PHC and should be targeted accordingly in training of medical students and GPs. However, further studies involving a broader range of clinical and non-clinical factors are needed to allow more precise assessment of gender differences.

Acknowledgements The authors thank the support of Zsófia Falusi and László Pál for providing the data for this study.

Contributors NK, LK and JS performed and designed the study. LK participated in the database preparation. NK, AP, VS and AN analysed and interpreted the data. NK and OV prepared the manuscript. RÁ and JS provided professional instruction on the subject and approved the final version to be submitted. JS revised the manuscript. All authors read and approved the final manuscript.

Funding The reported study was carried out in the framework of the 'Public Health Focused Model Programme for Organising Primary Care Services Backed by a Virtual Care Service Centre' (SH/8/1). The Model Programme is funded by the Swiss Government via the Swiss Contribution Programme (SH/8/1) in agreement with the Government of Hungary. Additional source of funding was from GINOP-2.3.2-15-2016-00005 project which was co-financed by the European Union and the European Regional Development Fund, and from EFOP-3.6.3-VEKOP-16-2017-00009 co-financed by EU and the European Social Fund. This work was also supported by the Portugal/Hungary Bilateral Project FCT/NKFIH - (TÉT_16-1-2016-0093) and the János Bolyai Scholarship of the Hungarian Academy of Sciences (MTA) to O.V.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The databases that we analysed were anonymised. The research protocol was reviewed and permitted by and performed in concordance with the Internal Data Safety and Patient Rights Board of the National Institute of Health Insurance Fund Management (E01/317-1/2014).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. OECD. OECD health statistics 2017. Available: <http://stats.oecd.org/>
2. OECD. Health at glance 2015, 2015. Available: <http://apps.who.int/medicinedocs/documents/s22177en/s22177en.pdf>
3. Canadian Institute for Health Information. *Supply, distribution and migration of Canadian physicians*, 2012. Ottawa: Canadian Institute for Health Information, 2013. http://publications.gc.ca/collections/collection_2013/icis-cihi/H115-23-2012-eng.pdf
4. Lambert EM, Holmboe ES. The relationship between specialty choice and gender of U.S. medical students, 1990-2003. *Acad Med* 2005;80:797-802.
5. Bertakis KD, Helms LJ, Callahan EJ, *et al.* The influence of gender on physician practice style. *Med Care* 1995;33:407-16.
6. Bensing JM, van den Brink-Muinen A, de Bakker DH. Gender differences in practice style: a Dutch study of general practitioners. *Med Care* 1993;31:219-29.
7. Frank E, Harvey LK. Prevention advice rates of women and men physicians. *Arch Fam Med* 1996;5:215-9.
8. Dahrouge S, Seale E, Hogg W, *et al.* A comprehensive assessment of family physician gender and quality of care: a cross-sectional analysis in Ontario, Canada. *Med Care* 2016;54:277-86.

9. Ewing Get *et al.* Self-Report of delivery of clinical preventive services by U.S. physicians comparing specialty, gender, age, setting of practice, and area of practice. *Am J Prev Med* 1999;17:62–72.
10. Franks P, Bertakis KD. Physician gender, patient gender, and primary care. *Journal of Women's Health* 2003;12:73–80.
11. Lurie N, Margolis KL, McGovern PG, *et al.* Why do patients of female physicians have higher rates of breast and cervical cancer screening? *J Gen Intern Med* 1997;12:34–43.
12. Chamot E, Charvet A, Perneger TV. Women's Preferences for Doctor's Involvement in Decisions about Mammography Screening. *Med Decis Making* 2004;24:379–85.
13. Levy S, Dowling P, Boulton L, *et al.* The effect of physician and patient gender on preventive medicine practices in patients older than fifty. *Fam Med* 1992;24:58–61.
14. Henderson JT, Weisman CS. Physician gender effects on preventive screening and counseling: an analysis of male and female patients' health care experiences. *Med Care* 2001;39:1281–92.
15. Tabenkin H, Eaton CB, Roberts MB, *et al.* Differences in cardiovascular disease risk factor management in primary care by sex of physician and patient. *Ann Fam Med* 2010;8:25–32.
16. Rosell-Murphy M, Rodriguez-Blanco T, Morán J, *et al.* Variability in screening prevention activities in primary care in Spain: a multilevel analysis. *BMC Public Health* 2015;15.
17. Diehl K, Gansefort D, Herr RM, *et al.* Physician gender and lifestyle counselling to prevent cardiovascular disease: a nationwide representative study. *J Public Health Res* 2015;4.
18. Ince-Cushman D, Correa JA, Shuldiner J, *et al.* Association of primary care physician sex with cervical cancer and mammography screening. *Can Fam Physician* 2013;59:e11–18.
19. Baumhäkel M, Müller U, Böhm M. Influence of gender of physicians and patients on guideline-recommended treatment of chronic heart failure in a cross-sectional study. *Eur J Heart Fail* 2009;11:299–303.
20. Berthold HK, Gouni-Berthold I, Bestehorn KP, *et al.* Physician gender is associated with the quality of type 2 diabetes care. *J Intern Med* 2008;264:340–50.
21. Kim C, McEwen LN, Gerzoff RB, *et al.* Is physician gender associated with the quality of diabetes care? *Diabetes Care* 2005;28:1594–8.
22. Roter DL, Hall JA. Why physician gender matters in shaping the physician-patient relationship. *J Womens Health* 1998;7:1093–7.
23. Beck RS, Daughtridge R, Sloane PD. Physician-Patient communication in the primary care office: a systematic review. *J Am Board Fam Pract* 2002;15:25–38.
24. Roter D, Lipkin M, Korsgaard A. Sex differences in patients' and physicians' communication during primary care medical visits. *Med Care* 1991;29:1083–93.
25. Sandhu H, Adams A, Singleton L, *et al.* The impact of gender dyads on doctor-patient communication: a systematic review. *Patient Educ Couns* 2009;76:348–55.
26. Jefferson L, Bloor K, Hewitt C. The effect of physician gender on length of patient consultations: observational findings from the UK hospital setting and synthesis with existing studies. *J R Soc Med* 2015;108:136–41.
27. Roter DL, Hall JA, Aoki Y. Physician gender effects in medical communication: a meta-analytic review. *JAMA* 2002;288:756–64.
28. Arnold RM, Martin SC, Parker RM. Taking care of patients--does it matter whether the physician is a woman? *West J Med* 1988;149:729–33.
29. Mercer SW, Higgins M, Bikker AM, *et al.* General practitioners empathy and health outcomes: a prospective observational study of consultations in areas of high and low deprivation. *Ann Fam Med* 2016;14:117–24.
30. Hojat M, Mangione S, Gonnella JS, *et al.* Empathy in medical education and patient care. *Acad Med J Assoc Am Med Coll* 2001;76.
31. Del Canale S, Louis DZ, Maio V, *et al.* The relationship between physician empathy and disease complications: an empirical study of primary care physicians and their diabetic patients in Parma, Italy. *Acad Med J Assoc Am Med Coll* 2012;87:1243–9.
32. Rueckert L, Naybar N. Gender differences in empathy: the role of the right hemisphere. *Brain Cogn* 2008;67:162–7.
33. Hojat M, Gonnella JS, Nasca TJ, *et al.* Physician empathy: definition, components, measurement, and relationship to gender and specialty. *AJP* 2002;159:1563–9.
34. Hojat M, Zuckerman M. Personality and specialty interest in medical students. *Med Teach* 2008;30:400–6.
35. Ádány R, Papp M. *Preventive services in primary care*. Budapest: Medicina, 2017.
36. National Institute of Health Insurance Fund Management. Quality indicators for performance evaluation of family doctors from January 2016 (in Hungarian). Budapest: : National Institute of Health Insurance Fund Management, 2016. Available: http://neak.gov.hu/data/cms1010747/Haziorvosok_indikator_alapu_teljesitmeny_NEAK.pdf [Accessed 30 May 2017].
37. Gaal P, Szigeti S, Csere M, *et al.* Hungary health system review. *Health Syst Transit* 2011;13:1–266.
38. Sándor J, Pálkás A, Vincze F, *et al.* Association between the general practitioner workforce crisis and premature mortality in Hungary: cross-sectional evaluation of health insurance data from 2006 to 2014. *Int J Environ Res Public Health* 2018;15:1388. 10.3390/ijerph15071388.
39. Schmittiel JA, Traylor A, Uratsu CS, *et al.* The association of patient-physician gender concordance with cardiovascular disease risk factor control and treatment in diabetes. *J Womens Health* 2009;18:2065–70.
40. Tran AT, Bakke Åsne, Berg TJ, *et al.* Are general practitioners characteristics associated with the quality of type 2 diabetes care in general practice? results from the Norwegian ROSA4 study from 2014. *Scand J Prim Health Care* 2018;36:170–9.
41. Lurie N, Slater J, McGovern P, *et al.* Preventive care for women -- does the sex of the physician matter? *N Engl J Med* 1993;329:478–82.
42. Journath G, Hellénus M-L, Manhem K, *et al.* Association of physician's sex with risk factor control in treated hypertensive patients from Swedish primary healthcare. *J Hypertens* 2050;2008.
43. Nilsson PM, Journath G. Effective Consultation for Patients with Hypertension: Does the Physician's Gender Matter for Risk-Factor Control? *Womens Health* 2008;4:433–4.
44. Fung V, Schmittiel JA, Fireman B, *et al.* Meaningful variation in performance: a systematic literature review. *Med Care* 2010;48:140–8.
45. Wong HJ, Siström CL, Benzer TI, *et al.* Use of imaging in the emergency department: physicians have limited effect on variation. *Radiology* 2013;268:779–89.
46. Nagy A, Nagy B, Adány R, *et al.* Determinants of low referral rates for ophthalmologic examination in people with type 2 diabetes in Hungary. *Diabetes Res Clin Pract* 2013;102:e29–31.
47. Fennema K, Meyer DL, Owen N. Sex of physician: patients' preferences and stereotypes. *J Fam Pract* 1990;30:441–6.
48. Jung HP, Baerveldt C, Olesen F, *et al.* Patient characteristics as predictors of primary health care preferences: a systematic literature analysis. *Health Expectations* 2003;6:160–81.
49. Graffy J. Patient choice in a practice with men and women general practitioners. *Br J Gen Pract J R Coll Gen Pract* 1990;40:13–15.
50. Delgado A, López-Fernández L-A, Luna JdeD, *et al.* The role of expectations in preferences of patients for a female or male general practitioner. *Patient Educ Couns* 2011;82:49–57.
51. Andel C, Davidow SL, Hollander M, *et al.* The economics of health care quality and medical errors. *J Health Care Finance* 2012;39:39–50.
52. Jefferson L, Bloor K, Birks Y, *et al.* Effect of physicians' gender on communication and consultation length: a systematic review and meta-analysis. *J Health Serv Res Policy* 2013;18:242–8.
53. Haq C, Steele DJ, Marchand L, *et al.* Integrating the art and science of medical practice: innovations in teaching medical communication skills. *Fam Med* 2004;36 Suppl:S43–50.
54. Choudhary A, Gupta V. Teaching communications skills to medical students: introducing the fine art of medical practice. *Int J App Basic Med Res* 2015;5:41–4.
55. Yedidia MJet *et al.* Effect of communications training on medical student performance. *JAMA* 2003;290:1157–65.
56. Karlberg L, Lindgren C. [Communication skills in the encounter with patients--current examination subject for medical students. Beneficial educational investment]. *Lakartidningen* 2004;101:3072–4.
57. European Institute for Gender Equality. *Gender impact assessment*. Luxembourg: Publications Office of the European Union, 2016. <https://eige.europa.eu/sites/default/files/mh0416171enn.pdf>
58. Hedden L, Barer ML, Cardiff K, *et al.* The implications of the feminization of the primary care physician workforce on service supply: a systematic review. *Hum Resour Health* 2014;12.