

Appendix 5 - Participant details

| PRM | Sample size | Setting | Age | % male | Education level | Ethnicity |
|---|-------------|------------------------|------------------------------------|--------|--|--|
| Satisfaction with Stroke Care questionnaire (SASC) | 149 | Community | Mean 71 | 50% | Not given | 80% white; 7% black Caribbean; 5% Bangladeshi, 2% black African, 2% Indian, 2% Pakistani, 1% other Asian, 1% not completed |
| Stroke Patient Reported Outcome Measure (Stroke-PROM) | 475 | Community and hospital | 11.6%≤45; 48.4%45-65; 40%≥65 | 60% | 38.5% primary school or lower; 30.3% junior high school; 18.5% senior high school; 12.6% college or higher | Not given |
| Barriers to Physical Activity after Stroke (BAPAS) | 109 | Community | Mean 60.6 (SD 12.8) | 63% | Not given | Not given |
| World Health Organisation Quality of Life-100 (WHOQOL-100) | 70 | Hospital | Mean 60.16 (SD 11.30) | 61% | 67.1% primary school graduates or less educated; 40% were retired. | Not given |
| Patient's Questionnaire on Participation in Discharge Planning (PQPDP) | 188 | Hospital | Mean 74 (SD 11.2) | 56% | 75% elementary school; 19% sec/high school; 6% university | Not given |
| Chao Perception of Continuity scale (Chao-PC) | 168 | Community | Mean 67.65 (SD 12.54) | 58% | 79% <16 yrs; 14.3% <18 years; 6% >18 yrs | 98.2% white British |