

Appendix 2. Data for Figure 2

Prevalence of key health status determinants by recorded diagnosis in 13,630 regular Aboriginal and Torres Strait Islander adults attending 18 participating health services in Queensland, April -September 2011.

Clinical Variable	Prevalence	IQR
Tobacco Use	0.51	0.40, 0.50
Alcohol	0.45	0.40, 0.50
BMI > 25	0.69	0.66, 0.73
BMI > 35 – Females only	0.45	0.43, 0.51
Diabetes Mellitus Type 2	0.20	0.15, 0.23
Hypertension	0.24	0.21, 0.27
eGFR < 60ml/min	0.11	0.09, 0.15

Appendix 3. Data for Figure 3

Performance of health assessments and care planning for regular Aboriginal and Torres Strait Islander patients, across 17 participating health services in Queensland, April - September 2011.

Care Activity - EPC items	Median	IQR
Type 2 Diabetes on GPMP	0.34	0.15, 0.51
CHD patients on GPMP	0.25	0.11, 0.45
Child Health Check (0-5)	0.38	0.26, 0.53
Child Health Check (6-14)	0.46	0.30, 0.61
Adult Health Check (15-54)	0.43	0.29, 0.50
Adult Health Check (55+)	0.60	0.23, 0.68