

Supplemental Table 1: Annual incidence of eating disorders by gender and eating disorder type per 100,000 population

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
All Eating Disorders	789/2,393,179	838/2,673,109	953/2,703,550	967/2,740,803	904/2,827,344	906/2,819,275	882/2,786,786	980/2,680,119	946/2,553,593	897/2,440,256
Incidence (95% CI)	33.0 (30.7-35.3)	31.4 (29.3-33.5)	35.3 (33.1-37.5)	35.3 (33.1-37.6)	32.0 (29.9-34.1)	32.1 (30.0-34.3)	31.7 (29.6-33.8)	36.6 (34.3-38.9)	37.1 (34.7-39.5)	36.8 (34.4-39.2)
Females										
All ED	732/1,374,900	777/1,504,681	884/1,513,075	899/1,518,646	839/1,532,386	853/1,515,225	805/1,486,937	894/1,434,592	873/1,367,358	816/1,302,016
Incidence (95% CI)	53.2 (49.5-57.2)	51.6 (48.0-55.4)	58.4 (54.6-62.4)	59.2 (55.4-63.2)	54.8 (51.1-58.5)	56.3 (52.6-60.2)	54.1 (50.5-58.0)	62.3 (58.3-66.5)	63.8 (59.7-68.2)	62.7 (58.4-67.1)
AN	195/1,374,900	189/1,504,681	177/1,513,075	234/1,518,646	196/1,532,386	188/1,515,225	172/1,486,937	234/1,434,592	196/1,367,358	177/1,302,016
Incidence (95% CI)	14.2 (12.3-16.3)	12.6 (10.9-14.5)	11.7 (10.1-13.5)	15.4 (13.5-17.5)	12.8 (11.1-14.7)	12.4 (10.7-14.3)	11.6 (10.0-13.4)	16.3 (14.3-18.5)	14.3 (12.4-16.5)	13.6 (11.7-15.7)
BN	294/1,374,900	330/1,504,681	361/1,513,075	378/1,518,646	323/1,532,386	346/1,515,225	312/1,486,937	303/1,434,592	324/1,367,358	269/1,302,016
Incidence (95% CI)	21.4 (19.0-24.0)	21.9 (19.7-24.4)	23.9 (21.5-26.4)	24.9 (22.5-27.5)	21.1 (18.9-23.5)	22.8 (20.5-25.3)	21.0 (18.8-23.4)	21.1 (18.8-23.6)	23.7 (21.2-26.4)	20.7 (18.3-23.2)
EDNOS	243/1,374,900	258/1,504,681	346/1,513,075	287/1,518,646	320/1,532,386	319/1,515,225	321/1,486,937	357/1,434,592	353/1,367,358	370/1,302,016
Incidence (95% CI)	17.7 (15.5-20.0)	17.2 (15.1-19.3)	22.9 (20.6-25.4)	18.9 (16.8-21.2)	20.9 (18.7-23.3)	21.1 (18.8-23.5)	21.6 (19.3-24.0)	24.9 (22.4-27.6)	25.8 (23.2-28.6)	28.4 (25.6-31.4)
Males										
All ED	57/1,018,279	61/1,168,428	69/1,190,475	68/1,222,157	65/1,294,958	53/1,304,050	77/1,299,849	86/1,245,527	73/1,186,235	81/1,138,240
Incidence (95% CI)	5.6 (4.3-7.2)	5.2 (4.0-6.6)	5.8 (4.5-7.3)	5.6 (4.3-7.0)	5.0 (3.9-6.3)	4.1 (3.1-5.3)	5.9 (4.7-7.4)	6.9 (5.6-8.5)	6.1 (4.9-7.7)	7.1 (5.7-8.8)
AN	8/1,018,279	12/1,168,428	21/1,190,475	17/1,222,157	4/1,294,958	6/1,304,050	4/1,299,849	18/1,245,527	12/1,186,235	15/1,138,240
Incidence (95% CI)	0.8 (0.4-1.5)	1.0 (0.6-1.7)	1.7 (1.1-2.7)	1.5 (0.9-2.3)	0.3 (0.1-0.7)	0.5 (0.2-0.9)	0.3 (0.1-0.7)	1.4 (0.9-2.2)	1.0 (0.5-1.7)	1.3 (0.8-2.1)
BN	14/1,018,279	14/1,168,428	14/1,190,475	12/1,222,157	19/1,294,958	15/1,304,050	20/1,299,849	17/1,245,527	21/1,186,235	18/1,138,240
Incidence (95% CI)	1.4 (0.8-2.2)	1.2 (0.7-2.0)	1.2 (0.7-1.9)	1.0 (0.7-1.9)	1.5 (0.9-2.3)	1.2 (0.7-1.9)	1.5 (1.0-2.3)	1.4 (0.8-2.1)	1.8 (1.1-2.8)	1.6 (1.0-2.4)
EDNOS	35/1,018,279	45/1,168,428	34/1,190,475	39/1,222,157	42/1,294,958	32/1,304,050	53/1,299,849	51/1,245,527	40/1,186,235	48/1,138,240
Incidence (95% CI)	3.4 (2.4-4.7)	3.8 (2.8-5.1)	2.9 (2.0-3.9)	3.2 (2.3-4.3)	3.2 (2.4-4.3)	2.4 (1.7-3.4)	4.1 (3.1-5.3)	4.1 (3.1-5.3)	3.4 (2.4-4.5)	4.2 (3.1-5.5)

ED: eating disorder; AN: anorexia nervosa, BN: bulimia nervosa, EDNOS: eating disorder not otherwise specified