

Table A1. Subgroup analysis: multivariate logistic regression results for subgroups based on control level

Variables	CAM usage by asthma control level, <i>OR (95% CI), p</i>		
	Controlled (N=91)	Partially Controlled (N=187)	Uncontrolled (N=202)
Age	1.01 (0.98;1.05), p=0.43	1.00 (0.97;1.01), p=0.33	1.00 (0.98-1.02), p=0.81
Gender			
Male	Reference	Reference	Reference
Female	1.77 (0.58;5.41), p=0.31	1.88 (0.95;3.72), p=0.07	1.31 (0.68-2.52), p=0.43
Household income			
<\$60,000	Reference	Reference	Reference
≥\$60,000	0.57 (0.18;1.76), p=0.32	0.52 (0.23;1.14), p=0.10	1.18 (0.63-2.20), p=0.60
Education			
Less than post-secondary education	Reference	Reference	Reference
Post-secondary education	1.13 (0.38;3.38), p=0.83	1.86 (0.77;4.47), p=0.17	0.77 (0.39-1.52), p=0.46
Ethnicity			
Caucasian	Reference	Reference	Reference
Asian	1.50 (0.38;5.70), p=0.56	1.02 (0.40;2.60), p=0.96	1.33 (0.59-3.01), p=0.49
Other	NA	1.92 (0.31;11.83), p=0.48	1.95 (0.44-8.61), p=0.38
Intake of controller medication			
Low	Reference	Reference	Reference
Moderate	0.92 (0.15-5.74), p=0.93	1.08 (0.43;2.70), p=0.87	1.70 (0.73-3.99), p=0.22
High	0.70 (0.22-2.19), p=0.54	0.80 (0.38;1.68), p=0.56	1.57 (0.79-3.11), p=0.20

OR, odds ratios; CI, confidence interval.