Authors	Year of publication	Disease	Drug	Number of participants	Adverse event	Other drugs described	Additional medical therapy according to standard procedures	NSAIDs use	Use of gastroprotection drugs	Randomization	Selection/exclusion	Peptic ulcer exclusion	Treatment level (hospitalization, AL, Ambulant)	Treatment Days	Follow up days	Definition of adverse effect	Methods of monitoring
Bradman, B., Schouten, H. J. A.,	1983	Traumatic injury, head	Dexamethasone	161	Major episode of GI bleeding which required blood transfusion	Probably. Artificially ventilation.				Randomization into two groups. Randomization procedure not described. Dexa/placebo vials were indistinguishable, and prepared by the pharmacy.	Pts who had received steroid therapy prior to admission were not entered into the study. Steroid/ventilation was stopped in 3 pts with bowel perforation or history of peptic ulcer.			180	Major episode of GI bleeding which required blood transfusion		None
Bracken, M. B., Shepard, M. J.,	1990	Traumatic injury, spinal cord	Methylprednisolone	333	Steroid group	Probably				Randomization by pharmacist. Three treatments were randomized in blocks of 3	Pts with pre-maintenance steroids, or received more than 100 mg methylpred or dexa			180	GI bleeding		
Dearden, N. M., Gibson, J. S.,	1986	Traumatic injury, head	Dexamethasone	130	Minor GI bleeding (blood staining-nasogastric aspirate). Major: GI bleeding necessitating transfusion	Antacids, osmotic diuretic agents, barbiturates, hypocapnic ventilation	Management according to a standard protocol Each center was expected to administer best treatment as indicated, and no attempt was made to standardize other drug treatment or neurosurgical interventions.			Randomization by pharmacist. Three treatments were randomized in blocks of 3 "Pts were randomly allocated" Randomization procedure not described.	Pts primarily excluded if gastric had been adm before randomization, known recent gastric or duodenal ulcer, or if gcs were inadvertently given, apart from the study medication.			180	Minor GI bleeding (blood staining nasogastric aspirate). Major: GI bleeding necessitating transfusion		Routine
Gaab, M. R., Trost, H. A.,	1994	Traumatic injury, head	Dexamethasone	299	Upper GI bleeding					Random permuted block design was used; randomization in blocks of 6. Vials containing the drug/placebo were prerandomized and assigned to each succeeding pt. No further description.	Each if preexisting history of peptic ulcer, steroids 2 weeks before the study. Contraindications were contraindications to steroids. Pts with seizures of steroids prior to admission were not excluded.			7/5	Upper GI bleeding		Systematic survey
Giannotta, S.L., Weiss, M.H.,	1984	Traumatic injury, head	Methylprednisolone	88	GI bleeding	Intubation, osmotic diuretics, barbiturates				Randomization by pharmacy	Each if preexisting history of peptic ulcer, steroids 2 weeks before the study. Contraindications were contraindications to steroids. Pts with seizures of steroids prior to admission were not excluded.			180	GI bleeding		Routine
Grumme, T., Baethmann, A.,	1995	Traumatic injury, head	Triamcinolone	396	GI bleeding	Osmotic agents, gastro-protective drugs, sedatives, antibiotics				Randomization by telephone to the pharmacist who obtained the patients height/weight and assigned the pt to a treatment group.	Pts who had received steroids, or received methylpred or equivalent before admission to hospital.			365	GI bleeding		Systematic survey
Matsumoto, T., Tamaki, T.,	2007	Traumatic injury, spinal cord	Methylprednisolone	66	GI bleeding	Broad-spectrum antibiotics, H2-blocker.				Central telephone randomization service provided by the CTSU or local pack system. Treatment allocation balanced for sex, age, time since injury, GCS-psyd fracture/severe and country.				60	GI bleeding		Systematic survey
Roquilly, A., Mahe, P. J.,	2011	Traumatic injury, multiple trauma	Hydrocortisone	150	Gastrointestinal bleeding or digestive perforation	Vasopressors, antibiotic prophylaxis, etomidate				Central randomization center. Randomized in 1:1 ration in fixed blocks of 4 and 6 by using a log to the treatment center who by a computerized number generator list.	Each if steroids last 6 months.			28	GI bleeding or digestive perforation		Routine
Chomnegard, V., Jepmonsant, S.,	1994	Tuberculous meningitis	Prednisolone	56		Tuberculostatics				Randomization by block of 4, coded treatments A and B					GI bleeding		Systematic survey
Chomnegard, V., Sawanyawisuth, K.,	2005	Encephalitic meningitis	Prednisolone	110		Alum milk, paracetamol				Stratification according to severity of headache and CSF opening pressure. Block 1 of 4 randomization.				14	GI bleeding not defined		Systematic survey
de Gans, J., Van De Beek, D.,	2002	Meningitis adults	Dexamethasone	301	Gastrointestinal bleeding (clinically relevant bleeding with a decreased serum hemoglobin level	Parenteral antibiotics				Computer-generated list of random numbers in blocks of six. Code was not broken before the last pt had completed the follow-up.	Each if hypersensitv to steroids or recent peptic ulcer disease.			7/5	GI bleeding (clinically relevant bleeding with a decreased serum hemoglobin level)		Systematic survey
Gijsen, D., Kumhar, M.R.,	2000	Meningitis adults	Dexamethasone	40	GI bleeding defined as oral or rectal gastrointestinal bleeding on visual inspection					Randomization was done by odd and even number of pts. Odd numbered pts received the drug and even the placebo.	Each if hepatic or renal dysfunction.			14	GI tract bleeding		None
Kilpi, T., Peltola, H.,	1995	Meningitis children	Dexamethasone	58		Ceftriaxone				A computer-generated list of random therapy assignments was kept at the hospital; the next adjunctive treatment regimen was obtainable by phone 24 hours a day.					GI bleeding not defined		Routine
King, S.M., Law, B.,	1984	Meningitis children	Dexamethasone	101	Occult blood in stool, GI bleed (perforation)	Cefuroxime/ceftriaxone.	Additional supportive care according to standard clinical practice.			Randomization was stratified by centre and by age. Double-blind manner for four days.	Each if current use of steroids or contraindications to the use ofsteroids, such as peptic ulcer disease.			4	Occult blood in stool, GI bleed (perforation)		Routine
Lebel, M.H., Freij, B.J.,	1988	Meningitis children	Dexamethasone	102	Gastrointestinal bleeding and heme-positive stools	Cefuroxime / ceftriaxone				Computer-generated list of random therapy assignments was used. The code was not broken until follow-up of all pts.	Each if hepatic or renal impairment.			14	GI bleeding and heme-positive stools		Routine
Lebel, M.H., Hoyt, M.J.,	1989	Meningitis children	Dexamethasone	60		Cefuroxime / ceftriaxone				Computer-generated list of random therapy assignments. The code not broken until follow-up of all pts. Double-blind manner.				14	GI bleeding not defined		Routine
Nguyen, T.H., Tran, T.H.,	2007	Meningitis adolescents adults	Dexamethasone	435	Minor GI bleeding. Withdrawal due to bleeding in the upper gastrointestinal tract.	Ceftriaxone				A computer-generated sequence of random numbers was used to assign treatment in blocks of 100 patients	Each if the attending physician believed corticosteroids were contraindicated			180	Minor GI bleeding. Withdrawal due to bleeding in the upper gastrointestinal tract.		None
Odio, C.M., Faingezicht, I.,	1991	Meningitis children	Dexamethasone	101	Gastric ulcer with overt hemorrhage	Ceftriaxone, intubation and mechanically ventilated if GCS <=7				Computer-generated list of random therapy assignments. Blinded fashion. The code was not broken until the last pt had completed therapy.				14	Gastric ulcer with overt hemorrhage		Routine
Peltola, H., Roine, I.,	2010	Meningitis children	Dexamethasone	329	Visible blood-I stool. Massive gastrointestinal bleeding	Ceftriaxone, antipyretics when needed, anticonvulsives.				Stratified block randomization (blocks of 20), but in Argentina the block-size was 10 Ns. The code was broken after the study was completed.	Previous immunosuppression			365	Visible blood-I stool. Massive gastrointestinal hemorrhage		None
Qazi, S.A., Khan, M.A.,	1996	Meningitis children	Dexamethasone	89	GI bleeding assessed by occult blood testing	Ampicillin, chloramphenicol (ceftriaxone), supportive care				Randomization: Unconstrained computer generated list of random treatment assignments.	Each if prior renal, hepatic or CNS disease.			7/5	GI bleeding assessed by occult blood testing		Routine
Sankar, J., Singh, P.,	2007	Meningitis children	Dexamethasone	252	GI bleeding	Ceftriaxone to all for a minimum of 7 days				A randomization list was prepared using simple random number table. Randomization codes were broken after the trial analysis was done. Duration of treatment not described.	Each if liver, renal or CNS disease			30	GI bleeding		Routine
Scarborough, M., Gordon, S.B.,	2007	Meningitis adults	Dexamethasone	465	Proven GI bleed	Ceftriaxone to all				Randomization was performed by computer-generated lists of eight in a two-way factorial design.	Each if ster received in the previous 48 hour.			40	Proven GI bleed		Systematic survey
Schaad, U. B., Lips, U.,	1993	Meningitis children	Dexamethasone	115	GI bleeding. Withdrawal due to bleeding in the upper gastrointestinal tract.	Ceftriaxone.	Additional medical therapy was provided according to standard procedures			Computer-generated list of random therapy assignments. The code not broken until the last pt had completed therapy.				7	Occult blood in stool. Overt GI bleeding, substantial falls in blood hemoglobin conc.		Routine
Thomas, R., Le Tutor, Y.,	1999	Meningitis adults	Dexamethasone	60	Gastric ulcer with overt hemorrhage	Amoxicillin				Randomization was stratified and equilibrated by center. The code was only broken at the end of the follow-up period.	Each if hypersensitivity to steroids.			30	Gastric ulcer with overt hemorrhage		Routine
Thwaites, G. E., Nguyen D.B.	2004	Tuberculous meningitis adults	Dexamethasone	545	GI bleeding defined as oral or rectal gastrointestinal bleeding on visual inspection	Isoniazid, rifampin, pyrazinamide, streptomycin, ethambutol (if HIV).				Computer-generated sequence of random numbers was used to allocate treatment in blocks of 30. All pts and staff remained blinded to the treatment allocation until the last pt completed follow-up	Each if the attending physician believed that steroids were contraindicated, if if the pt had received more than one dose of any steroid before study entry.			270	GI bleeding defined as oral or rectal gastrointestinal bleeding on visual inspection		Routine
Wald, E.R., Kaplan, S.L.,	1999	Meningitis children	Dexamethasone	143	Occult blood in stool and grossly bloody stools	Ceftriaxone. Anticonvulsants if needed.	Similar supportive care.			Computer-generated list of random therapy assignments in blocks of 10 for each participating institution. Investigators remained blinded to treatment assignment	Each if admin of steroids before enrollment in the study.			365	Occult blood in stool and grossly bloody stools		Routine
Bolbart, P. E., Charpentier, C	1999	Sepsis	Hydrocortisone	41	Gastrointestinal bleeding or endoscopic evidence of symptomatic gastric ulcerations	Vasopressors.	All concomitant treatments were accepted except NSAIDs and corticosteroids for 28 days.			Randomization was by computer-generated list in blocks of 8 and stratified by center by the hospital pharmacy.	Each if recent bleeding or endoduodenal ulcer, gastrointestinal bleeding, or corticosteroid treatment		5 % or more	28	GI bleeding endoscopic evidence of symptomatic gastric ulcerations		Routine
Briegel, J., Forst, H.,	1999	Septic shock	Hydrocortisone	40	Intestinal hemorrhage	Vasopressors, antibiotics				Randomization 1:1 ratio according to study center in blocs of four with the use of a computer-generated random-number generator list provided by statistician. All staff blinded etc.	Each if treatment with long-term steroids within the past 6 months or short-term steroids within the past 4 weeks.		1-Dec-6	14	Intestinal hemorrhage		Routine
Sprung, C. L., Annane, D.,	2008	Septic shock	Hydrocortisone	499	GI bleeding	Vasopressors, antibiotics?				A randomization table determined the order of inclusion for the pts to receive placebo among the expected 30 admissions. Eligible pts were prospectively randomized after the completion of randomization.	Each if recent GI hemorrhage, history of steroid use for over two weeks within the last year or upon admission to hospital.			28	GI bleeding		Routine
Sprung, C. L., Annane, D.,	2008	Septic shock	Hydrocortisone	499	GI bleeding defined as decrease hematocrit by 3%, with evidence of blood or coffee-ground material from upper GI tract. If tidal volume 12 hours after drug adm., considered complication of study drug.	Vasopressors, antibiotics, prophylactic antacids\cimetidine or both to all.				Randomization 1:1 ratio according to study center in blocs of four with the use of a computer-generated random-number generator list provided by statistician. All staff blinded etc.	Each if treatment with long-term steroids within the past 6 months or short-term steroids within the past 4 weeks.		5-11	28	GI bleeding		Routine
Sprung, C.L., Panagiota, V.C.,	1986	Septic shock	Methylprednisolone	59						Computer-generated randomization scheme. One or two doses given as intravenous infusion	None described			1-3			
Veterans Administration Systemic Sepsis Cooperative Study Group	1987	Sepsis	Methylprednisolone	223	GI bleeding	Antibiotics.	Additional therapeutic agents except naloxone, prostaglandin inh or steroid-containing compounds.			Within each hospital, pts were randomized to receive methylpred sod succinate or placebo. Randomization was performed by computer-generated list in blocks of 8	Each if glucocorticoid adm during the previous two weeks or a history of hypersensitivity to glucocorticoid adm.			5	GI bleeding		Routine
Vilde O. Dagsnæs M.,	2002	Sepsis	Prednisolone	40			Standard therapy for sepsis and septic shock: antibiotics, fluid replacement, vasoactive drugs, mechanical ventilation			Computer-generated randomization procedure. Pts and primary physicians were blinded as to which therapy was administered. No further description.	Each if use of steroids within the 3 months before the admission.		6-10	14	GI bleeding not defined		Routine
Svitila, F., Pehrsenn, O.,	2005	Bronchopulmonary dysplasia	Dexamethasone	109	Gastrointestinal bleeding and gastrointestinal perforations	Surfactant to 91% Open glucocorticoid therapy was common in both groups (66% in DX vs 73% in PL)				Randomization was performed blindly by the study investigators in the pharmacy. Open-label postnatal DX was allowed when deemed necessary by the attending physician, but its use was discouraged.			5-2	507	GI bleeding and GI perforations		Systematic survey
Romanta, F., Latsera, G.,	2005	Bronchopulmonary dysplasia	Prednisolone	50	GI perforation	Ibuprofen 21/25 (84%) steroid, 19/24 (79%) placebo. Surfactant.				Randomization: centralized, computer-generated, stratification into six risk groups. All infants with GI perforation received ibuprofen	Each if use of steroids before and during the 12 days of treatment.		5-12	28	GI perforation		Systematic survey
Lin, Y. J., Yeh, T. F.,	1998	RDS children	Dexamethasone	40		Surfactant, ampicillin, gentamicin				Randomization, achieved by use of a random number table, was stratified. Treatment allocation was recorded on cards enclosed in sequentially numbered envelo. Follow up until 36 weeks postmenstrual age.				28	GI bleeding not defined		Systematic survey
Brozanski, B. S., Jones, J.-G.,	1995	Bronchopulmonary dysplasia	Dexamethasone	76	Gastrointestinal hemorrhage	Diuretics, surfactant, bronchodilators, and methylxanthines				Trial entry and assignment by phone to CTSU for random allocation. The neonatologist could give open steroids at a later date if this was clinically indicated because of life-threatening deterioration.	Each if the clinician believed that the baby's condition was so poor that steroids were mandatory, or in mild that steroids were contraindicated.		0-49	252	GI bleeding		None
Collaborative Dexamethasone Trial Group	1993	Bronchopulmonary dysplasia	Dexamethasone	283	GI bleeding	Oxygen Treatment for patent ductus arteriosus (indomethacin), assisted ventilation, hospitalists, bronchodilators, diuretics.				The infants were assigned randomly to one of three treatment groups. Group selection was determined by sequential assignment from a table of random numbers. Two-way stratification by birth weight and center.			0-18-42	450	Gastric perforation		Systematic survey
Cummings, J. J., D'Eugenio, D. B.,	1989	Bronchopulmonary dysplasia	Dexamethasone	36	Gastric perforation. Gastric positive stools.					Randomization was by computer-generated, stratified, prerassigned treatment designations in sealed, opaque envelopes.			5-5	54	GI bleeding		Routine
Efird, M. M., Heerens, A. T.,	2001	Hypotension preterm infants	Hydrocortisone	34	Intestinal perforation	Vasopressors. Indomethacin to 2 HC, 6 placebo Indomethacin to many. Indomethacin treatment not randomly assigned.	Additional therapeutic agents except naloxone, prostaglandin inh or steroid-containing compounds.			Randomization by center within 4 strata (weight and medical risk ratios).	Each if glucocorticoid adm during the previous two weeks or a history of hypersensitivity to glucocorticoid adm.			14	Intestinal perforation		Routine
Garland, J.S., Alex, C.P.,	2004	Bronchopulmonary dysplasia	Dexamethasone	241	Gastrointestinal perforation. Gastric positive stools.	Surfactant to 91%. Open glucocorticoid therapy was common in both groups (66% in DX vs 73% in PL)				Randomization was performed blindly by the study investigators in the pharmacy. Open-label postnatal DX was allowed when deemed necessary by the attending physician, but its use was discouraged.			5-2	507	GI bleeding and GI perforations		Systematic survey
Kazzi, N. J., Bruns, T. W.	1999	Bronchopulmonary dysplasia	Prednisolone	53	GI perforation	Ibuprofen 21/25 (84%) steroid, 19/24 (79%) placebo. Surfactant.				Randomization: centralized, computer-generated, stratification into six risk groups. All infants with GI perforation received ibuprofen	Each if use of steroids before and during the 12 days of treatment.		5-12	28	GI perforation		Systematic survey
Lin, Y. J., Yeh, T. F.,	1998	RDS children	Dexamethasone	40		Surfactant, ampicillin, gentamicin				Randomization, achieved by use of a random number table, was stratified. Treatment allocation was recorded on cards enclosed in sequentially numbered envelo. Follow up until 36 weeks postmenstrual age.				28	GI bleeding not defined		Systematic survey
Ng, P. C., Lee, C. H.,	2004	Hypotension preterm infants	Hydrocortisone	68	Gastrointestinal perforation	Indomethacin 79.2%, vasopressor 100 % 7, dopamine, surfactant. VLBW infants who weighted <1250 g were given prophylactic indomethacin.				Random assignment in each participating center. Study drug/placebo prepared into identical syringes. 3 of the 4 with GI perforation had received indomethacin/ibuprofen	Each if postnatal systemic or inhaled corticosteroids before receiving the trial drug. Use of antenatal steroids were allowed.			5	Brown gastric aspirate (coffee ground). Gastrointestinal perforation?		Routine
Peltoniemi, O., Kari, A.,	2003	Bronchopulmonary dysplasia	Hydrocortisone	51	GI bleeding and GI perforations					Random assignment in each participating center. Study drug/placebo prepared into identical syringes. 3 of the 4 with GI perforation had received indomethacin/ibuprofen			5-10		GI bleeding and GI perforations		Systematic survey
Rastogi, A., Akintorin, S. M.,	1996	RDS children	Dexamethasone	76		Surfactant				The pharmacist consulted the appropriate assignment list, with "treatment" and "control" randomly ordered for that infant's weight stratum.			5-12	28	GI bleeding		Routine
Sanders, R. J., Cox, C.,	1994	Bronchopulmonary dysplasia	Dexamethasone	40		Surfactant, mechanical ventilation				Randomization: An order was sent to the pharmacy. Set of sealed envelopes. Investigators and care takers were blinded to the study drug.			5-3	120	GI bleeding not defined		Routine
Shinwell, E. S., Karplus, M.	1999	Bronchopulmonary dysplasia	Dexamethasone	248	Gastrointestinal haemorrhage	Surfactant				Phone to central randomization unit. Syringe sets were numbered according to a random number list. Randomization stratified by center and birthweight. Staff blinded.	Each if contraindications to steroids, such as bleeding tendency.		5-3	28	GI bleeding		Systematic survey
Sinkin, R. A., Dweck, H. S.,	2000	Bronchopulmonary dysplasia	Dexamethasone	366	Gastrointestinal perforation. Did not collect data on GI hemorrhage.	Surfactant				Randomized with stratification by center from a set of sealed envelopes in the pharmacy. Study drug/placebo prepared practice. Use of dex was at the discretion of the attending physician, not open-label steroid therapy.			5-12	365	GI perforation. Did not collect data on GI hemorrhage.		Systematic survey
Watterberg, K. L., Gerdes, J. S.,	1999	Chronic lung disease premature infants	Hydrocortisone	40	Upper gastrointestinal bleeding (a heme-positive gastric aspirate or emesis), gastrointestinal perforation	Indomethacin treatment to many but not randomly assigned. Surfactant.				Random permuted block algorithm. Stratified according to center and birth weight.	Each: prior postnatal treatment with a glucocorticoid. Use of any glucocorticoid therapy during hospitalization were recorded.		5-10		Upper GI bleeding (a heme-positive gastric aspirate or emesis), gastrointestinal perforation		Systematic survey
Kakela, J., Mosley, J. W.,	1991	Acute hepatic failure	Hydrocortisone	62	Upper GI bleeding that required blood transfusion					Pts were randomized by a computer program. The investigator had 3 options to have the pt randomized to any of the 3 groups or randomised to either hydrocortisone or assigned to one of the hydrocort gr				8	Upper GI bleeding that required blood transfusion		None
Sherman, S., Bilut, Lr.,	2005	Post-ERCP pancreatitis	Prednisone	1115		Surfactant. Group receiving indomethacin exc. in the present study but are available in the article.				Randomization: A computer-generated randomization was performed by computer for each participating center. Random sequences of drug or placebo were prepared by the pharmacy.	Excl: Pts with GI bleeding unless they could be effectively treated within 48 hours. Also gastric or duodenal ulcers and ulcerated esophagitis at endoscopy were excluded.			3	GI perforation. Upper or lower GI bleeding.		Routine
Soh, T. F., Lin, Y. J.,	1997	Bronchopulmonary dysplasia	Dexamethasone	262	Gastrointestinal haemorrhage	Ampicillin, gentamicin. Not surfactant.				Randomization: The numbers 1 through 270 were assigned at random either to the study group or to the control group.			5-28	28	GI bleeding		Systematic survey
Arabi, Y. M., Alyunuh, A	2010	Cirrhosis and septic shock	Hydrocortisone	75	GI bleeding	H2-blockers or PPIs for stress-ulcer prophylaxis. Vasopressors. Monopresen.				Pts were randomized by a central pharmacy using computer-generated random sequences. Sealed envelopes and based on computer-generated random allocation in blocks to one of two arms.	Each if prior systemic steroid usage or contraindications for systemic steroids		5-9	28	GI bleeding		Routine
Carithers, R. L., Jr., Herlong, H. F.,	1989	Alcoholic hepatitis	Methylprednisolone	66	GI bleeding	Spironolactone, B-complex multivitamins, folic acid, benzodiazepins				Randomization: Random sequences for drug or placebo were submitted to the supplm. A random code was prepared for each center, the code kept by an independent source.	Each if gastrointestinal hemorrhage requiring transfusions		5-42		GI bleeding		None
European Association for the Study of the Liver	1995	Chronic hepatitis B	Methylprednisolone	99	Hematemesis					Pts were allocated at a 1:1 ratio to a treatment plan. No further description.	Each if treatment with steroids for more than 4 weeks less than 3 mo before randomization, or if contraindications to steroid treatment		Essog time	454	Hematemesis		Routine
Kingsgaard, K., Marcellin, P.,	1990	Chronic hepatitis B	Methylprednisolone	288	GI bleeding	Interferon				Randomization: Central envelope procedure stratified according to sex, bleeding and cirrhosis	Each if contraindication for steroid therapy		5-28	28	GI bleeding		Systematic survey
Rakela, J., Mosley, J. W.,	1991	Acute hepatic failure	Hydrocortisone	62	Upper GI bleeding that required blood transfusion					Pts were randomized by a computer program. The investigator had 3 options to have the pt randomized to any of the 3 groups or randomised to either hydrocortisone or assigned to one of the hydrocort gr				8	Upper GI bleeding that required blood transfusion		None
Raymond, M. J., Poynard, T.,	1992	Alcoholic hepatitis	Methylprednisolone	81	GI bleeding	Spironolactone, B-complex multivitamins, folic acid, antacids				Randomization: Computer-generated numbers by pharmacy. A sequence. Concealed envelopes with active drug or placebo were dispensed.	Excl: Pts with GI bleeding unless they could be effectively treated within 48 hours. Also gastric or duodenal ulcers and ulcerated esophagitis at endoscopy were excluded.		5-28	180	GI bleeding		Routine
Aaronson, D., Kaiser, H.,	1996	Asthma	Prednisone	26		Albuterol sulfate				Randomization was carried out by a pharmacist.	Each if systemic to steroids, pts receiving other corticosteroids or who had received any form of corticosteroid therapy in the previous 6 months.		5-42	56	GI bleeding not defined	Patient checklist or interview	
Keane, D., Ment, L. s.,	1996	Proktocolitis children	Dexamethasone	66	Presence of blood routinely checked in gastric secretions and stool. Occult GI hemorrhage	Ranitidin				Randomization was carried out by a pharmacist. Pts were randomly assigned to receive dexa or placebo. Vials numbered sequentially and filled with steroid or placebo, were stored in each center in multiple blocks, with six sets to a block.	Each if steroids 7 days before inclusion Each if they had received corticosteroids of any kind or amount in the 48 hours before study screening the study or the presence of a disease process for which corticosteroids are required.		5-12	367	Presence of blood routinely checked in gastric secretions and stool. Occult GI hemorrhage		Routine
Bernard, G. R., Luce, J. M.,	1987	ARDS	Methylprednisolone	99		Mechanical ventilation				Randomization was performed in blocks of 4 pts at each study site. Pts were randomized in a 1:1 ratio.	Each if use of systemic corticosteroids within the preceding 3 weeks		5-4	45	Gastrointestinal bleeding		Routine
Buckingham, S. C., Jaffi, H. S.,	2003	RSV infection children	Dexamethasone	41		Intubation, mechanical ventilation.				Randomization was carried out by a computer-generated random-number table.	Each if regular corticosteroid medication before or during admission		5-3	7	GI bleeding		Routine
Confalonieri, M., Urbino, R.,	2005	Pneumonia	Methylprednisolone	86	Upper GI bleeding	Antibiotic therapy				Randomization schemes were generated in blocks of 10 for each participating ICUs by a central randomization center. Sealed envelopes.	Each if major gastrointestinal bleed within 3 months of the current hospitalization. Each if active GI bleeding event requiring transfusion		5-7	60	Upper GI bleeding		Routine
Cosser, Sc., Ferrara B.,	1992	Exfoliation failure, preterm babies	Dexamethasone	59		Methylxanthines, surfactant				Pts were randomly assigned to receive either dexa or sham treatment. The hospital pharmacy prepared the drugs. No further description.	Each if previously treated with DX		5-3	5	GI bleeding not defined		Routine
Elliott, A. M., Lucas, W.,	2004	HIV associated tuberculosis	Methylprednisolone	197		Ethambutol, isoniazid, rifampicin, pyrazinamide				The randomization sequence was generated by a statistician, by use of STATA. Randomization was done in blocks of 20. Staff gave pts the next number in the sequence in the order in which they were enrol	Each if they had risk factors for serious steroid-related Aes (a history of peptic ulcer disease). Inclusion if they had not recently received treatment with glucocorticoids.		0-56	1155	Gastric ulcers		Spontaneous reporting
Fanta, C. H., Rossing, T. H.,	1983	Asthma	Prednisone	20		Bronchodilator treatment: ith isoproterenol, epinephrine, methylxanthines, beta agents, aminophylline, terbutaline, isoetharine				Randomization was based on a one-on-one allocation. The allocation sequence was computer generated and was kept i a safe at the hospital pharmacy.	Each if currently taking oral corticosteroids		0-3		GI bleeding not defined	Patient checklist or interview	
Fernandez-Serrano, S., Dorca, J.,	2011	Pneumonia	Prednisone	213		Ceftriaxone and levofloxacin to all. Omeprazole				Randomization was based on a one-on-one allocation. The allocation sequence was computer generated and was kept i a safe at the hospital pharmacy.	Each if use of prednisone 15 mg or more for more than 24 hours or any condition requiring corticosteroids.		0-7	90	GI bleeding not defined		Routine
Sorkness, C. A., Lafornz C	1999	Asthma	Prednisone	58		Beta agonists, theophylline,cromolyn sodium.				Randomization was performed by the random numbers method in a 1:1 ratio by the pharmacy.	Each: Receipt of any corticosteroid therapy during the 3 months before visit 1; use of >240 mg prednisone or its equivalent in any dosage form or regimen during the 3 months before visit 1; use of inhaled steroids; historical or current evidence of significant peripheral disease		0-28	28	GI bleeding not defined		Routine
Walmsley, S., Levinton, C.,	1995	Pneumonia / AIDS	Methylprednisolone	78	Lower GI bleeding	Trimetoprim-sulfamethoxazole, pentamidine, dapsone, or eflornithine. Use of steroids outside the study protocol were discouraged.				Randomization was performed by the random numbers method in a 1:1 ratio by the pharmacy.	Each if active gastrointestinal bleeding in the previous 3 months or receiving systemic steroids in the 3 months before randomization		5-10	180	Lower GI bleeding		Systematic survey
Weigelt, J. A., Norcross, J. F.,	1985	Respiratory failure	Methylprednisolone	81		Pluid administration, crystalloid solutions, blood, colloids, antibiotics were used as indicated				Randomization was according to a computer-generated random-number table.	Each if on long-term steroid therapy		5-3	168	GI bleeding not defined		Routine
Weiss, C., Welz, A.,	1982	Tuberculous pleurisy	Prednisone	41		Standard treatment: bekä-agonists, ipratropium bromide, oral antibiotics, supplemental oxygen. Methylxanthines.				Pts randomly assigned in a double-blind fashion. No further description.	Each if previous or current oral steroid treatment, contraindication to parenteral steroids (recent gastric ulcer perforation or bleed).		0-730	730	GI bleeding not defined		Routine
Choy, E. H., Kingsley, G. H.,	2005	RA	Dexamethasone	91		NSAIDs, analgesics, DMARD (gold, penicillamine, mtx, azathioprine or cyclosporin).				Pts were randomly assigned to receive steroid or placebo. Injection were given by nurses not involved in assessing pts.	Each if previous or current oral steroid treatment, contraindication to parenteral steroids (recent gastric ulcer perforation or bleed).		0-730	730	GI bleeding not defined		Routine
Kirwan, J. R.	1995	RA	Prednisone	126		Any other treatment, except systemic steroids. NSAIDs, gold, penicillamine, sulfasalazine. methotrexate. other.				Randomization was performed in blocks of six subjects at each center, and except in an emergency, the randomization codes were not broken until the main analysis.	Each if unstable conditions that might respond to prednisone, recent peptic ulcerations, or regular use of anti-inflammatory medications within the previ 2 mo.		0-730	730	GI bleeding not defined		Routine
Kirwan, J. R., Hallgren, R.,	2004	RA	Prednisone	70		No treatment with or a stable dose of NSAIDs or analgesics/DMARDs for at least 30/90 days				Drugs were dispensed by the pharmacy in relation to the study number and in accordandwith a predefined sequence of randomly generated allocations kept in sealed envelopes.	Each if steroid 30 days before inclusion, active peptic ulcer		0-84	112	GI bleeding not defined		Routine
Van Everdingen, A. A., Jacobs, J. W.,	2002	RA	Prednisone	81	Gastrointestinal bleeding,ulcers, and peptic symptoms. PUD with bleeding confirmed by gastroscopy	NSAIDs, paracetamol, sulfasalazine, intra-articular steroid injections to SLNd and and LEJd1 plus. The overall use of NSAIDs was considerably lower in the prednisone group than i the placebo group.				Computer-generated randomization procedure in blocks of 10. Only pharmacist could access the allocation table.	Each if contraindications to prednisone or NSAIDs, active gastrointestinal problems, hemorrhagic diathesis		0-730	730	GI bleeding, ulcers, and peptic symptoms. PUD with bleeding confirmed by gastroscopy		Routine
Wassenberg, S., Rau, R.,	2005	RA	Prednisone	150	Gastric ulcer, but no bleeding described	Gold or MTX to all. NSAIDs/osteoporosis prophylaxis/nitrogen therapy permitted.				Randomization was carried out across all study centers on the basis of a randomization plan drawn up by Merck using the RAPHSIS computer program.	Each if glucocorticoidsdependent disease such as asthma, previous oral glucocorticoidtreatment (<3 months).		0-730	730	Gastric ulcer, but no bleeding described		Systematic survey
Adams, M., Soskop, M.	1995	Chronic	Methylprednisolone	126	Hematemesis	Tropisetron and chemotherapy to all				Randomized, no further description of procedure	Each if previous peptic ulcer or adverse reactions to steroids or taking other corticosteroid medication		0-5		Hematemesis		Systematic survey
Kim, M. J., Lee, J.,	1994	Chronic	Methylprednisolone	45		Cisplatin, ondansetron, levamisole				Randomized, no further description of procedure			5-5	5	GI bleeding not defined		Routine
Kraan, F. S., Davis, K. L.,	2000	Alzheimer's disease	Prednisone	138						Drugs were assembled into identical containers with coded labels. Codes were sealed at the packaging center, the randomization scheme was approved only by the statistical core.	Each if comorbid conditions that might respond to prednisone, recent peptic ulcerations, or regular use of anti-inflammatory medications within the previ 2 mo.		0-477	365	GI bleeding not defined		Routine
Al-Shehri, A. M	2004	Tonsillectomy	Dexamethasone	30		Postoperative analgesics (2000 mg paracetamol per day), no additional medication.				Randomly assigned to treatments according to a double-blind randomization.	Each if contraindications to steroid use such as gastrointestinal ulcer		0-5	14	GI bleeding not defined		Spontaneous reporting
Arana, G. W., Santos, A. B	1985	Depression	Dexamethasone	37						Randomly assigned. Pls, investigators, and the research assistent were blind to the treatment assignment during the study period. No further description.	Each if current treatment with steroids.		0-56	14	GI bleeding not defined	Patient checklist or interview	
Austin, J. R., Pesland, S. P.	1993	Facial nerve paralysis	Prednisone	76						The randomization was performed by the pharmacy under the auspices of the clinical director of pharmacological research.	Each if contraindications to steroid use such as peptic ulcer disease.		0-10	180	GI bleeding not defined		Routine
Azuara-Blanco, A., Spaeth, G. L.,	1999	Glaucoma	Prednisone	30		Regional anesthesia, mitomycin C, 5-fluorouracil, topical antibiotic and cycloplegic				Randomization according to Moses-Oakford blocked assignment algorithm.	Excl: active peptic ulcer disease.		0-3	240	GI bleeding not defined		Routine
Barrilleaux, P. S., Martin, J. N	2007	Preeclamptic patients	Dexamethasone	157		Mg, antihypertensive therapy.				Randomization via computer-generated randomization sequence via personnel in pharmacy research group. Infusion sets identical for treatment and control patients.	Each if recent or concurrent use of steroids		0-3	7	GI bleeding not defined		Routine
Berglund, F.-I., Sjoberg, V.	1958	Thrombosis	Prednisone	112	Upper GI discomfort reported, not included.	Standard ant