

**Table S3.** Demographic characteristics and clinical diagnosis (per Reference Clinical Diagnosis) by study – PP population (N = 622)

		Study				
		<b>DP008-003 (N = 157)</b>	<b>PDT304 (N = 100)</b>	<b>PDT301 (N = 288)</b>	<b>PDT408 (N=77)</b>	<b>Total (N = 622)</b>
<b>Age (yr)</b>	Mean (SD)	63.1 (8.51)	60.5 (10.97)	74.2 (7.02)	64.1 (12.05)	67.9 (10.61)
	Min, Max	40, 80	33, 79	54, 90	25, 84	25, 90
	Median	64.0	61.5	75.0	67.0	69.0
<b>Gender</b>	Male	99 (63%)	57 (57%)	160 (56%)	40 (52%)	356 (57%)
	Female	58 (37%)	43 (43%)	128 (44%)	37 (48%)	266 (43%)
<b>Race</b>	Caucasian	153 (97%)	100 (100%)	288 (100%)	76 (99%)	617 (99%)
	Black	3 (2%)	0 (0%)	0 (0%)	0 (0%)	3 (<1%)
	Asian	1 (1%)	0 (0%)	0 (0%)	1 (1%)	2 (<1%)
	Other	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
<b>PS (SDDD)</b>		115 (73%)	69 (69%)	0 (0%)	47 (61%)	231 (37%)
<b>Possible PS</b>		115 (73%)	5 (5%)	0 (0%)	47 (61%)	167 (27%)
<b>Probable PS</b>		0 (0%)	64 (64%)	0 (0%)	0 (0%)	64 (10%)
<b>DLB (SDDD)</b>		0 (0%)	0 (0%)	110 (38%)	0 (0%)	110 (18%)
<b>Possible DLB</b>		0 (0%)	0 (0%)	25 (9%)	0 (0%)	25 (4%)
<b>Probable DLB</b>		0 (0%)	0 (0%)	85 (30%)	0 (0%)	85 (14%)
<b>Non-PS/Non-DLB (no SDDD)</b>		42 (27%)	31 (31%)	123 (43%)	30 (39%)	226 (36%)
<b>ET</b>		16 (10%)	14 (14%)	0 (0%)	23 (30%)	53 (9%)
<b>AD</b>		0 (0%)	0 (0%)	122 (42%)	0 (0%)	122 (20%)
<b>Other</b>		26 (17%)	17 (17%)	1 (<1%)	7 (9%)	51 (8%)
<b>SDDD Present<sup>a</sup></b>		115 (73%)	69 (69%)	110 (38%)	47 (61%)	341 (55%)
<b>SDDD Absent</b>		42 (27%)	31 (31%)	123 (43%)	30 (39%)	226 (36%)

<sup>a</sup>Includes Possible and Probable PS and Possible and Probable DLB diagnoses.

AD = Alzheimer's disease; DLB = Dementia with Lewy bodies; ET = Essential tremor; N = number of subjects in the study; PP = Per protocol; PS = Parkinsonian syndrome; SD = standard deviation; SDDD = striatal dopaminergic deficit disorder.