

**Title: Cohort profile: COVID-19 in a cohort of pregnant women and their descendants, the MOACC-19 study**

Authors: Javier Llorca<sup>1,2</sup>, Carolina Lechosa-Muñiz<sup>1,3</sup>, Pilar Gortázar<sup>1,3</sup>, María Fernández-Ortiz<sup>4</sup>, Yolanda Jubete<sup>1,3</sup>, María J. Cabero<sup>1,3,4</sup> and the MOACC-19 group

## Affiliations:

1. Universidad de Cantabria, Santander, Spain
2. CIBER de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain
3. University Hospital Marques de Valdecilla, Santander, Spain
4. IDIVAL, Santander, Spain

## Correspondence:

Javier Llorca  
Facultad de Medicina  
Universidad de Cantabria  
Avda. Herrera Oria s/n  
39011 Santander  
Spain  
Phone # 34-942 201 993  
E-mail: [javier.llorca@unican.es](mailto:javier.llorca@unican.es)

## Members of the MOACC-19 group:

Jéssica Alonso-Molero<sup>1,4</sup>, Bárbara Arozamena<sup>3</sup>, Laura Conde-Gil<sup>3</sup>, Elsa Cornejo del Río<sup>3</sup>, Rocío Cuesta-González<sup>3</sup>, Trinidad Dierssen-Sotos<sup>1,2,4</sup>, Pelayo Frank de Zulueta<sup>3</sup>, Inés Gómez-Acebo<sup>1,2,4</sup>, Coral Llano-Ruiz<sup>3</sup>, Lorena Lasarte-Oria<sup>3</sup>, Sonia López-Gómez<sup>3</sup>, Sonia Mateo-Sota<sup>3</sup>, Victoria Orallo<sup>3</sup>, Rosa Pardo<sup>3</sup>, Daniel Pérez González<sup>3</sup>, María Sáez de Adana Herrero<sup>3</sup>.

Supplementary Table 1. Characteristics of women tested positive for SARS-CoV-2 infection via RT-PCR, subcohorts 1 and 2

Variable	Category	N (%)
<b>Study cohort</b>	Sub-cohort 1 (Women delivering before 26/May/2020)	5
	Sub-cohort 2 (Women delivering from 26/May/2020 on)	9
<b>Partner RT-PCR</b>	Negative	7
	Positive	7
<b>Age</b>	<25 years	1 (7.1)
	25-29 years	3 (21.4)
	30-34 years	4 (28.6)
	35-39 years	4 (28.6)
	≥40 years	2 (14.3)
<b>Education level</b>	Primary studies	3 (21.4)
	Secondary studies	2 (14.3)
	Vocational training	2 (14.3)
	University studies	7 (50.0)
<b>Work status</b>	Unemployed / no active worker	5 (35.7)
	Employed	9 (64.3)
	Student	0
<b>Occupation</b>	Health care worker	4 (28.6)
	Working in restaurant or commerce	2 (14.3)
	Others	3 (21.4)
<b>Fertilization type</b>	Natural	12 (92.3)
	Artificial insemination	1 (7.7)
	In vitro fertilization (own ovule)	0
	In vitro fertilization (ovodonation)	0
	Missing	1 (7.7)
<b>Smoker in pregnancy</b>	No	14 (100.0)
	Yes	0
<b>Alcohol consumption in pregnancy</b>	No	14 (100.0)
	Yes	0
<b>Parity</b>	1	9 (64.3)
	2	1 /7.1)
	≥3	4 (28.6)
<b>Type of delivery</b>	Eutocic	8 (57.1)
	Caesarean rate	4 (28.6)
	Instrumental	2 (14.3)
<b>Age (years)</b>	mean±sd (min, max)	32.9±5.1 (23 – 40)
<b>Gestational age leave work (weeks)</b>	mean±sd (min, max)	23.4±12.7 (0 – 36)
<b>Pregestational BMI (kg/m<sup>2</sup>)</b>	mean±sd (min, max)	23.3±4.0 (18.6 – 32.0)

<b>Gestational weight gain (kg)</b>	mean±sd (min, max)	11.2±4.1 (8 – 22)
<b>Symptoms associated with COVID-19</b>	Fever	3
	Chills	3
	Intense tiredness	2
	Sore throat	3
	Cough	3
	Short of breathing	2
	Headache	3
	Nausea	1
	Loss of sense	2

Supplementary Table 2. Characteristics of children born from women tested positive to SARS-CoV-2 infection using RT-PCR

Variable	Category	N = 7 (%)
Gestational age at birth	<34 weeks	0
	34-36 weeks	0
	≥37 weeks	14 (100.0)
Birth weight	<2500 g	0
	2500-4000 g	14 (100.0)
	>4000 g	0
Gender	Women	4 (28.6)
	Men	10 (71.4)
Twin or singleton	Singleton	14 (100.0)
	Twin	0
Breastfeeding at hospital discharge	Exclusive breastfeeding	4 (28.6)
	Mixed	6 (42.9)
	Infant formula	4 (28.6)
Neonatology admission	No	11 (78.6)
	Yes	3 (21.4)
Birth size	mean±sd (min, max)	50.0±2.4 (47 – 53)
Birth cephalic perimeter	mean±sd (min, max)	34.7±1.5 (33 – 39)
Apgar 1'	mean±sd (min, max)	8.4±1.4 (4 – 9)
Apgar 5'	mean±sd (min, max)	9.2±0.9 (7 – 10)
pH at birth	mean±sd (min, max)	7.27±0.05 (7.20 – 7.36)
SARS-CoV-2 via RT-PCR		All negative