

Appendix 5: All advice provided to patients during all registered contacts, and separately for patients suspected of COVID-19.

All countries	Advice for home isolation			
	<i>All patients</i>	67.2% (65.5-68.8)		
	<i>COVID-19 suspected</i>	83.2% (81.1-85.1)		
	Advice for symptomatic treatment			
	<i>All patients</i>	75.6% (74-77)		
	<i>COVID-19 suspected</i>	70.4% (67.9-72.8)		
	A scheduled follow-up visit/call			
	<i>All patients</i>	49.9% (48.1-51.6)		
	<i>COVID-19 suspected</i>	51.5% (48.8-54.2)		
	Advice for family members		Home isolation	
<i>All patients</i>	51.9% (50.1-53.7)	<i>All patients</i>	34.0% (32.3-35.7)	
<i>COVID-19 suspected</i>	69.6% (67.1-72.1)	<i>COVID-19 suspected</i>	54.7% (52-57.4)	
		Social distancing		
		<i>All patients</i>	40.2% (38.5-41.9)	
		<i>COVID-19 suspected</i>	53.1% (52-57.4)	
Preventive measures for patient		Extra handwashing		
<i>All patients</i>	64.1% (62.4-65.8)	<i>All patients</i>	56.7% (55-58.5)	
<i>COVID-19 suspected</i>	68.6% (66.1-71.1)	<i>COVID-19 suspected</i>	62.4% (59.8-65)	
		Sneezing in sleeve		
		<i>All patients</i>	47.2% (45.5-49)	
		<i>COVID-19 suspected</i>	48.4% (45.7-51.1)	
		Social distancing		
		<i>All patients</i>	51.1% (49.3-52.9)	
		<i>COVID-19 suspected</i>	56.3% (53.6-59)	
		Nose/mouth protection		
		<i>All patients</i>	37.3% (35.6-39)	
		<i>COVID-19 suspected</i>	42.2% (39.5-44.9)	
		Staying in separate room		
		<i>All patients</i>	27.3% (25.8-28.9)	
		<i>COVID-19 suspected</i>	40.6% (37.9-43.3)	
Where to find reliable information				
<i>All patients</i>	20.0% (18.6-21.5)			
<i>COVID-19 suspected</i>	28.0% (25.6-30.5)			
Armenia	Advice for home isolation			
	<i>All patients</i>	47.6% (40.3-55.1)		
	<i>COVID-19 suspected</i>	64.9% (25.6-30.5)		
	Advice for symptomatic treatment			
	<i>All patients</i>	81.2% (74.6-86.3)		
	<i>COVID-19 suspected</i>	83.3% (75.4-89.1)		
	A scheduled follow-up visit/call			
	<i>All patients</i>	49.4% (42-56.8)		
	<i>COVID-19 suspected</i>	46.5% (37.6-55.6)		
	Advice for family members		Home isolation	
<i>All patients</i>	47.6% (40.3-55.1)	<i>All patients</i>	44.1% (36.9-51.6)	
<i>COVID-19 suspected</i>	62.3% (53.1-70.6)	<i>COVID-19 suspected</i>	58.8% (49.6-67.4)	
		Social distancing		
		<i>All patients</i>	9.4% (5.9-14.7)	
		<i>COVID-19 suspected</i>	11.4% (6.8-18.5)	

	Preventive measures for patient <i>All patients</i>	44.1% (36.9-51.6)	Extra handwashing <i>All patients</i>	42.4% (35.2-49.9)	
	<i>COVID-19 suspected</i>	55.3% (46.1-64.1)	<i>COVID-19 suspected</i>	53.5% (44.462.4)	
			Sneezing in sleeve <i>All patients</i>	5.9% (3.2-10.5)	
			<i>COVID-19 suspected</i>	7.0% (3.6-13.2)	
			Social distancing <i>All patients</i>	6.5% (3.7-11.2)	
			<i>COVID-19 suspected</i>	7.0% (3.6-13.2)	
			Nose/mouth protection <i>All patients</i>	5.9% (3.2-10.5)	
		<i>COVID-19 suspected</i>	7.9% (4.2-14.3)		
		Staying in separate room <i>All patients</i>	2.9% (1.3-6.7)		
		<i>COVID-19 suspected</i>	4.4% (1.9-9.9)		
	Where to find reliable information <i>All patients</i>	2.4% (1-5.9)			
	<i>COVID-19 suspected</i>	1.8% (0.5-6.2)			
Belgium	Advice for home isolation <i>All patients</i>	90.7% (86.1-93.9)			
	<i>COVID-19 suspected</i>	89.0% (82.9-93.1)			
	Advice for symptomatic treatment <i>All patients</i>	71.3% (64.9-76.9)			
	<i>COVID-19 suspected</i>	69.2% (61.3-76.1)			
	A scheduled follow-up visit/call <i>All patients</i>	30.1% (24.4-36.5)			
	<i>COVID-19 suspected</i>	34.9% (27.7-43)			
	Advice for family members <i>All patients</i>	50.0% (43.3-56.6)	Home isolation <i>All patients</i>	22.2% (17.2-28.2)	
			<i>COVID-19 suspected</i>	18.5% (13-25.6)	
	<i>COVID-19 suspected</i>	50.7% (42.7-58.7)	Social distancing <i>All patients</i>	42.1% (35.7-48.8)	
			<i>COVID-19 suspected</i>	42.5% (34.7-50.6)	
	Preventive measures for patient <i>All patients</i>	31.5% (25.7-38)	Extra handwashing <i>All patients</i>	23.6% (18.4-29.7)	
			<i>COVID-19 suspected</i>	19.9% (14.2-27.1)	
		<i>COVID-19 suspected</i>	26.7% (20.2-34.4)	Sneezing in sleeve <i>All patients</i>	18.1% (13.5-23.7)
				<i>COVID-19 suspected</i>	13.7% (9.1-20.2)
			Social distancing <i>All patients</i>	30.6% (24.8-37)	
			<i>COVID-19 suspected</i>	25.3% (19-33)	
			Nose/mouth protection <i>All patients</i>	6.0% (3.6-10)	
		<i>COVID-19 suspected</i>	5.5% (2.8-10.4)		
		Staying in separate room <i>All patients</i>	12.5% (8.7-17.6)		
		<i>COVID-19 suspected</i>	11.0% (6.9-17.1)		
	Where to find reliable information <i>All patients</i>	15.3% (11.1-20.7)			
	<i>COVID-19 suspected</i>	14.4% (9.6-20.1)			

Germany	Advice for home isolation			
	<i>All patients</i>		19.6% (14.6-25.8)	
	<i>COVID-19 suspected</i>		88.2% (73.4-95.3)	
	Advice for symptomatic treatment			
	<i>All patients</i>		96.3% (92.6-98.2)	
	<i>COVID-19 suspected</i>		100% (89.9-100)	
	A scheduled follow-up visit/call			
	<i>All patients</i>		91.0% (86.1-94.3)	
	<i>COVID-19 suspected</i>		94.1% (80.9-98.4)	
	Advice for family members			
<i>All patients</i>	9.5% (6.1-14.6)	Home isolation	<i>All patients</i>	6.3% (3.7-10.8)
<i>COVID-19 suspected</i>	52.9% (36.7-68.6)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	35.3% (21.5-52.1)
		Social distancing	<i>All patients</i>	7.4% (4.5-12.1)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	41.2% (26.4-57.8)
Preventive measures for patient		Extra handwashing	<i>All patients</i>	39.7% (33-46.8)
<i>All patients</i>	48.7% (41.7-55.8)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	97.1% (85.1-99.5)
<i>COVID-19 suspected</i>	97.1% (85.1-99.5)	Sneezing in sleeve	<i>All patients</i>	40.2% (33.5-47.3)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	91.2% (77-97)
		Social distancing	<i>All patients</i>	37.0% (30.5-44.1)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	94.1% (80.9-98.4)
		Nose/mouth protection	<i>All patients</i>	22.8% (17.4-29.2)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	85.3% (70-93.6)
		Staying in separate room	<i>All patients</i>	0.0% (0-2)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	0.0% (0-10)
Where to find reliable information				
<i>All patients</i>				10.6% (7-15.8)
<i>COVID-19 suspected</i>				58.8% (42.2-73.6)
Denmark	Advice for home isolation			
	<i>All patients</i>		88.2% (82.1-92.4)	
	<i>COVID-19 suspected</i>		92.7% (86.3-96.3)	
	Advice for symptomatic treatment			
	<i>All patients</i>		46.1% (38.3-54)	
	<i>COVID-19 suspected</i>		44.5% (35.6-53.9)	
	A scheduled follow-up visit/call			
	<i>All patients</i>		18.4% (13.1-25.3)	
	<i>COVID-19 suspected</i>		17.3% (11.4-25.4)	
	Advice for family members			
<i>All patients</i>	24.3% (18.2-31.7)	Home isolation	<i>All patients</i>	13.8% (9.2-20.2)
<i>COVID-19 suspected</i>	23.6% (16.7-32.4)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	14.5% (9.2-22.3)
		Social distancing	<i>All patients</i>	13.8% (9.2-20.2)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	10.9% (6.4-8.1)
Preventive measures for patient		Extra handwashing	<i>All patients</i>	21.7% (15.9-28.9)
<i>All patients</i>	37.5% (30.2-45.4)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	19.1% (12.8-27.4)

	<i>COVID-19 suspected</i>	35.5% (27.1-44.8)	Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	18.4% (13.1-25.4) 18.2% (12.1-26.4)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	23.0% (17.1-30.3) 20.9% (14.4-29.4)
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	2.0% (0.7-5.6) 2.7% (0.9-7.7)
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	0.7% (0.1-3.6) 0.9% (0.2-4.9)
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	7.9% (4.6-13.3) 7.3% (3.7-13.7)		

Spain	Advice for home isolation <i>All patients</i> <i>COVID-19 suspected</i>	81.2% (75.5-85.8) 89.4% (84.1-93)		
	Advice for symptomatic treatment <i>All patients</i> <i>COVID-19 suspected</i>	84.8% (79.5-88.9) 84.0% (78.1-88.6)		
	A scheduled follow-up visit/call <i>All patients</i> <i>COVID-19 suspected</i>	74.4% (68.3-79.7) 81.9% (75.8-86.8)		
	Advice for family members <i>PPAS COVID</i> <i>COVID-19 suspected patients</i>	69.5% (63.2-75.2) 78.7% (72.3-84)	Home isolation <i>All patients</i> <i>COVID-19 suspected</i>	65.5% (59-71.4) 73.9% (67.2-79.7)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	51.1% (44.6-57.6) 57.4% (50.3-64.3)
	Preventive measures for patient <i>All patients</i> <i>COVID-19 suspected</i>	80.7% (75-85.4) 87.2% (81.7-91.3)	Extra handwashing <i>All patients</i> <i>COVID-19 suspected</i>	78.0% (72.1-83) 84.0% (78.1-88.6)
			Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	39.5% (33.3-46) 43.1% (36.2-50.2)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	58.3% (51.7-64.6) 63.8% (56.8-70.4)
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	56.5% (50-62.8) 60.6% (53.5-67.3)
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	62.3% (55.8-68.4) 70.2% (63.3-76.3)
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	20.6% (15.8-26.4) 23.9% (18.4-30.5)		

France	Advice for home isolation			
	<i>All patients</i>	79.4%	(72.3-85)	
	<i>COVID-19 suspected</i>	89.4%	(81.5-94.1)	
	Advice for symptomatic treatment			
	<i>All patients</i>	80.0%	(73-85.5)	
	<i>COVID-19 suspected</i>	81.9%	(72.9-88.4)	
	A scheduled follow-up visit/call			
	<i>All patients</i>	47.7%	(40-55.6)	
	<i>COVID-19 suspected</i>	61.7%	(51.6-70.9)	
	Advice for family members			
<i>All patients</i>	64.5%	(56.7-71.6)	Home isolation	
<i>COVID-19 suspected</i>	72.3%	(62.6-80.4)	<i>All patients</i>	51.0%
			<i>COVID-19 suspected</i>	57.4%
			Social distancing	
			<i>All patients</i>	54.8%
			<i>COVID-19 suspected</i>	61.7%
Preventive measures for patient			Extra handwashing	
<i>All patients</i>	80.6%	(73.7-86.1)	<i>All patients</i>	74.2%
<i>COVID-19 suspected</i>	84.0%	(75.3-90.1)	<i>COVID-19 suspected</i>	80.9%
			Sneezing in sleeve	
			<i>All patients</i>	63.9%
			<i>COVID-19 suspected</i>	73.4%
			Social distancing	
			<i>All patients</i>	76.8%
			<i>COVID-19 suspected</i>	81.9%
			Nose/mouth protection	
			<i>All patients</i>	41.3%
			<i>COVID-19 suspected</i>	44.7%
			Staying in separate room	
			<i>All patients</i>	40.6%
			<i>COVID-19 suspected</i>	56.4%
Where to find reliable information				
<i>All patients</i>	9.0%	(5.5-14.6)		
<i>COVID-19 suspected</i>	11.7%	(6.7-19.7)		
Georgia*	Advice for home isolation			
	<i>All patients</i>	86.9%	(81.9-90.7)	
	<i>COVID-19 suspected</i>	100%		
	Advice for symptomatic treatment			
	<i>All patients</i>	82.9%	(77.4-87.3)	
	<i>COVID-19 suspected</i>	88.9%		
	A scheduled follow-up visit/call			
	<i>All patients</i>	49.5%	(43-56.1)	
	<i>COVID-19 suspected</i>	55.6%		
	Advice for family members			
<i>All patients</i>	58.6%	(52-64.8)	Home isolation	
<i>COVID-19 suspected</i>	88.9%		<i>All patients</i>	7.7%
			<i>COVID-19 suspected</i>	22.2%
			Social distancing	
			<i>All patients</i>	54.5%
			<i>COVID-19 suspected</i>	88.9%
Preventive measures for patient			Extra handwashing	
<i>All patients</i>	93.2%	(89.2-95.9)	<i>All patients</i>	73.9%
			<i>COVID-19 suspected</i>	66.7%

	<i>COVID-19 suspected</i>	100%	Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	73.0% (66.8-78.4) 55.6%
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	89.6% (84.9-93) 88.9%
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	72.5% (66.3-78) 66.7%
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	10.8% (7.4-15.6) 55.6%
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	11.3% (7.8-16.1) 0.0%		

Greece	Advice for home isolation <i>All patients</i> <i>COVID-19 suspected</i>	75.6% (69.5-80.8) 83.1% (73.7-89.7)		
	Advice for symptomatic treatment <i>All patients</i> <i>COVID-19 suspected</i>	63.8% (57.3-70) 50.6% (40.1-61.1)		
	A scheduled follow-up visit/call <i>All patients</i> <i>COVID-19 suspected</i>	71.0% (64.7-76.6) 77.1% (67-84.8)		
	Advice for family members <i>All patients</i> <i>COVID-19 suspected</i>	73.3% (67.1-78.7) 96.4% (89.8-98.7)	Home isolation <i>All patients</i> <i>COVID-19 suspected</i>	51.0% (44.6-57.6) 74.7% (64.4-82.8)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	62.4% (55.9-68.6) 94.0% (86.7-97.4)
	Preventive measures for patient <i>PPAS COVID</i> <i>COVID-19 suspected patients</i>	82.4% (76.8-86.8) 92.8% (85.1-96.6)	Extra handwashing <i>All patients</i> <i>COVID-19 suspected</i>	72.9% (66.6-78.3) 91.6% (83.6-95.8)
			Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	71.9% (65.7-77.5) 91.6% (83.6-95.8)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	71.0% (64.7-76.6) 91.6% (83.6-95.8)
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	63.3% (56.8-69.4) 89.2% (80.7-94.2)
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	48.0% (41.5-54.5) 75.9% (65.7-83.8)
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	0.0% (0-1.7) 0.0% (0-4.4)		

Hungary*	Advice for home isolation			
	<i>All patients</i>		45.7% (37.6-54)	
	<i>COVID-19 suspected</i>		83.3%	
	Advice for symptomatic treatment			
	<i>All patients</i>		55.1% (46.8-63.1)	
	<i>COVID-19 suspected</i>		33.3%	
	A scheduled follow-up visit/call			
	<i>All patients</i>		39.1% (31.4-47.5)	
	<i>COVID-19 suspected</i>		0.0%	
	Advice for family members			
<i>All patients</i>	42.8% (34.8-51.1)	Home isolation	<i>All patients</i>	32.6% (25.4-40.8)
<i>COVID-19 suspected</i>	83.3%	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	66.7%
		Social distancing	<i>All patients</i>	35.5% (28-43.8)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	55.6%
Preventive measures for patient		Extra handwashing	<i>All patients</i>	58.0% (49.6-65.9)
<i>All patients</i>	61.6% (53.3-69.3)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	88.9%
<i>COVID-19 suspected</i>	100%	Sneezing in sleeve	<i>All patients</i>	48.6% (40.4-56.8)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	77.8%
		Social distancing	<i>All patients</i>	25.4% (18.8-33.2)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	72.2%
		Nose/mouth protection	<i>All patients</i>	25.4% (18.8-33.2)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	66.7%
		Staying in separate room	<i>All patients</i>	11.6% (7.3-18)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	55.6%
Where to find reliable information				
<i>All patients</i>		0.0% (0-2.7)		
<i>COVID-19 suspected</i>		0.0%		
Ireland	Advice for home isolation			
	<i>All patients</i>		76.5% (70.4-81.7)	
	<i>COVID-19 suspected</i>		91.7% (83.8-95.9)	
	Advice for symptomatic treatment			
	<i>All patients</i>		85.7% (80.4-89.8)	
	<i>COVID-19 suspected</i>		86.9% (78.1-92.5)	
	A scheduled follow-up visit/call			
	<i>All patients</i>		10.6% (7.2-15.4)	
	<i>COVID-19 suspected</i>		14.3% (8.4-23.3)	
	Advice for family members			
<i>All patients</i>	65.9% (59.4-71.9)	Home isolation	<i>All patients</i>	52.5% (45.9-59.1)
<i>COVID-19 suspected</i>	81.0% (71.3-87.9)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	70.2% (59.8-79)
		Social distancing	<i>All patients</i>	44.7% (38.2-51.4)
		<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	57.1% (46.5-67.2)
Preventive measures for patient		Extra handwashing	<i>All patients</i>	58.1% (51.4-64.4)
<i>All patients</i>	67.7% (61.3-73.6)	<i>COVID-19 suspected</i>	<i>COVID-19 suspected</i>	75.0% (64.8-83)

	<i>COVID-19 suspected</i>	88.1% (79.5-93.4)	Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	46.5% (40-53.2) 60.7% (50-70.5)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	56.2% (49.6-62.7) 73.8% (63.5-82)
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	41.9% (35.6-48.6) 54.8% (44.1-65)
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	43.8% (37.3-50.4) 58.3% (47.7-68.3)
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	54.8% (48.2-61.3) 73.8% (63.5-82)		

Moldova	Advice for home isolation <i>All patients</i> <i>COVID-19 suspected</i>	77.3% (71.6-82.2) 75.0% (67.2-81.4)		
	Advice for symptomatic treatment <i>All patients</i> <i>COVID-19 suspected</i>	88.2% (83.5-91.7) 80.7% (73.4-86.4)		
	A scheduled follow-up visit/call <i>All patients</i> <i>COVID-19 suspected</i>	82.4% (77-86.7) 73.6% (65.7-80.2)		
	Advice for family members <i>All patients</i> <i>COVID-19 suspected</i>	84.0% (78.8-88.1)	Home isolation <i>All patients</i> <i>COVID-19 suspected</i>	49.2% (42.9-55.5) 79.3% (71.8-85.2)
		98.6% (94.9-99.6)	Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	81.9% (76.6-86.3) 95.7% (91-98)
	Preventive measures for patient <i>All patients</i> <i>COVID-19 suspected</i>	95.8% (92.4-97.7)	Extra handwashing <i>All patients</i> <i>COVID-19 suspected</i>	95.4% (91.9-97.4) 94.3% (89.1-97)
		95.0% (90-97.6)	Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	95.4% (91.9-97.4) 94.3% (89.1-97)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	95.0% (91.4-97.1) 93.6% (88.2-96.6)
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	94.5% (90.9-96.8) 93.6% (88.2-96.6)
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	55.0% (48.7-61.2) 63.6% (55.3-71.1)
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	52.9% (46.6-59.2) 75.0% (67.2-81.4)		

Netherlands	Advice for home isolation			
	<i>All patients</i>	71.3%	(65.2-76.8)	
	<i>COVID-19 suspected</i>	81.5%	(74.1-87.1)	
	Advice for symptomatic treatment			
	<i>All patients</i>	49.1%	(42.7-55.6)	
	<i>COVID-19 suspected</i>	45.2%	(37-53.6)	
	A scheduled follow-up visit/call			
	<i>All patients</i>	26.1%	(20.8-32.1)	
	<i>COVID-19 suspected</i>	34.8%	(27.3-43.2)	
	Advice for family members	<i>All patients</i>	54.8%	(48.3-61.1)
<i>COVID-19 suspected</i>		63.0%	(54.6-70.6)	<i>All patients</i>
				<i>COVID-19 suspected</i>
				Social distancing
				<i>All patients</i>
				<i>COVID-19 suspected</i>
Preventive measures for patient	<i>All patients</i>	56.1%	(49.6-62.3)	Extra handwashing
	<i>COVID-19 suspected</i>	56.3%	(47.9-64.4)	<i>All patients</i>
				<i>COVID-19 suspected</i>
				Sneezing in sleeve
				<i>All patients</i>
				<i>COVID-19 suspected</i>
				Social distancing
				<i>All patients</i>
			<i>COVID-19 suspected</i>	
			Nose/mouth protection	
			<i>All patients</i>	
			<i>COVID-19 suspected</i>	
			Staying in separate room	
			<i>All patients</i>	
			<i>COVID-19 suspected</i>	
Where to find reliable information	<i>All patients</i>	24.8%	(19.7-30.7)	
	<i>COVID-19 suspected</i>	25.9%	(19.2-33.9)	
Poland*	Advice for home isolation			
	<i>All patients</i>	57.4%	(50.9-63.6)	
	<i>COVID-19 suspected</i>	61.5%		
	Advice for symptomatic treatment			
	<i>All patients</i>	73.9%	(67.9-79.2)	
	<i>COVID-19 suspected</i>	69.2%		
	A scheduled follow-up visit/call			
	<i>All patients</i>	54.3%	(47.9-60.6)	
	<i>COVID-19 suspected</i>	69.2%		
	Advice for family members	<i>All patients</i>	25.2%	(20-31.2)
<i>COVID-19 suspected</i>		84.6%	(49.6-62.3)	<i>All patients</i>
				<i>COVID-19 suspected</i>
				Social distancing
				<i>All patients</i>
				<i>COVID-19 suspected</i>
Preventive measures for patient	<i>All patients</i>	56.1%	(49.6-62.3)	Extra handwashing
				<i>All patients</i>
				<i>COVID-19 suspected</i>

	<i>COVID-19 suspected</i>	84.6%	Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	43.0% (36.8-49.5) 76.9%
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	31.7% (26.1-38) 76.9%
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	13.5% (9.7-18.5) 84.6%
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	7.8% (5-12) 38.5%
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	30.4% (24.9-36.7) 53.8%		

Romania*	Advice for home isolation <i>All patients</i> <i>COVID-19 suspected</i>	32.5% (24.6-41.5) 75.0%		
	Advice for symptomatic treatment <i>All patients</i> <i>COVID-19 suspected</i>	94.7% (89-97.6) 87.5%		
	A scheduled follow-up visit/call <i>All patients</i> <i>COVID-19 suspected</i>	41.2% (32.6-50.4) 87.5%		
	Advice for family members <i>All patients</i> <i>COVID-19 suspected</i>	35.1% (27-44.2) 87.5%	Home isolation <i>All patients</i> <i>COVID-19 suspected</i>	3.5% (1.4-8.7) 37.5%
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	34.2% (26.1-43.3) 87.5%
	Preventive measures for patient <i>All patients</i> <i>COVID-19 suspected</i>	68.4% (59.4-76.2) 100%	Extra handwashing <i>All patients</i> <i>COVID-19 suspected</i>	68.4% (59.4-76.2) 100%
			Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	64.9% (55.8-73) 87.5%
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	60.5% (51.4-69) 100%
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	57.9% (48.7-66.6) 100%
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	29.8% (22.2-38.8) 87.5%
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	4.4% (1.9-9.8) 37.5%		

Ukraine	Advice for home isolation			
	<i>All patients</i>	61.8%	(55.4-67.9)	
	<i>COVID-19 suspected</i>	85.3%	(75-91.8)	
	Advice for symptomatic treatment			
	<i>All patients</i>	83.3%	(78-87.6)	
	<i>COVID-19 suspected</i>	64.7%	(52.8-75)	
	A scheduled follow-up visit/call			
	<i>All patients</i>	64.5%	(58.1-70.4)	
	<i>COVID-19 suspected</i>	64.7%	(52.8-75)	
	Advice for family members			
<i>All patients</i>	54.4%	(47.9-60.7)	Home isolation	
<i>COVID-19 suspected</i>	88.2%	(78.5-93.9)	<i>All patients</i>	39.9%
			<i>COVID-19 suspected</i>	83.8%
			Social distancing	
			<i>All patients</i>	45.6%
			<i>COVID-19 suspected</i>	70.6%
Preventive measures for patient			Extra handwashing	
<i>All patients</i>	64.9%	(58.5-70.8)	<i>All patients</i>	57.0%
<i>COVID-19 suspected</i>	76.5%	(65.1-85)	<i>COVID-19 suspected</i>	69.1%
			Sneezing in sleeve	
			<i>All patients</i>	50.4%
			<i>COVID-19 suspected</i>	61.8%
			Social distancing	
			<i>All patients</i>	52.2%
			<i>COVID-19 suspected</i>	66.2%
			Nose/mouth protection	
			<i>All patients</i>	51.3%
			<i>COVID-19 suspected</i>	66.2%
			Staying in separate room	
			<i>All patients</i>	50.4%
			<i>COVID-19 suspected</i>	67.6%
Where to find reliable information				
<i>All patients</i>	28.9%	(23.5-35.1)		
<i>COVID-19 suspected</i>	58.8%	(47-69.7)		
United Kingdom	Advice for home isolation			
	<i>All patients</i>	50.0%	(41.2-58.8)	
	<i>COVID-19 suspected</i>	66.7%	(54.1-77.3)	
	Advice for symptomatic treatment			
	<i>All patients</i>	66.7%	(57.8-74.5)	
	<i>COVID-19 suspected</i>	68.3%	(55.8-78.7)	
	A scheduled follow-up visit/call			
	<i>All patients</i>	15.8%	(10.4-23.4)	
	<i>COVID-19 suspected</i>	21.7%	(13.1-33.6)	
	Advice for family members			
<i>All patients</i>	40.0%	(31.7-48.9)	Home isolation	
<i>COVID-19 suspected</i>	51.7%	(39.3-63.8)	<i>All patients</i>	31.7%
			<i>COVID-19 suspected</i>	40.0%
			Social distancing	
			<i>All patients</i>	14.2%
			<i>COVID-19 suspected</i>	25.0%
Preventive measures for patient			Extra handwashing	
<i>All patients</i>	28.3%	(21-37)	<i>All patients</i>	20.0%
			<i>COVID-19 suspected</i>	23.3%

	<i>COVID-19 suspected</i>	31.7% (21.3-44.2)	Sneezing in sleeve <i>All patients</i> <i>COVID-19 suspected</i>	5.8% (2.9-11.5) 5.0% (1.7-13.7)
			Social distancing <i>All patients</i> <i>COVID-19 suspected</i>	15.8% (10.4-23.4) 23.3% (14.4-35.4)
			Nose/mouth protection <i>All patients</i> <i>COVID-19 suspected</i>	5.0% (2.3-10.5) 8.3% (3.6-18)
			Staying in separate room <i>All patients</i> <i>COVID-19 suspected</i>	8.3% (4.6-14.7) 11.7% (5.8-22.2)
	Where to find reliable information <i>All patients</i> <i>COVID-19 suspected</i>	12.5% (7.7-19.6) 10% (4.7-20.1)		

* For countries with less than 20 COVID-19 suspected patients, 95% CIs were not calculated and data should be interpreted with care.