

**Supplemental Table 1: Univariate associations between patient characteristics and outcomes**

<b>A. Heard of preeclampsia before first diagnosis</b>	<i>OR</i>	<i>95% CI</i>	<i>p-value</i>
Maternal age (years)			
<25	0.40	0.25-0.66	<0.001
25-29	ref		
30-34	1.73	1.16-2.57	0.007
≥35	1.48	0.93-2.36	0.098
Year of delivery			
Before 2011	0.25	0.15-0.41	<0.001
2011-2013	0.61	0.36-1.03	0.064
2014-2016	0.66	0.41-1.06	0.088
From 2017 onwards	ref		
Highest level of education completed			
High school or less and technical/vocational school	0.33	0.20-0.55	<0.001
Some college	0.67	0.42-1.06	0.086
College	ref		
Graduate school	2.13	1.42-3.19	<0.001
Multiparity	1.39	0.78-2.67	0.262

<b>B. Serious mental/emotional impact</b>	<i>OR</i>	<i>95% CI</i>	<i>p-value</i>
Maternal age (years)			
<25	0.92	0.58-1.48	0.740
25-29	ref		
30-34	1.09	0.78-1.51	0.620
≥35	1.41	0.95-2.09	0.089
Year of delivery			
Before 2011	0.67	0.45-1.01	0.055
2011-2013	0.49	0.32-0.75	0.001
2014-2016	0.81	0.56-1.17	0.257
From 2017 onwards	ref		
Multiparity	1.10	0.70-1.75	0.677
Perinatal loss	3.97	2.36-6.68	<0.001
Cesarean section	1.09	0.82-1.45	0.565
Maternal ICU-stay	1.60	1.12-2.28	0.009
Baby admitted to the NICU	1.82	1.36-2.43	<0.001
Gestational age at delivery (weeks+days)			
<28+0	3.11	1.91-5.05	<0.001
28+0 - 31+6	1.74	1.13-2.68	0.013
32+0 - 36+6	1.30	0.94-1.80	0.112
≥37+0	ref		
Not involved in making decisions	2.20	1.58-3.07	<0.001
Not aware of preeclampsia before diagnosis	1.52	1.10-2.08	0.010
Not aware of the symptoms of preeclampsia before diagnosis	1.26	0.95-1.67	0.109
Healthcare provider did not convey the seriousness of the condition	1.50	1.11-2.02	0.008

**C. Family planning: wanted more children but decided not to have another pregnancy**

	OR	95% CI	p-value
Maternal age (years)			
<25	0.55	0.29-1.06	0.075
25-29	ref		
30-34	0.97	0.65-1.44	0.868
≥35	1.63	1.05-2.53	0.031
Multiparity	2.14	1.31-3.49	0.002
Gestational age at delivery (weeks+days)			
<28+0	1.08	0.62-1.90	0.784
28+0 - 31+6	1.45	0.88-2.39	0.148
32+0 - 36+6	1.26	0.85-1.86	0.247
≥37+0	ref		
Perinatal loss	0.32	0.15-0.68	0.003
No counseling about the risk of preeclampsia in future pregnancies	1.20	0.85-1.69	0.296
No counseling about later-life health risks	1.27	0.82-1.97	0.286
Maternal ICU-stay	1.24	0.83-1.86	0.293
Baby admitted to the NICU	1.34	0.95-1.90	0.094

Results are from univariate logistic regression analysis. OR=Odds Ratio; 95% CI=95% confidence interval; ICU=intensive care unit; NICU=neonatal intensive care unit