

Appendix 1

Comparison of characteristics of participants^a versus non-participants within the eligible intention-to-screen population of the D₂AF study.

Characteristic	Participants (n=4339)	Non-participants (n=5061)	p-value
Female, n (%)	2336 (53.8)	2831 (55.9)	0.041
Age in years, M (SD)	73.5 (5.5)	76.7 (7.4)	<0.001
History ^b			
Hypertension, n (%)	2212 (51.1)	2416 (48.3)	0.008
Stroke/TIA, n (%)	329 (7.6)	603 (12.1)	<0.001
Diabetes, n (%)	783 (18.1)	1029 (20.6)	0.002
Heart failure, n (%)	80 (1.8)	304 (6.1)	<0.001
Thromboembolism, n (%)	200 (4.8)	271 (5.4)	0.077
Vascular disease, n (%)	644 (14.9)	968 (19.4)	<0.001

Abbreviations: M (mean), TIA (transient ischemic attack), SD (standard deviation).

^a In the current diagnostic accuracy study, we analyse 4339 patients whereas we included 4106 patients in the intention-to-screen arm of the D₂AF randomized controlled trial. The screening of 233 patients occurred after the end of the study year, and they were therefore not eligible for the randomized controlled trial. However, we did include them in the diagnostic accuracy study.

^b For nine participants and 64 non-participants, history was missing.