

List of Supplementary tables

Supplementary table 1: Health post and child factors associated with referrals of 0 – 59 months sick young children in four regions of Ethiopia

| Characteristics | Referral status of 0 – 59 days old sick young infants | | | Referral status of 2 – 59 months old children | | |
|---|---|--------------------------------|---------|---|---------------------------------|---------|
| | Referred N=39 n (%) | Not referred N=190 n (%) | P-value | Referred N=78 n (%) | Not referred N=1045 n (%) | P-value |
| <i>The health post</i> | | | | | | |
| Mean score of basic amenities | | | | | | |
| 0 to 33 percent | 13 (24) | 42 (76) | 0.466 | 20 (13) | 129 (87) | 0.043 |
| 34 to 66 percent | 9 (15) | 50 (85) | | 19 (6) | 292 (94) | |
| 67 to 100 percent | 17 (15) | 98 (85) | | 39 (6) | 624 (94) | |
| Mean score of basic equipment | | | | | | |
| 0 to 33 percent | 2 (22) | 7 (78) | 0.825 | 3 (4) | 73 (96) | 0.512 |
| 34 to 66 percent | 10 (19) | 43 (81) | | 12 (8) | 147 (92) | |
| 67 to 100 percent | 27 (16) | 140 (84) | | 63 (7) | 825 (93) | |
| Mean score of standard precaution | | | | | | |
| 0 to 33 percent | 4 (9) | 40 (91) | 0.527 | 30 (8) | 353 (92) | 0.279 |
| 34 to 66 percent | 14 (19) | 59 (81) | | 31 (8) | 355 (92) | |
| 67 to 100 percent | 21 (19) | 91 (81) | | 17 (5) | 337 (95) | |
| Rapid diagnostic test (RDT) | | | | | | |
| 0 to 33 percent | 10 (17) | 50 (83) | 0.947 | 47 (9) | 467 (91) | 0.026 |
| 34 to 66 percent | 0 | 0 | | 0 | 0 | |
| 67 to 100 percent | 29 (17) | 140 (83) | | 31 (5) | 578 (95) | |
| Mean score of essential medicines | | | | | | |
| 0 to 33 percent | 1 (8) | 11 (92) | 0.026 | 17 (18) | 77 (82) | 0.001 |
| 34 to 66 percent | 32 (21) | 113 (79) | | 43 (6) | 697 (94) | |
| 67 to 100 percent | 6 (8) | 66 (92) | | 18 (6) | 271 (94) | |
| Health post's overall service readiness index | | | | | | |
| 0 to 33 percent | 0 (0) | 1 (0) | 0.737 | 7 (18) | 31 (82) | 0.193 |
| 34 to 66 percent | 22 (19) | 93 (81) | | 43 (7) | 575 (93) | |
| 67 to 100 percent | 17 (15) | 96 (85) | | 28 (6) | 439 (94) | |
| Number of health extension worker per health post | | | | | | |

| | | | | | | | |
|--|---|---------|----------|-------|---------|-----------|-------|
| | 1 | 2 (5) | 39 (95) | | 20 (9) | 196 (81) | |
| | 2 or more | 37 (20) | 151 (80) | 0.024 | 58 (6) | 849 (94) | 0.264 |
| <i>Distance between health post and the nearest referral health centre</i> | | | | | | | |
| Distance to the next referral site ^b | | | | | | | |
| | 0 - 8 kms | 16 (21) | 61 (79) | | | | |
| | 8.01 - 20 kms | 10 (12) | 71 (88) | 0.554 | | | |
| | >20 kms | 13 (19) | 57 (81) | | | | |
| | 0 - 8 kms | | | | 24 (7) | 340 (93) | |
| | 8.01 - 14 kms | | | | 25 (8) | 301 (92) | 0.699 |
| | >14 kms | | | | 22 (6) | 362 (94) | |
| <i>The child</i> | | | | | | | |
| Region | | | | | | | |
| | Amhara | 3 (25) | 9 (75) | | 34 (7) | 428 (93) | |
| | Oromia | 10 (13) | 65 (87) | 0.722 | 26 (7) | 342 (93) | 0.691 |
| | SNNPR ^a | 9 (21) | 33 (79) | | 11 (8) | 128 (92) | |
| | Tigray | 17 (17) | 83 (83) | | 7 (5) | 147 (95) | |
| Age | | | | | | | |
| | 1 st week | 14 (24) | 45 (76) | | | | |
| | 2 - 4 weeks | 16 (16) | 82 (84) | 0.353 | | | |
| | 5 - 8 weeks | 9 (13) | 63 (87) | | | | |
| | 2 - 11 months | | | | 26 (8) | 294 (92) | |
| | 12 - 23 months | | | | 28 (8) | 325 (92) | 0.249 |
| | 24 - 59 months | | | | 24 (5) | 426 (95) | |
| Sex | | | | | | | |
| | Boy | 18 (15) | 106 (85) | 0.395 | 40 (7) | 550 (93) | 0.810 |
| | Girl | 21 (20) | 84 (80) | | 38 (7) | 495 (93) | |
| Disease classification | | | | | | | |
| | Non-severe illnesses not requiring referral | 21 (12) | 159 (88) | 0.013 | 56 (5) | 1023 (95) | 0.001 |
| | Severe illnesses requiring referral | 18 (37) | 31 (63) | | 22 (50) | 22 (50) | |

^a SNNPR - Southern Nations, Nationalities, and Peoples Region

^b GIS based data of distance from health post to health centre were missing for 0-59 days sick young infant (1) and 2-59 months sick children (49).

Supplementary table 2: Individual health post readiness factors associated with referrals of 0-59 months sick children in four regions of Ethiopia

| Characteristics | Referral status of 0 – 59 days sick young infants | | | Referral status of 2 – 59 months old children | | |
|---|---|--------------------------------|---------|---|-----------------------------|---------|
| | Referred N=39 n (%) | Not referred N=190 n (%) | P-value | Referred N=78 n (%) | Referred N=1045 n (%) | P-value |
| <i>Availability of basic amenities</i> | | | | | | |
| Water source | Not available | 15 (22) | 0.296 | 30 (8) | 356 (92) | 0.558 |
| | Available | 24 (15) | | 48 (7) | 689 (93) | |
| Electric power | Not available | 18 (17) | 0.886 | 51 (8) | 602 (92) | 0.257 |
| | Available | 21 (18) | | 27 (6) | 443 (94) | |
| Facility owned toilet | Not available | 6 (18) | 0.880 | 14 (7) | 207 (93) | 0.981 |
| | Available | 33 (17) | | 64 (7) | 838 (93) | |
| Facility owned landline or mobile phone | Not available | 35 (19) | 0.133 | 51 (6) | 828 (94) | 0.038 |
| | Available | 4 (10) | | 27 (11) | 217 (89) | |
| <i>Availability of basic equipment</i> | | | | | | |
| Thermometer | Not available | 8 (14) | 0.594 | 12 (5) | 228 (95) | 0.207 |
| | Available | 31 (18) | | 66 (8) | 817 (92) | |
| Child weight scale | Not available | 5 (13) | 0.526 | 20 (9) | 214 (91) | 0.388 |
| | Available | 34 (18) | | 58 (7) | 831 (93) | |
| Stethoscope | Not available | 18 (19) | 0.709 | 35 (8) | 399 (92) | 0.336 |
| | Available | 21 (16) | | 43 (6) | 646 (94) | |
| MUAC tape measure | Not available | 1 (100) | 0.040 | 0 (0) | 14 (100) | 0.701 |
| | Available | 38 (17) | | 78 (7) | 1031 (93) | |
| <i>Availability diagnostics</i> | | | | | | |
| Rapid diagnostic test (RDT) | Not available | 10 (17) | 0.947 | 47 (9) | 467 (91) | 0.026 |
| | Available | 29 (17) | | 31 (5) | 578 (95) | |

| <i>Availability of essential medicines</i> | | | | | | | |
|--|---------------|---------|----------|-------|---------|----------|-------|
| Vitamin A | Not available | 9 (27) | 25 (73) | 0.211 | 11 (5) | 207 (95) | 0.214 |
| | Available | 30 (15) | 165 (85) | | 67 (7) | 838 (93) | |
| Amoxicillin | Not available | 6 (14) | 36 (86) | 0.745 | 28 (15) | 159 (85) | 0.001 |
| | Available | 33 (18) | 154 (82) | | 50 (5) | 886 (95) | |
| Gentamycin | Not available | 20 (14) | 127 (86) | 0.201 | | | |
| | Available | 19 (23) | 63 (77) | | | | |
| Tetracycline eye ointment | Not available | 26 (19) | 110 (81) | 0.457 | 46 (8) | 567 (92) | 0.554 |
| | Available | 13 (14) | 80 (86) | | 32 (6) | 478 (94) | |
| Oral rehydration solution (ORS) | Not available | 2 (13) | 13 (87) | 0.627 | 10 (17) | 48 (83) | 0.048 |
| | Available | 37 (17) | 177 (83) | | 68 (6) | 997 (94) | |
| Zinc-ORS | Not available | | | | 69 (7) | 947 (93) | 0.608 |
| | Available | | | | 9 (8) | 98 (92) | |
| Zinc | Not available | | | | 16 (9) | 152 (91) | 0.331 |
| | Available | | | | 62 (7) | 893 (93) | |
| Coartem | Not available | 18 (20) | 71 (80) | 0.494 | 47 (9) | 507 (91) | 0.087 |
| | Available | 21 (15) | 119 (85) | | 31 (5) | 535 (95) | |
| Plumpy Nut | Not available | 8 (23) | 27 (77) | 0.404 | 30 (8) | 372 (92) | 0.688 |
| | Available | 31 (16) | 163 (84) | | 48 (7) | 673 (93) | |
| BP100 | Not available | 38 (17) | 180 (83) | 0.012 | 77 (7) | 993 (93) | 0.107 |
| | Available | 1 (9) | 10 (91) | | 1 (2) | 52 (98) | |