

TABLE S2: Diagnostic properties of CoMiSS for various cut-off values**A. Baseline CoMiSS (n=250)**

CoMiSS cut-off	Sensitivity	Specificity	PPV	NPV
≥ 5	88.5%	33.3%	89.7%	30.6%
≥ 6	78.8%	51.5%	91.4%	27.0%
≥ 7	68.2%	57.6%	91.4%	21.6%
≥ 8	56.2%	60.6%	90.4%	17.4%
≥ 9	43.8%	69.7%	90.5%	15.9%
≥ 10	35.9%	69.7%	88.6%	14.2%
≥ 11	25.8%	81.8%	90.3%	14.4%
≥ 12	20.3%	87.9%	91.7%	14.4%

B. Baseline CoMiSS plus ≥50% reduction from baseline to Visit 2 (n=247)

CoMiSS cut-off	Sensitivity	Specificity	PPV	NPV
≥ 5	38.1%	62.5%	87.2%	13.1%
≥ 6	35.3%	68.8%	88.4%	13.7%
≥ 7	34.4%	71.9%	89.2%	14.0%
≥ 8	30.2%	71.9%	87.8%	13.3%
≥ 9	25.6%	78.1%	88.7%	13.5%
≥ 10	20.5%	78.1%	86.3%	12.8%
≥ 11	16.3%	84.4%	87.5%	13.0%
≥ 12	14.0%	87.5%	88.2%	13.1%