## **TRUFFLE 2 flowchart**

32+0 to 36+6 weeks, singleton, non-anomalous pregnancy, age ≥ 18 AND EFW or AC <10<sup>th</sup> centile **OR** decrease by 50 percentiles AND Normal computerised CTG (STV ≥ 4.5msec) AND **Umbilical Cerebral Ratio** ≥ 1.0 at 32+0 - 33+6 weeks ≥ 0.8 at 34+0 - 36+6 weeks (UCR must be repeated within 15 minutes – 24 hours to confirm abnormal reading) AND No contra-indication to either trial arm RANDOMISATION (at time of 2nd abnormal UCR) 1:1 central by centre and gestational age (above or below 34 weeks) **IMMEDIATE DELIVERY DELAYED DELIVERY** Twice weekly ultrasound Doppler and Delivery or start of induction of labour within cCTG 48 hours of randomisation Delivery mandatory if: cCTG STV < 4.5 msec To allow for administration of repeated decelerations corticosteroids and magnesium OR if UA EDF is reversed sulphate (see guidance) ≥ 34 + 0 if UA EDF absent **PRIMARY OUTCOME** – Composite perinatal outcome **SECONDARY OUTCOMES –** 2 year neonatal follow-up & maternal outcomes

TRUFFLE-2\_Protocol\_4.8\_02-12-2022