

Supplemental Table 3: Crude associations between sub-types of LUTS and the information collection component of JST-IC

Sub-types of LUTS	JST-IC (information collection)					
	Total	p	Males	p	Females	p
Frequency	No	3.4±0.9	0.21	3.4±0.8	0.74	3.5±0.9
	Yes	3.3±0.9		3.3±1.0		3.3±1.0
Nocturia (Once or more)	No	3.5±0.8	0.18	3.4±0.8	0.57	3.5±0.8
	Yes	3.3±0.9		3.3±0.9		3.3±1.0
Nocturia (Twice or more)	No	3.5±0.8	<0.01	3.5±0.8	<0.01	3.5±0.8
	Yes	3.2±1.0		3.3±0.9		3.2±1.1
Nocturia (Three times or more)	No	3.4±0.8	<0.01	3.4±0.8	<0.01	3.4±0.9
	Yes	3.0±1.1		3.1±1.1		2.8±1.3
Urgency	No	3.4±0.9	0.02	3.4±0.8	0.15	3.4±0.9
	Yes	3.1±1.1		3.2±1.1		3.1±1.1
Urinary incontinence	No	3.4±0.9	<0.01	3.4±0.9	0.14	3.4±0.9
	Yes	2.9±1.2		3.1±1.1		2.8±1.2
OAB	No	3.4±0.9	0.02	3.4±0.8	0.13	3.4±0.9
	Yes	3.1±1.1		3.2±1.1		3.0±1.1

LUTS: lower urinary tract symptom.

JST-IC: Japan Science and Technology Agency Index of Competence.

Student's t-tests were performed.